



Summary Report on Adverse Events Following Immunisation of COVID-19 Vaccines in Malaysia

(Data as of 20th September 2022)¹⁻³

#5

Introduction

The COVID-19 vaccines have been proven to be highly effective in preventing severe illnesses and deaths from COVID-19 infection.⁴ In an effort to stall the spread of the infection and end the pandemic, a mass vaccination roll-out was initiated on 24th February 2021 under the **National Immunisation Program for COVID-19 (PICK)** in Malaysia.

Similar to any medicines, vaccines may cause side-effects commonly known as adverse event following immunisation (AEFIs). An AEFI is defined as **any untoward medical occurrence that follows the administration of a vaccine and may not necessarily be causally related to the vaccine itself.**⁵

It is inaccurate to assume that all AEFIs reported in this summary report are directly caused by the vaccine.

When millions of vaccines are administered in a short period of time, it is expected to see a surge in the number of adverse events being reported including serious ones. However, reviews of these individual reports often reveal that vaccine does not play a role in the vast majority of these events.

The reported adverse events can also occur due to **fear of injections or the immunisation process, previously undiagnosed illnesses, underlying diseases, or medications being taken concurrently by the vaccine recipients. These events may also happen coincidentally, shortly after a vaccine was administered.**

How does NPRA collect adverse event reports?

Similar to other regulatory agencies, NPRA monitors drugs and vaccines safety through passive surveillance. AEFI reports are collected via the existing **NPRA Adverse Drug Reaction (ADR)/AEFI Reporting System.**

In addition to the current reporting system, during PICK, **notifications of documented minor AEFIs** from the vaccine recipients are also collected through the **MySejahtera Application** in their smartphones.

Documented AEFI are known adverse events that already described in the product leaflet and are based on global safety data from clinical trials and/or post-marketing safety monitoring.

NPRA receives all medicines adverse event reports including AEFIs from pharmaceutical companies, healthcare professionals in government and private health institutions as well as consumers in Malaysia.

NPRA has issued the Malaysian ADR/AEFI Reporting Manual for Healthcare Providers to guide reporters on how to report adverse events.⁵ NPRA also continuously highlights the importance of ADR/AEFI reporting and how to report them through periodic communications, several trainings, and awareness campaigns.

How does NPRA process AEFI reports and monitor safety of COVID-19 vaccines?

As with any medicine and vaccine, NPRA closely monitors the safety of COVID-19 vaccines. Every adverse event report received at the national centre is carefully processed and assessed by trained pharmacists. When clinically important information is missing from a report, utmost efforts will be made to obtain additional information.

NPRA requires healthcare providers to report all suspected AEFIs including any death after COVID-19 vaccination, **even if it is unclear whether the vaccine was the cause**. AEFIs are initially categorised into **serious** and **non-serious**. **Serious** AEFIs include those that require hospitalisation, prolonged existing hospitalisation, are life-threatening, cause persistent or significant disability/incapacity, a congenital anomaly/birth defect, or suspected to cause death. **While these events may happen after vaccination, they are rarely caused by the vaccine**. Serious AEFIs are investigated thoroughly by the healthcare facility involved. The investigation report will then be reviewed by a committee of experts, the **COVID-19 Vaccines Pharmacovigilance Special Committee (JFK)** to determine if the events are causally linked to the vaccine.

All reports will be subsequently presented to the **Malaysian Adverse Drug Reactions Advisory Committee (MADRAC)** before they are submitted to the **World Health Organisation (WHO) global database**.

Reports recorded in the databases are constantly reviewed and monitored to identify unexpected adverse events or potential safety **signals** for further evaluation. In addition to monitoring local AEFI reports, NPRA also collaborates with Product Registration Holders to monitor and detect any emerging safety issues. NPRA also network with other regulatory agencies to keep abreast with new safety concerns raised globally. This allows rapid detection and assessment of all available safety information on the vaccines to ensure the overall benefit–risk profile of the vaccines remains positive. Any emerging risks of the vaccines will be communicated promptly to healthcare professionals and the public so that suitable actions can be taken accordingly.

*Reports help to identify **signals** that **alert** scientists of possible cause-and-effect relationships that **need to be investigated**.*

Total Doses of COVID-19 Vaccines Administered¹⁻²


In Malaysia, there are currently eight (8) types of COVID-19 vaccines conditionally registered with the Drug Control Authority (DCA). These vaccines have met NPRA's high standards for quality, safety and efficacy —namely Comirnaty (Pfizer), CoronaVac (Sinovac), Vaxzevria (AstraZeneca), Convidecia (Cansino), Covilo (Sinopharm), Spikevax (Moderna), Jcovden (Jassen), and Covaxin (Bharat).

Till date, only Comirnaty, CoronaVac, Vaxzevria, Convidecia, and Covilo vaccines are being used in the National COVID-19 Immunisation Programme (PICK). As of 20th September 2022, a total of 72,091,311 COVID-19 vaccine doses have been administered.

Table 1: Total COVID-19 vaccine doses administered by product (as of 20th September 2022)

Comirnaty (Pfizer)	44,605,452 doses
CoronaVac (Sinovac)	21,519,759 doses
Vaxzevria (AstraZeneca)	5,696,251 doses
Convidecia (Cansino)	227,572 doses
Covilo (Sinopharm)	42,277 doses

Rate of Reported Adverse Events²⁻³



AEFI responses received via MySejahtera Application

From the start of the vaccine roll-out up to 20th September 2022, the total of minor and documented AEFIs recorded via the MySejahtera Application is 978,549 which is equivalent to 13,574 responses* for every 1,000,000 doses of vaccines administered.

The most common side effects notified were consistent with those typically observed following vaccination. These include injection site pain, headache, fatigue, muscle or joint pain, lethargy and fever, which will usually recover in a few days with or without treatment. The reporting trend of these common side-effects were also consistent throughout the monitoring period.

*Important note: *The total number of responses collected may not accurately represent the total number of individuals who experience an adverse event, as each individual may report more than one response.*

AEFI reports received via NPRA Reporting System

While through the existing NPRA Reporting System, the total of AEFI reports received was 26,613 which is equivalent to **370 reports per 1,000,000 doses** administered. The majority of reports received, at **93%, were non-serious**, short-term, and do not pose any potential risk to the health of the vaccine recipients.

Serious adverse events after COVID-19 vaccination occurred but very rarely. During this monitoring period, only 1,865 or 7% of the total AEFI reports received were categorised as serious AEFIs. Note that the **ACTUAL proportion of serious AEFIs could be much smaller**, when the relatively high number of minor AEFI responses received via the MySejahtera Application are taken into account. **The reporting rate of serious AEFIs was recorded at 0.0026% of total doses administered, or 26 reports per 1,000,000 doses.**




Figure 1: Number of AEFI reports received via NPRA Reporting System

It should be noted that the number of reports received for different COVID-19 vaccines are not directly comparable as the total number of doses used for each vaccine differs as they were rolled out at different times during the vaccination programme.

The **most frequently reported serious AEFIs** were shortness of breath, chest pain, palpitations and anaphylactic reaction. The majority of these serious cases require only **short-term hospitalisation for treatment or further observation and at the time of AEFI reporting, the affected recipients have mostly recovered or in the state of recovering from the reported event.**

It should be stressed that the causal links of the event to the vaccination in these reports have not been ascertained meaning that the vaccines do not necessarily cause the serious events, as explained in the [Introduction Section](#).

Out of **1,865 serious AEFI reports received**, 612 were reports for death cases. However, after JFK's evaluation based on the investigation findings up to 20th September 2022, none of the fatal outcomes reported among the vaccine recipients could be directly linked to the vaccine administered.

Booster doses (PICK-B)

The Comirnaty, CoronaVac and Vaxzevria vaccines have been approved for use as boosters in adults under the PICK-Booster (PICK-B) which began on 13th October 2021. It is not expected that the type of side effects will be different to first and second vaccine doses based on the results of clinical trials and observations in other countries where booster doses have been rolled out earlier.⁶⁻¹²

As of 20th September 2022, a total of 16,704,848 booster doses of the Comirnaty, CoronaVac and Vaxzevria vaccines have been administered. Till date, NPRA has received 1,717 reports (103 reports per 1,000,000 doses administered) following the use of boosters, of which 162 of reports (**10 reports per 1,000,000 doses administered, or 9.4% of total AEFI reports received**) are categorised as **serious**. This is **lower** than the AEFI reporting rate for COVID-19 vaccines for all vaccine doses combined.

The types of reactions experienced after a booster dose were also found to be similar to those experienced after primary doses, including in heterologous* vaccination. The most common side effects reported include fever, headache, acute stress reaction, injection site pain and muscle pain, dizziness, and shortness of breath.

Review of booster dose AEFI reports at this point of time does not raise any new safety concerns. NPRA will continue to closely monitor AEFI data following the use of boosters.

**Heterologous booster refers to the administration of a vaccine product that differs from the product(s) previously used for primary vaccine series (e.g. a vectored vaccine followed by an mRNA vaccine), once an initially sufficient immune response rate in a vaccinated population has waned over time.¹³*




Figure 2: AEFI reporting rate per 1,000,000 doses (all doses versus booster doses)

COVID-19 vaccination in children aged 5 to 11 years

The National Covid-19 Immunisation Programme for children aged 5 to 11 years (PICKids) in Malaysia has been launched on the 3rd February 2022 following the conditional approval of Comirnaty 10 mcg Concentrate for Dispersion for Injection (Pfizer-BioNTech). This special COVID-19 vaccine formulation has a lower dose compared to those for aged 12 years and above (30 mcg). CoronaVac Injection Suspension COVID-19 Vaccine (Vero Cell), Inactivated, the second COVID-19 vaccine conditionally approved for children aged 5 to 11 years in Malaysia, has been made available to PICKids on 7th March 2022.

The NPRA is closely monitoring the AEFI reported in these 5 to 11-years-old children. As of 20th September 2022, a total of 3,306,118 doses of the Comirnaty and CoronaVac vaccines have been administered to children aged 5 to 11 years. There have been 522 AEFI reports received for this age group, equating to a rate of **158 reports per 1,000,000 doses administered**. The AEFI reporting rate for children aged 5 to 11 years is **lower than the overall AEFI reporting rate** thus far (370 for per 1,000,000 doses administered).

Of these, the **vast majority (94%)** were non-serious effects. The most frequently reported side effects for PICKids were fever, immunisation stress-related response (ISSR), skin itchiness or redness, dizziness, and headache. From the total of 522 AEFI reports received for children aged 5 to 11 years, there were 34 reports involving serious AEFIs such as exacerbation of asthma and Bell's palsy.

The AEFI reporting rate among vaccine recipients aged 5 to 11 years old in Malaysia (158 reports per 1,000,000 doses) is comparable to global scenario, for instances, in Canada (242 reports per 1,000,000 doses) and in Australia (709 reports per 1,000,000 doses). These countries also reported lower AEFI reporting rates among children aged 5 to 11 years compared to adult vaccine recipients.⁸⁻⁹ Similarly,

the most frequently reported adverse effects were mild side effects, such as pain, swelling, injection site itchiness or redness, headache and fatigue.

Overall, the trend of AEFI reported to NPRA among children aged 5 to 11 years is similar to what is being reported by the reference regulatory agencies. To date, no safety issues have been identified locally and globally following COVID-19 vaccine use in this age group.^{7-9,14-16} NPRA will continue to monitor the safety of COVID-19 vaccines used in children aged 5 to 11 years, including for the CoronaVac vaccine.

Adverse Events of Special Interest

NPRA is also closely monitoring the occurrences and outcome of specific adverse events known as ‘adverse events of special interest’, including anaphylaxis, acute facial paralysis, myocarditis/pericarditis, and vaccine-induced immune thrombocytopenia and thrombosis (VITT). These events have mainly been reported in the vaccines clinical trials and post-marketing surveillance globally or observed previously with the use of other vaccines.⁶⁻¹²

Anaphylaxis is a well-known adverse reaction and the most commonly reported serious AEFI associated with vaccines in general. It usually occurs 15-30 minutes after vaccination. Acknowledging this risk, as part of the risk management plan, PICK requires all vaccine recipients to be observed for 15-30 minutes post vaccination at the vaccination centres. In rare instances should anaphylaxis reaction occurs, the affected recipient can be treated immediately and effectively.¹⁷

- During the monitoring period, NPRA had received a total of 108 AEFI reports for anaphylaxis (Brighton level 1-3) with COVID-19 vaccines equivalent to 1.5 reports per million doses administered. Comirnaty, the mRNA vaccines recorded the highest number of cases at 72 reports or 1.6 per million doses administered. This locally observed rate of anaphylaxis is similar to those reported overseas.⁶⁻¹⁰ All anaphylaxis cases were treated accordingly and have recovered or were recovering at the time of reporting.

Acute facial paralysis, also known as Bell’s palsy, is a condition that causes temporary weakness on one side of the face. It typically develops gradually and most people recover within a few months. Acute facial paralysis can strike at any age and has been linked to a number of infectious diseases, including the SARS-CoV-2 virus.^{6,18} Acute facial paralysis has been reported in clinical trials of mRNA vaccines, including Comirnaty vaccine. Recent findings also showed an overall increased risk of Bell’s palsy after immunisation with CoronaVac, the inactivated virus vaccine.¹⁹

- NPRA had received a total of 157 AEFI reports of acute facial paralysis associated with COVID-19 vaccines during the monitoring period, equating to 2.2 reports per million doses administered. Comirnaty vaccine was associated with 90 reports or 2 per million doses administered, CoronaVac vaccine with 47 reports or 2.2 per million doses administered, and AstraZeneca vaccine with 20 reports or 3.5 per million doses administered. All acute facial paralysis cases have recovered or were recovering at the time of reporting.

Myocarditis/Pericarditis is a known but very rare side effect of mRNA vaccines e.g. Comirnaty, especially in male adolescents and young adults.^{6-9,20} Symptoms typically include palpitation, chest pain, arrhythmia and dyspnoea. It is usually mild and temporary, with most people getting better within a few days with minimal treatment.²¹

- As of 20th September 2022, NPRA has received 63 reports which have been assessed as likely to be myocarditis/pericarditis following about 44.6 million doses of the Comirnaty vaccine. This is equivalent to 1.4 reported myocarditis/pericarditis cases per million doses of Comirnaty vaccine. There has also been one (1) report of likely myocarditis/pericarditis associated with AstraZeneca vaccine, equating to 0.2 per million doses. Myocarditis/pericarditis is found to be more common in adolescents (0.7 cases per million doses). Most of the cases were mild in nature and the vaccine recipients responded well to the treatment and have recovered or were recovering at the time of reporting.

Vaccine-induced immune thrombocytopenia and thrombosis (VITT), which may also be called thrombotic thrombocytopenia syndrome (TTS) following COVID-19 vaccination, is another known but very rare serious side effect of adenoviral vector COVID-19 vaccines such as AstraZeneca vaccine. The exact mechanism of how VITT is triggered is still under investigation, however the majority of cases are associated with the finding of thrombosis (frequently in uncommon locations, such as cerebral venous sinus or splanchnic veins), low platelet count, markedly elevated D-dimer and positive anti-platelet factor 4 (anti-PF4) antibodies.²² Individuals with VITT generally present symptoms between 4 and 42 days (most commonly 4-30 days) after vaccination with an adenoviral vector vaccine, which include persistent headaches which do not respond to simple pain killers, blurred vision, difficulty with speech, seizures, and bleeding or bruising.

- Up to 20th September 2022, four (4) cases of VITT were reported following about 5.7 million doses of AstraZeneca vaccine and one (1) case following about 44.6 million doses of Comirnaty vaccine.

Summary

Similar to global scenario⁶⁻¹², **the vast majority (93%)** of the reported AEFIs in Malaysia are **non-serious**. The most common reactions included injection site pain, headache, fatigue, muscle or joint pain, lethargy and fever.



The rate of **serious** AEFIs reported via the NPRA Reporting System is **small at 26 reports per 1,000,000 doses**, most requiring **short-term hospitalisation for observation and treatment**. However, following detailed investigations and reviews, these events are not directly caused by the vaccines given.

The **benefit-to-risk-ratio** of COVID-19 vaccines registered in Malaysia **remains very favourable**.

NPRA is continuously monitoring the safety of COVID-19 vaccines in the Malaysian population. We encourage all healthcare professionals and vaccine recipients to report any suspected AEFIs. This provides valuable data that helps us to identify new risks and appropriate safety measures and regulatory actions can be taken to mitigate the risks.

References

1. Ministry of Health, Malaysia. COVIDNOW in Malaysia [Internet]. 2022 [cited 2022 Sep 20]. Available from: <https://covidnow.moh.gov.my/>
2. Ministry of Health, Malaysia. COVID19-public [Internet]. Github. 2022 [cited 2022 Jun 11]. Available from: <https://github.com/MoH-Malaysia/covid19-public/>
3. National Pharmaceutical Regulatory Agency (NPRA). The Malaysian National ADR database [Internet]. 2022 [cited 2022 Sep 20]. Available from: <https://www.npra.gov.my> (access restricted)
4. World Health Organization (WHO). Coronavirus disease (COVID-19): Vaccines [Internet]. 2022 May 17 [cited 2022 Sep 20]. Available from: [https://www.who.int/news-room/questions-and-answers/item/coronavirus-disease-\(covid-19\)-vaccines](https://www.who.int/news-room/questions-and-answers/item/coronavirus-disease-(covid-19)-vaccines)
5. National Pharmaceutical Regulatory Agency (NPRA). The Malaysian Adverse Drug Reaction (ADR)/Adverse Event Following Immunisation (AEFI) Reporting Manual for Healthcare Providers [Internet]. 2022 [cited 2021 Sep 20]. Available from: <https://www.npra.gov.my/easyarticles/images/users/1047/Adverse-Drug-Reaction-ADR--Adverse-Event-Following-Immunisation-AEFI-Reporting-Manual-For-Healthcare-Providers.pdf>
6. Medicines and Healthcare Products Regulatory Agency (MHRA). Coronavirus vaccine - weekly summary of Yellow Card reporting [Internet]. 2022 Oct 7 [cited 2022 Oct 10]. Available from: <https://www.gov.uk/government/publications/coronavirus-covid-19-vaccine-adverse-reactions/coronavirus-vaccine-summary-of-yellow-card-reporting>
7. Health Sciences Authority. Safety updates on COVID-19 vaccines [Internet]. 2022 Sep 19 [cited 2022 Sep 20]. Available from: <https://www.hsa.gov.sg/COVID19-vaccines-safety-updates>
8. Ontario Agency for Health Protection and Promotion (Public Health Ontario). Adverse Events Following Immunization (AEFIs) for COVID-19 in Ontario: December 13, 2020 to October 9, 2022. [Internet]. 2022 [cited 2022 Oct 10]. Available from: https://www.publichealthontario.ca/-/media/documents/ncov/epi/covid-19-aefi-report.pdf?sc_lang=en
9. Therapeutic Goods Administration. COVID-19 vaccine weekly safety report 09-06-2022 [Internet]. 2022 Oct 6 [cited 2022 Oct 10]. Available from: <https://www.tga.gov.au/periodic/covid-19-vaccine-safety-report-09-06-2022>
10. Uppsala Monitoring Centre (UMC). COVID-19 vaccine reporting in VigiBase. Report 20, data extraction date: 2022-09-25 [Internet]. 2022 Sep 28 [cited 2022 Oct 10] Available from: <https://www.vigilyze.who-umc.org> (access restricted)
11. Hause AM, Baggs J, Gee J, et al. Safety Monitoring of an Additional Dose of COVID-19 Vaccine — United States, August 12–September 19, 2021. MMWR Morb Mortal Wkly Rep 2021;70:1379–1384. DOI: <http://dx.doi.org/10.15585/mmwr.mm7039e4>
12. Munro APS, Janani L, Cornelius V, Aley PK, Babbage G, Baxter D, et al. Safety and immunogenicity of seven COVID-19 vaccines as a third dose (booster) following two doses of ChAdOx1 nCov-19 or BNT162b2 in the UK (COV-BOOST): a blinded, multicentre, randomised, controlled, phase 2 trial. Lancet. 2021 Dec 18;398(10318):2258–76. Available from: [https://doi.org/10.1016/S0140-6736\(21\)02717-3](https://doi.org/10.1016/S0140-6736(21)02717-3)

13. World Health Organization (WHO). Interim recommendations for heterologous COVID-19 vaccine schedules [Internet]. 2021 Dec 16 [cited 2022 Jun 11]. Available from: <https://www.who.int/publications/i/item/WHO-2019-nCoV-vaccines-SAGE-recommendation-heterologous-schedules>
14. Han B, Song Y, Li C, Yang W, Ma Q, Jiang Z, Li M, Lian X, Jiao W, Wang L, Shu Q, Wu Z, Zhao Y, Li Q, Gao Q. Safety, tolerability, and immunogenicity of an inactivated SARS-CoV-2 vaccine (CoronaVac) in healthy children and adolescents: a double-blind, randomised, controlled, phase 1/2 clinical trial. *Lancet Infect Dis*. 2021 Dec;21(12):1645-1653. Available from: [https://dx.doi.org/10.1016%2FS1473-3099\(21\)00319-4](https://dx.doi.org/10.1016%2FS1473-3099(21)00319-4)
15. Wu Q, Dudley MZ, Chen X, Bai X, Dong K, Zhuang T, Salmon D, Yu H. Evaluation of the safety profile of COVID-19 vaccines: a rapid review. *BMC Med*. 2021 Jul 28;19(1):173. Available from: <https://dx.doi.org/10.1186%2Fs12916-021-02059-5>
16. Walter EB, Talaat KR, Sabharwal C, Gurtman A, Lockhart S, Paulsen GC, et al. Evaluation of the BNT162b2 Covid-19 Vaccine in Children 5 to 11 Years of Age. *N Engl J Med*. 2021;386(1):35-46. Available from: <https://doi.org/10.1056/nejmoa2116298>
17. Ministry of Health, Malaysia. ANNEX 48: Clinical Guidelines on COVID-19 Vaccination in Malaysia. 4th ed. [Internet]. 2021 Oct [cited 2022 Sep 20]. Available from: https://covid-19.moh.gov.my/garis-panduan/garis-panduan-kkm/ANNEX_48_CLINICAL_GUIDELINES_FOR_COVID_IN_MALAYSIA_4th_EDITION_19102021_FINALE.pdf
18. Johns Hopkins Medicine. Bell's Palsy [Internet]. 2022 [cited 2022 Sep 20]. Available from: <https://www.hopkinsmedicine.org/health/conditions-and-diseases/bells-palsy#:~:text=Bell's%20palsy%20is%20an%20unexplained,strike%20anyone%20at%20any%20age>
19. Cirillo N, Doan R. The association between COVID-19 vaccination and Bell's palsy. *Lancet Infect Dis*. 2022 Jan;22(1):5-6. Available from: [https://doi.org/10.1016/S1473-3099\(21\)00467-9](https://doi.org/10.1016/S1473-3099(21)00467-9)
20. Marshall M, Ferguson ID, Lewis P, Jaggi P, Gagliardo C, Collins JS, Shaughnessy R, Caron R, Fuss C, Corbin KJE, Emuren L, Faherty E, Hall EK, Di Pentima C, Oster ME, Painsil E, Siddiqui S, Timchak DM, Guzman-Cottrill JA. Symptomatic Acute Myocarditis in 7 Adolescents After Pfizer-BioNTech COVID-19 Vaccination. *Pediatrics*. 2021 Sep;148(3):e2021052478. Available from: <https://doi.org/10.1542/peds.2021-052478>
21. Truong DT, Dionne A, Muniz JC, McHugh KE, Portman MA, Lambert LM, Thacker D, Elias MD, Li JS, Toro-Salazar OH, Anderson BR, Atz AM, Bohun CM, Campbell MJ, Chrisant M, D'Addese L, Dummer KB, Forsha D, Frank LH, Frosch OH, Gelehrter SK, Giglia TM, Hebson C, Jain SS, Johnston P, Krishnan A, Lombardi KC, McCrindle BW, Mitchell EC, Miyata K, Mizzi T, Parker RM, Patel JK, Ronai C, Sabati AA, Schauer J, Sexson Tejtel SK, Shea JR, Shekerdemian LS, Srivastava S, Votava-Smith JK, White S, Newburger JW. Clinically Suspected Myocarditis Temporally Related to COVID-19 Vaccination in Adolescents and Young Adults: Suspected Myocarditis After COVID-19 Vaccination. *Circulation*. 2022 Feb;145(5):345-356. Available from: <https://doi.org/10.1161/circulationaha.121.056583>
22. World Health Organization (WHO). Guidance for clinical case management of thrombosis with thrombocytopenia syndrome (TTS) following vaccination to prevent coronavirus disease (COVID-19) [Internet]. 2021 Jul 19 [cited 2022 Sep 20]. Available from: <https://apps.who.int/iris/bitstream/handle/10665/342999/WHO-2019-nCoV-TTS-2021.1-eng.pdf?sequence=1&isAllowed=y>



How to report adverse events?

For more information, visit www.npra.gov.my

By reporting suspected adverse events to the NPRA, you help us learn more about the benefits and risks of vaccines – so we can all make better informed decisions.

Always report – so we can learn more about that vaccine and improve how it is used.



For healthcare professionals:

Report adverse events to the NPRA ([as healthcare professional](#)) through

- Pharmacy hospital information system (PHIS), OR
- [Online web form](#), OR
- [Submission of manual form](#) via mail/email



For consumers:

Inform your healthcare providers at your health facility to make a report on your behalf.

Alternatively, report adverse events to the NPRA ([as consumer](#)) through

- [Online web form \(ConSERF\)](#), OR
- [Submission of manual form \(ConSERF\)](#) via mail/email

You are encouraged to first discuss with your healthcare providers regarding the adverse events before reporting directly to NPRA.