

NO	PRODUCT (ACTIVE INGREDIENT)	ADDITIONAL INDICATION	MARKETING AUTHORIZATION HOLDER
1.	1.1 ZELDOX CAPSULE 20MG 1.2 ZELDOX CAPSULE 1.3 40MG 1.4 ZELDOX CAPSULE 60MG ZELDOX CAPSULE 80MG (Ziprasidone hydrochloride monohydrate)	Ziprasidone is indicated for the treatment of manic or mixed episodes associated with bipolar disorder, with or without psychotic features. Treatment of mania with ziprasidone for more than 3 weeks and prophylactic use in bipolar disorder have not been systematically evaluated in controlled clinical trials.	Pfizer (Malaysia) Sdn. Bhd. Level 3 & 4, Bangunan Palm Grove, No. 14, Jalan Glenmarie (Persiaran Kerjaya), Section U1, 40150 Shah Alam, Selangor.
2.	2.1 EBIXA FILM-COATED TABLET 10MG (Memantine hydrochloride)	Ebixa may be useful as monotherapy or as adjunctive therapy with cholinesterase inhibitors for the symptomatic treatment of patients with moderate to severe dementia of the Alzheimer's type.	JDH Pharmaceutical Sdn. Bhd. Lot 6, Persiaran Perusahaan Seksyen 23, Kawasan Perusahaan Shah Alam, 40300 Shah Alam, Selangor.
3.	3.1 SIFROL TABLET 0.125MG 3.2 SIFROL TABLET 3.3 0.25MG SIFROL TABLET 1MG (Pramipexole)	Sifrol is indicated for the symptomatic treatment of idiopathic Restless Legs Syndrome.	Boehringer Ingelheim (Malaysia) Sdn. Bhd. 2nd Floor, Wisma UEP, Jalan USJ 10/1A, Pusat Perniagaan USJ 10, 46720 Subang Jaya, Selangor.
4.	4.1 REQUIP TABLET 0.25MG 4.2 REQUIP TABLET 4.3 1MG 4.4 REQUIP TABLET	Treatment of idiopathic/ primary Restless Legs Syndrome.	GlaxoSmithKline Pharmaceutical Sdn. Bhd. 8th Floor, Menara Lien Hoe, No. 8, Persiaran Tropicana,

		2MG REQUIP TABLET 5MG (Ropinirole hydrochloride)		47410 Petaling Jaya, Selangor.
5.	5.1	CILODEX (Ciprofloxacin 0.3% and dexamethasone 0.1%) Sterile Otic Suspension (Ciprofloxacin and Dexamethasone)	Acute Otitis Media in paediatric patients (age 6 months and older) with tympanostomy tubes due to Staphylococcus aureus, Streptococcus pneumoniae, Haemophilus influenzae, Moraxella catarrhalis and Pseudomonas aeruginosa.	Alcon Laboratories (M) Sdn. Bhd. No. 1001A & 1001D, 9th Floor, Wisma Glomac 3, Kompleks Kelana Centre Point, No. 3, Jalan SS7/19, Kelana Jaya, 47301 Petaling Jaya, Selangor.
6.	6.1	ARIXTRA 2.5MG/0.5ML SOLUTION FOR INJECTION (Fondaparinux sodium)	Prevention of Venous Thrombotic Events (VTE) in medical patients who are at risk of thromboembolic complications due to restricted mobility during acute illness.	GlaxoSmithKline Pharmaceutical Sdn. Bhd. 8th Floor, Menara Lien Hoe, No. 8, Persiaran Tropicana, 47410 Petaling Jaya, Selangor.
7.	7.1 7.2 7.3 7.4	MABTHERA 100MG/10ML VIALS MABTHERA 500MG/50ML VIALS MABTHERA VIALS 100MG/10ML MABTHERA VIALS 500MG/50ML (Rituximab)	Mabthera is indicated for the treatment of patients with follicular lymphoma as maintenance treatment, after response to induction therapy.	Roche (Malaysia) Sdn. Bhd. 14th Floor, West Block, Wisma Selangor Dredging, 142, Jalan Ampang, 50450 Kuala Lumpur.

DCA 187 (23/11/2006)

NO	PRODUCT (ACTIVE INGREDIENT)	ADDITIONAL INDICATION	MARKETING AUTHORIZATION HOLDER
----	-----------------------------------	-----------------------	--------------------------------------

1.	1.1	TOPAMAX TABLET 25MG	Topamax is indicated as monotherapy in patients with newly diagnosed epilepsy or for conversion to monotherapy in patients with epilepsy.	Johnson & Johnson Sdn. Bhd. Lot 3 & 5, Jalan Tandang, 46050 Petaling Jaya, Selangor.
	1.2	TOPAMAX TABLET		
	1.3	50MG		
	1.4	TOPAMAX TABLET 100MG	<u>Dosage :</u> <i>Monotherapy Epilepsy</i>	
	1.5	TOPAMAX TABLET	<i>General</i>	
	1.6	200MG	When concomitant antiepileptic drugs are withdrawn to achieve monotherapy with topiramate, consideration should be given to the effects this may have on seizure control. Unless safety concerns require an abrupt withdrawal of the concomitant AED, a gradual discontinuation at the rate of approximately one-third of the concomitant AED dose every 2 weeks is recommended.	
	1.7	TOPAMAX TABLET		
	1.8	25MG		
	1.9	TOPAMAX TABLET 50MG		
	1.10	TOPAMAX TABLET 100MG		
		TOPAMAX TABLET 200MG	When enzyme inducing drugs are withdrawn, topiramate levels will increase. A decrease in Topamax dosage may be required if clinically indicated.	
		TOPAMAX SPRINKLE CAPSULE 15MG	<i>Adults</i> Titration should begin at 25mg nightly for 1 week. The dosage should then be increased at 1 or 2-week intervals by increments of 25 or 50mg/day, administered in two divided doses. If the patients is unable to tolerate the titration regimen, smaller increments or longer intervals between increments can be used. Dose and titration rate should be guided by clinical outcome.	
		TOPAMAX SPRINKLE CAPSULE 25MG		
		(Topiramate)		
			The recommended initial target dose for topiramate monotherapy in adults is 100mg/day and the maximum recommended daily dose is 500mg. Some patients with refractory forms of epilepsy have tolerated topiramate monotherapy at doses of 1000mg/day. These dosing recommendations apply to all adults including elderly in absence of underlying renal disease.	
			<i>Children aged 2 and above</i> Treatment of children aged 2 years and above should begin at 0.5 to 1mg/kg nightly for the first week. The dosage should then be increased at 1 or 2-week intervals by increments of 0.5 to 1mg/kg/day, administered in 2 divided doses.	

			The recommended initial target dose range for topiramate monotherapy in children aged 2 years and above is 3 to 6mg/kg/day. Children with recently diagnosed partial onset seizures have received doses of up to 500mg/day.	
2.	2.1 2.2 2.3 2.4	LEXAPRO 5 MG TABLET LEXAPRO 10 MG TABLET LEXAPRO 15MG TABLET LEXAPRO 20 MG TABLET (Escitalopram oxalate)	i) Treatment of Social Anxiety Disorder (social phobia). ii) Treatment of Generalized Anxiety Disorder.	JDH Pharmaceutical Sdn. Bhd. Lot 6, Persiaran Perusahaan Seksyen 23, Kawasan Perusahaan Shah Alam, 40300 Shah Alam, Selangor.
3.	3.1 3.2 3.3	EFEXOR XR 37.5MG CAPSULE EFEXOR XR 75MG CAPSULE EFEXOR XR 150MG CAPSULE (Vanlafaxine hydrochloride)	Panic disorder, including prevention of relapse.	Wyeth (M) Sdn. Bhd. 701 & 801 Block C (Menara Glomac), Kelana Business Centre, No. 97 Jalan SS 7/2, Kelana Jaya, 47301 Petaling Jaya, Selangor.

DCA 186 (19/10/2006)

NO		PRODUCT (ACTIVE INGREDIENT)	ADDITIONAL INDICATION	MARKETING AUTHORIZATION HOLDER
1.	1.1 1.2 1.3 1.4	MABTHERA 100MG/10ML VIALS MABTHERA 500MG/50ML VIALS MABTHERA VIALS 100MG/10ML	Mabthera in combinations with methotrexate is indicated for the treatment of adult patients with severe active rheumatoid arthritis who have had an inadequate response or intolerance to other disease-modifying anti-rheumatic drugs including one or more tumour necrosis factor (TNF) inhibitor therapies.	Roche (Malaysia) Sdn. Bhd. 14th Floor, West Block, Wisma Selangor Dredging, 142, Jalan Ampang, 50450 Kuala Lumpur.

		MABTHERA VIALS 500MG/50ML (Rituximab)		
2.	2.1	HERCEPTIN VIAL 440MG POWDER FOR CONCENTRATE FOR SOLUTION FOR INFUSION	<i>Early Breast Cancer (EBC)</i> Herceptin is indicated for the treatment of patients with HER2 positive early breast cancer following surgery, chemotherapy (neoadjuvant or adjuvant) and radiotherapy (if applicable).	Roche (Malaysia) Sdn. Bhd. 14th Floor, West Block, Wisma Selangor Dredging, 142, Jalan Ampang, 50450 Kuala Lumpur.
	2.2	HERCEPTIN VIAL 150MG POWDER FOR CONCENTRATE FOR SOLUTION FOR INFUSION		

DCA 185 (29/9/2006)

NO		PRODUCT (ACTIVE INGREDIENT)	ADDITIONAL INDICATION	MARKETING AUTHORIZATION HOLDER
1.	1.1	SAIZEN 1.33MG FREEZE-DRIED POWDER	Treatment of growth disturbance (current height SDS <-2.5 and parental adjusted height SDS<-1) in short children born small for gestational age (SGA) with a birth weight and/or length below -2 SD, who failed to show catch-up growth (HV SDS < 0 during the last year) by 4 years of age or later.	Diethelm Malaysia Sdn. Bhd. 74, Jalan Universiti, 46200 Petaling Jaya, Selangor.
	1.2	SAIZEN 3.33MG INJECTION		
	1.3	SAIZEN 8MG CLICK.EASY (Recombinant Human Somatropin)		
2.	2.1	ERBITUX 2MG/ML SOLUTION FOR INFUSION (Cetuximab)	Erbitux in combination with radiation therapy is indicated for the treatment of patients with locally advanced squamous cell cancer of the head and neck.	Merck Sdn. Bhd. No. 4, Jalan U1/26, Section U1, Hicom Glenmarie Industrial Park, 40150 Shah Alam, Selangor.

DCA 184 (24/8/2006)

NO		PRODUCT (ACTIVE INGREDIENT)	ADDITIONAL INDICATION	MARKETING AUTHORIZATION HOLDER
1.	1.1	FLUDARA 10MG FILM COATED TABLETS	<u>Fludara Oral Tablet</u> Treatment of B-cell chronic lymphocytic leukemia (CLL) in patients with sufficient bone marrow reserves.	Zuellig Pharma Sdn. Bhd. Level 3A, Block B, No.10 Jalan Bersatu 13/4, 46200 Petaling Jaya, Selangor.
	1.2	FLUDARA INJECTION (Fludarabine phosphate)	First line treatment with Fludara oral should only be initiated in patients with advanced disease, Rai stages III/IV (Binet stage C), Rai stages I/II (Binet stage A/B) where the patient has related symptoms or evidence of progressive disease. <u>Fludara Injection</u> Treatment of B-cell chronic lymphocytic leukemia (CLL) in patients with sufficient bone marrow reserves. First line treatment with Fludara should only be initiated in patients with advanced disease, Rai stages III/IV (Binet stage C), or Rai stages I/II (Binet stage A/B) where patient has related symptoms or evidence of progressive disease.	

DCA 183 (27/7/2006)

NO		PRODUCT (ACTIVE INGREDIENT)	ADDITIONAL INDICATION	MARKETING AUTHORIZATION HOLDER
1.	1.1	ARIXTRA 2.5MG/0.5ML SOLUTION FOR INJECTION (Fondaparinux sodium)	Prevention of venous thromboembolic events (VTE) in patients undergoing abdominal surgery who are at risk of thromboembolic complications.	GlaxoSmithKline Pharmaceutical Sdn. Bhd. 8th Floor, Menara Lien Hoe, No. 8, Persiaran Tropicana, 47410 Petaling Jaya, Selangor.
2.	2.1	KEPPRA 250MG FILM COATED TABLET	Keppra is indicated as adjunctive therapy in the treatment of partial onset seizures with or without secondary generalization in adults and <u>children from 4 years of age</u> with epilepsy.	UCB Pharma Asia Pacific Sdn. Bhd. Level 2 & 3, Block B, Plaza Damansara, No. 45 Medan Setia 1,
	2.2	KEPPRA 500MG FILM COATED TABLET		

	2.3	KEPPRA 1000MG FILM COATED TABLET (Levetiracetam)		Bukit Damansara, 50490 Kuala Lumpur.
3.	3.1 3.2	XELODA TABLET 150MG XELODA TABLET 500MG (Capecitabine)	Xeloda is indicated for the adjuvant treatment of patients following surgery of stage III (Dukes? stage C) colon cancer.	Roche (Malaysia) Sdn. Bhd. 14th Floor, West Block, Wisma Selangor Dredging, 142, Jalan Ampang, 50450 Kuala Lumpur.
4.	4.1	INVANZ INJECTION (Ertapenem sodium)	<p>1. <u>Pediatric use</u></p> <p>Invanz is indicated for the treatment of pediatric patients with moderate to severe infections caused by susceptible strains of microorganisms, as well as initial empiric therapy prior to identification or causative organisms in the infections listed below:-</p> <ol style="list-style-type: none"> 1. Complicated intra-abdominal infections. 2. Complicated skin and skin structure infections. 3. Community acquired pneumonia. 4. Acute pelvic infections including postpartum endomyometritis, septic abortion and post surgical gynaecologic infections. <p>1. <u>Diabetic foot infections in adult patients</u></p> <p>Invanz is indicated for the treatment of adult patients with moderate to severe infections caused by susceptible strains of microorganisms, as well as initial empiric therapy prior to identification or causative organisms in the infections listed below:-</p> <ol style="list-style-type: none"> 1. Complicated skin and skin structure infections including diabetic lower extremity and diabetic foot infections. 	Merck Sharp & Dohme (I.A.) Corp. Letter Box 1502, 15th Floor, Menara Merais, No.1, Jln 19/3, 46300 Petaling Jaya, Selangor.

5.	5.1	FEMARA 2.5MG FILM-COATED TABLET (Letrozole)	Adjuvant treatment of postmenopausal women with hormone receptor positive early breast cancer.	Novartis Corporation (Malaysia) Sdn. Bhd. Lot 9, Jalan 26/1, Seksyen 26, Kaw. Perindustrian HICOM, 40400 Shah Alam, Selangor.
6.	6.1 6.2 6.3 6.4 6.5 6.6 6.7 6.8 6.9 6.10	TOPAMAX TABLET 25MG TOPAMAX TABLET 50MG TOPAMAX TABLET 100MG TOPAMAX TABLET 200MG TOPAMAX TABLET 25MG TOPAMAX TABLET 50MG TOPAMAX TABLET 100MG TOPAMAX TABLET 200MG TOPAMAX SPRINKLE CAPSULE 15MG TOPAMAX SPRINKLE CAPSULE 25MG (Topiramate)	Topamax is indicated in adults for the prophylaxis of migraine headache. The usefulness of Topamax in the acute treatment of migraine headache has not been studied.	Johnson & Johnson Sdn. Bhd. Lot 3 & 5, Jalan Tandang, 46050 Petaling Jaya, Selangor.
7.	7.1	REMICADE POWDER FOR CONCENTRATE FOR SOLUTION FOR INFUSION (Infliximab)	Treatment of moderate to severe plaque psoriasis in adults who failed to respond to, or who have a contraindication to, or are intolerant to other systemic therapy including cyclosporine, methotrexate or PUVA.	Schering-Plough Sdn. Bhd. Level 3-1, CP Tower, No. 11, Jalan 16/11, Pusat Dagang Seksyen 16, 46350 Petaling Jaya, Selangor.

8.	8.1 8.2	<p>FEIBA (FACTOR VIII INHIBITOR BYPASSING FRACTION) POWDER FOR INJECTION 500U</p> <p>FEIBA (FACTOR VIII INHIBITOR BYPASSING FRACTION) POWDER FOR INJECTION 1000U</p> <p>(Human Plasma Protein with a Factor VIII Bypassing Inhibitor Activity)</p>	<ol style="list-style-type: none"> 1. Treatment and prophylaxis of haemorrhages in haemophilia B patients with inhibitors. 2. Treatment and prophylaxis of haemorrhages in non-haemophilic patients who have developed inhibitors to Factors VIII, IX and XI. 3. Individual reports exist about the used of FEIBA in the treatment of patients with acquired inhibitors to Factors X and XIII. 4. FEIBA was also used in combination with Factor VIII concentrate for a long-term therapy to achieve a complete and permanent elimination of the Factor VIII inhibitor so as to allow for regular treatment with Factor VIII concentrate as in patients without inhibitor. 	<p>United Italian Trading (M) Sdn. Bhd. No. 1, 2nd Floor, The Highway Centre, Jalan 51/205, 46050 Petaling Jaya, Selangor.</p>
----	----------------	---	--	--

DCA 182 (29/6/2006)

NO	PRODUCT (ACTIVE INGREDIENT)	ADDITIONAL INDICATION	MARKETING AUTHORIZATION HOLDER
1.	<p>1.1 Temodal Capsules 5mg</p> <p>1.2 Temodal Capsules 20mg</p> <p>1.3 Temodal Capsules 100mg</p> <p>1.4 Temodal Capsules 250mg</p> <p>(Temozolomide)</p>	<p>Temodal capsules are indicated for the treatment of patients with newly diagnosed glioblastoma multiforme concomitantly with radiotherapy and subsequently as adjuvant monotherapy treatment.</p>	<p>Schering-Plough Sdn. Bhd. Level 3-1, CP Tower, No. 11, Jalan 16/11, Pusat Dagang Seksyen 16, 46350 Petaling Jaya, Selangor.</p>
2.	<p>2.1 Cellcept Capsule 250mg</p> <p>2.2 Cellcept Tablet 500mg</p> <p>(Mycophenolate mofetil)</p>	<p>Cellcept is indicated for induction and maintenance treatment of lupus nephritis. Cellcept should be used concomitantly with corticosteroids.</p>	<p>Roche (Malaysia) Sdn. Bhd. 14th Floor, West Block, Wisma Selangor Dredging, 142, Jalan Ampang, 50450 Kuala</p>

DCA 181 (29/5/2006)

NO	PRODUCT (ACTIVE INGREDIENT)	ADDITIONAL INDICATION	MARKETING AUTHORIZATION HOLDER
1.	1.1 Lipitor 10mg tablet 1.2 Lipitor 20mg tablet 1.3 Lipitor 40mg tablet 1.4 Lipitor 80mg tablet 1.5 Lipitor 10mg tablet 1.6 Lipitor 20mg tablet 1.7 Lipitor 40mg tablet (Atorvastatin)	<p>Prevention of Cardiovascular Disease</p> <ol style="list-style-type: none"> 1. In adult patients without clinically evident coronary heart disease, but with multiple risk factors for coronary heart disease such as age, smoking, hypertension, low HDL-C, or a family history of early coronary heart disease, Lipitor is indicated to: <ol style="list-style-type: none"> 1. Reduce the risk of myocardial infarction. 2. Reduce the risk of stroke. 3. Reduce the risk for revascularization procedures and angina. 1. In patients with type 2 diabetes, and without clinically evident coronary heart disease, but with multiple risk factors for coronary heart disease such as retinopathy, albuminuria, smoking, or hypertension. Lipitor is indicated to: <ol style="list-style-type: none"> 1. Reduce the risk of myocardial infarction. 2. Reduce the risk of stroke. 	<p>Pfizer (Malaysia) Sdn. Bhd. Level 3 & 4, Bangunan Palm Grove, No. 14, Jalan Glenmarie (Persiaran Kerjaya), Section U1, 40150 Shah Alam, Selangor.</p>
2.	2.1 Mabthera 100mg/10ml Vial 2.2 Mabthera 500mg/50ml Vial 2.3 Mabthera 100mg/10ml Vial 2.4 Mabthera 500mg/50ml Vial (Rituximab)	<p>Mabthera is indicated for the treatment of previously untreated patients with stage III-IV follicular lymphoma in combination with CVP chemotherapy.</p>	<p>Roche (Malaysia) Sdn. Bhd. 14th Floor, West Block, Wisma Selangor Dredging, 142, Jalan Ampang, 50450 Kuala Lumpur.</p>

NO	PRODUCT (ACTIVE INGREDIENT)	ADDITIONAL INDICATION	MARKETING AUTHORIZATION HOLDER
1.	1.1 Aromasin 25mg Tablets (Exemestane)	Aromasin is indicated for adjuvant treatment of postmenopausal women with estrogen-receptor positive early breast cancer who have received 2 to 3 years of tamoxifen and are switched to Aromasin for completion of a total of 5 consecutive years of adjuvant hormonal therapy.	Pfizer (Malaysia) Sdn. Bhd. Level 3 & 4, Bangunan Palm Grove, No. 14, Jalan Glenmarie (Persiaran Kerjaya), Section U1, 40150 Shah Alam, Selangor.
2.	2.1 AERIUS Syrup 0.5 mg/ ml (Desloratadine)	To be used for children from age 1 year old.	Schering-Plough Sdn. Bhd. Level 3-1, CP Tower, No. 11, Jalan 16/11, Pusat Dagang Seksyen 16, 46350 Petaling Jaya, Selangor.
3.	3.1 HERCEPTIN 440MG VIALS 3.2 HERCEPTIN VIAL 150MG (Trastuzumab)	<p>Herceptin is indicated for the treatment of patients with metastatic breast cancer who have tumours that overexpress HER2 in combination with docetaxel for the treatment of those patients who have not received chemotherapy for their metastatic disease.</p> <p>Hence, the final indication would be:-</p> <p>Herceptin is indicated for the treatment of patients with metastatic breast cancer who have tumours that overexpress HER2:</p> <ul style="list-style-type: none">▪ as a monotherapy for the treatment of those patients who have received one or more chemotherapy regimens for their metastatic disease.▪ in combination with paclitaxel or docetaxel for the treatment of those patients who have not received chemotherapy for their metastatic disease.	Roche (Malaysia) Sdn. Bhd. 14th Floor, West Block, Wisma Selangor Dredging, 142, Jalan Ampang, 50450 Kuala Lumpur.

DCA 178 (23/2/2006)

NO		PRODUCT (ACTIVE INGREDIENT)	ADDITIONAL INDICATION	MARKETING AUTHORIZATION HOLDER
1.	1.1	REMICADE POWDER FOR CONCENTRATE FOR SOLUTION FOR INFUSION (Infliximab)	Psoriatic arthritis : <ul style="list-style-type: none"> ▪ Reduction of signs and symptoms of arthritis. ▪ Improvement in physical function. ▪ Reduction in psoriasis as measured by PASI (an index which combines symptom evaluation and body surface area). 	Schering-Plough Sdn. Bhd. Level 3-1, CP Tower, No. 11, Jalan 16/11, Pusat Dagang Seksyen 16, 46350 Petaling Jaya, Selangor.
2.	2.1 2.2	EMEND 80MG CAPSULE EMEND 125MG CAPSULE (Aprepitant)	Emend, in combination with other antiemetic agents, is indicated for the prevention of acute and delayed nausea and vomiting associated with initial and repeat courses of: <ul style="list-style-type: none"> ▪ moderately emetogenic cancer chemotherapy. 	Merck Sharp & Dohme (I.A.) Corp. Letter Box 1502, 15th Floor, Menara Merais, No.1, Jln 19/3, 46300 Petaling Jaya, Selangor.

DCA 177 (26/1/2006)

NO		PRODUCT (ACTIVE INGREDIENT)	ADDITIONAL INDICATION	MARKETING AUTHORIZATION HOLDER
1.	1.1 1.2	Zyrtec 1mg/mL Oral Solution Zyrtec Oral Drops 10mg/mL (Cetirizine hydrochloride)	Adults and children of 1 year and above. Symptomatic treatment of seasonal rhinitis, perennial allergic rhinitis and urticaria of allergic origin.	UCB Pharma Asia Pacific Sdn. Bhd. Level 2 & 3, Block B, Plaza Damansara, No. 45 Medan Setia 1, Bukit Damansara, 50490 Kuala Lumpur.
2.	2.1 2.2	Oxis Turbuhaler 4.5 Mcg/ Dose Oxis Turbuhaler 9.0 Mcg/ Dose (Formoterol Fumarate)	Oxis is indicated as an add on therapy to maintenance treatment with inhaled corticosteroids, for the relief of broncho-obstructive symptoms and prevention of exercise-induced symptoms in patients with asthma when adequate treatment with	AstraZeneca Sdn. Bhd. Wisma Prima 17, Jalan Seri Semantan Satu, Damansara Heights,

			corticosteroids is not sufficient.	50490 Kuala Lumpur.
3.	3.1	RECORMON PRE-FILLED SYRINGES 500IU/0.3ML	Recormon is indicated for the treatment of anemia in adult patients with multiple myeloma, low grade non-Hodgkin?s lymphoma or chronic lymphocytic leukemia, who have a relative erythropoietin deficiency and are receiving anti-tumour therapy. Deficiency is defined as an inappropriately low serum erythropoietin level in relation to the degree of anemia.	Roche (Malaysia) Sdn. Bhd. 14th Floor, West Block, Wisma Selangor Dredging, 142, Jalan Ampang, 50450 Kuala Lumpur.
	3.2	RECORMON PRE-FILLED SYRINGES 1000IU/0.3ML		
	3.3	RECORMON PRE-FILLED SYRINGES 2000IU/0.3ML		
	3.4	RECORMON PRE-FILLED SYRINGES 4000IU/0.3ML		
	3.5	RECORMON PRE-FILLED SYRINGES 5000IU/0.3ML		
	3.6	RECORMON PRE-FILLED SYRINGES 10000IU/0.6ML		
	3.7	RECORMON PRE-FILLED SYRINGES 6000IU/0.3ML		
	3.8	RECORMON 10,000 FOR RECO-PEN		
	3.9	RECORMON 20,000 IU FOR RECO-PEN		
	3.10	RECORMON 50,000 MULTIDOSE		
	3.11	RECORMON 100,000 MULTIDOSE		
	3.12	RECORMON VIAL 1000IU		
	3.13	RECORMON VIAL 2000IU		
	3.14	RECORMON VIAL 5000IU		
	3.15	RECORMON VIAL		

		10000IU (Epoetin Beta)		
4.	4.1	Enbrel Injection 25mg/Vial (Etanercept)	Enbrel is indicated in the treatment of adults with moderate to severe plaque psoriasis who failed to respond to, or who have contraindication to, or are intolerant to other systemic therapy including cyclosporine, methotrexate or PUVA.	Wyeth (M) Sdn. Bhd. 701 & 801 Block C (Menara Glomac), Kelana Business Centre, No. 97 Jalan SS 7/2, Kelana Jaya, 47301 Petaling Jaya Selangor.