

Maklumat tambahan indikasi untuk upload pada laman web

Year 2013

Products Approved For Additional Indication (DCA 261 – 28 Februari 2013)

| NO | PRODUCT<br>(ACTIVE INGREDIENT)   | ADDITIONAL INDICATION   | MARKETING<br>AUTHORIZATION HOLDER  |
|----|--|---|--|
| 1. | 1.1 <b>Xamiol Gel</b><br>[ Calcipotriol hydrate 50mcg & Betamethasone dipropionate 0.643mg (equivalent to 0.5mg betamethasone) ]   | <p>➤ Indication:<br/><i>Topical treatment of scalp and non-scalp plaque psoriasis vulgaris in adults.</i></p> <p>➤ Posology:<br/><i>Xamiol Gel should be applied to affected areas once daily. The recommended treatment period is 4 weeks for scalp areas and 8 weeks for “non-scalp” areas.</i></p>   | <b>DKSH MALAYSIA SDN BHD</b><br>74, Jalan Universiti,<br>P.O.Box 77,<br>46200 Petaling Jaya,<br>Selangor.  |
| 2. | 2.1 <b>Votrient Film-Coated Tablet 200mg</b><br>[ Pazopanib Hydrochloride 217 mg (equivalent to pazopanib 200mg) ]<br><br>2.2 <b>Votrient Film-Coated Tablet 400mg</b><br>[Pazopanib Hydrochloride 433 mg (equivalent to pazopanib 400mg)] | <p>➤ Indication:<br/><br/><i>Votrient is indicated for the treatment of patients with advanced Soft Tissue Sarcoma (STS), who had received prior chemotherapy.</i></p> <p><i>Limitation of use: The efficacy of Votrient for the treatment of patients with adipocytic STS or gastrointestinal stromal tumors has not been demonstrated.</i></p> <p>➤ Posology:<br/><br/><i>The recommended dose of Votrient for the treatment of STS is 800mg orally once daily.</i></p> | <b>GLAXOSMITHKLINE PHARMACEUTICAL SDN. BHD.</b><br>Level 6, Quill 9, 112, Jalan Semangat,<br>No.8, Persiaran Tropicana,<br>46300 Petaling Jaya,<br>Selangor. |