

HYCOLIC SYRUP

Hyoscine butylbromide (5mg/5ml)

What is in this leaflet

1. What *Hycolic* is used for
2. How *Hycolic* works
3. Before you use *Hycolic*
4. How to use *Hycolic*
5. While you are using it
6. Side effects
7. Storage and Disposal of *Hycolic*
8. Product description
9. Manufacturer and Product Registration holder
10. Date of revision

What *Hycolic* is used for

It is used for spasm of the gastrointestinal or genito-urinary tracts and in menstrual pain.

How *Hycolic* works

Hyoscine butylbromide reduces muscle spasm and diminishes gastric and intestinal movement. It is used in conditions associated with spasms.

Before you use *Hycolic*

- When you must not use it

- Do not take *Hycolic* if
- allergic to its active ingredient or to other ingredients in it.
 - enlargement of the prostate gland, intestinal obstruction due to paralysis of the intestinal muscles, and pyloric stenosis (blockage of stomach to the intestines). and increased eye pressure.

- Before you start to use it

- use with care in elderly men and in children, with fever.
- caution if there is increased heart rate in individuals with excess of thyroid hormone, heart failure and surgery.

- Taking other medicines

- Concurrent use of other medicines may increase the effects such as:
- amantadine (for flu)
 - some antihistamines
 - butyrophenones (for schizophrenia)
 - phenothiazines (for mental disorders)
 - tricyclic antidepressants (for depression)

How to use *Hycolic*

- How much to use

Adults: oral, 2 – 4 teaspoonfuls 3 – 4 times daily, the dosage adjusted as needed and tolerated. Children-over 6 years: oral, 2 teaspoonfuls 3 – 4 times daily. Below 6 years: oral, 1 – 2 teaspoonfuls 3 times daily. Infants: oral, 1 teaspoonful 3 times daily, or as directed. Follow all directions given by your doctor and pharmacist carefully. They may differ from the information contained in this leaflet. If you do not understand the instructions on the label, ask your doctor or pharmacist.

- When to use it

Use as directed by your doctor or pharmacist.

- How long to use it

Continue taking *Hycolic* for as long as your doctor recommends.

- If you forget to use it

Consult your doctor or pharmacist on what you should do if you forget to use it. Take the missed dose as soon as you remember. If it is almost time for your next dose, wait until then to take the medicine and skip the missed dose. Do not take a double dose to make up for the missed dose.

- If you use too much (overdose)

Contact your doctor immediately or go to the Emergency Department of your nearest hospital, if you think you or anyone else may have taken too much of this medicine. Overdose may cause increased heart rate, high fever, restlessness, confusion, excitement, hallucination (hear, smell, taste or feel things that do not exist), rash, circulatory failure and breathing difficulty.

While you are using *Hycolic*

- Things you must do

Take your medicine exactly as your doctor has told you. Tell the doctors, dentists and pharmacists treating you that you are taking *Hycolic*. Tell the doctor immediately if you are pregnant or breastfeeding.

- Things you must not do

Do not stop taking the medicine unless advised by your doctor. Do not take any new medicines without consulting your doctor. Do not give *Hycolic* to anyone else, even if they have the same symptoms or condition as you.

- Things to be careful of

Safety for use in pregnancy and breastfeeding has not been established.

Side effects

Dry mouth, increased eye pressure, blurred vision, flushing of skin, increased and irregular heartbeats, urinary retention, constipation, vomiting and giddiness may occur occasionally.

You may report any side effects or adverse drug reactions directly to the National Centre for Adverse Drug Reaction Monitoring by calling Tel: 03-78835550, or visiting the website npra.moh.gov.my (Public → Reporting Medicinal Problems / Side Effects / AEFI / Vaccine Safety).

Storage and Disposal of *Hycolic*

- Storage

Store in cool, dry place below 25°C. Protect from light. Keep out of reach of children.

- Disposal

Ask your pharmacist how to dispose of medicines no longer required.

Product description

- What it looks like

Plain coloured, raspberry flavoured syrup.

- Ingredients

- Active ingredient: hyoscine butylbromide
- Inactive ingredients: methyl parabens, propyl parabens, raspberry flavour, liquid glucose, sodium saccharin, sodium chloride, deionised water and syrup.

- MAL number:

MAL19930043AZ

Manufacturer and Product Registration Holder

Prime Pharmaceutical Sdn. Bhd.
1505, Lorong Perusahaan Utama 1
Taman Perindustrian Bukit Tengah
14000 Bukit Mertajam, Penang, Malaysia.

Date of revision

17/08/2017

Serial Number

NPRA(R4/1)170817/00217