

# PABRON DM SYRUP (KIDS)

Dextromethorphan HBr 7.5mg/5ml and Bromhexine HCl 4.0mg/5ml

## What is in this leaflet

1. What PABRON DM SYRUP (KIDS) is used for
2. How PABRON DM SYRUP (KIDS) works
3. Before you use PABRON DM SYRUP (KIDS)
4. How to use PABRON DM SYRUP (KIDS)
5. While you are using it
6. Side effects
7. Storage and disposal of PABRON DM SYRUP (KIDS)
8. Product description
9. Manufacturer and Product Registration Holder
10. Date of revision

This leaflet answers some common questions about PABRON DM SYRUP (KIDS). It does not contain all the available information. It does not take the place of talking to your pharmacist or doctor.

All medicines have risks and benefits. Your pharmacist or doctor has weighed the risks of you taking PABRON DM SYRUP (KIDS) against the benefits they expect it will have for you.

If you have any concerns about taking this medicine, ask your pharmacist or doctor.

Keep this leaflet with the medicine.

You may need to read it again.

## What PABRON DM SYRUP (KIDS) is used for

PABRON DM SYRUP (KIDS) is used for the short term symptomatic relief of cough associated with reduction in sputum viscosity.

## How PABRON DM SYRUP (KIDS) works

PABRON DM SYRUP (KIDS) contains the active ingredients Bromhexine HCl and Dextromethorphan HBr.

**Bromhexine HCl** belongs to a group of medicines called mucolytic agents and works by making your mucus thinner so that you can get rid of it more easily.

**Dextromethorphan** belongs to a group of medicines called antitussive (cough suppressant) which suppresses a cough by acting on the cough centre of the brain.

Ask your pharmacist or doctor if you have any questions about this medicine.

## Before you use PABRON DM SYRUP (KIDS)

### -When you must not use it

If you are sensitive to;

- Bromhexine
- Dextromethorphan
- other components of the formulation.

Unsuitable for phenylketonurics (an inherited inability to metabolize (process) the essential amino acid phenylalanine).

Not to be used in children less than 2 years of age.

### -Before you start to use it

Tell your pharmacist or doctor:

If you have peptic ulceration (painful sores or ulcers in the lining of the stomach), asthma, chronic bronchitis (an inflammation of the bronchial tubes, the airways that carry air to your lungs), emphysema (an abnormal increase in the size of the air spaces in the lungs), diabetes, liver function impairment and respiration depression (decrease in a person's ability to breath).

Inform your doctor if you are pregnant or breastfeeding. Safe use during pregnancy and lactation has not been established.

### -Taking other medicines

Tell your doctor if you are taking any other medicines, including any that you buy without a prescription

from a pharmacy, supermarket or health food shop.

Please inform your doctor if you are taking drug-metabolizing enzyme inhibitors like amiodarone, quinidine which may occasionally cause a rise in blood pressure.

## How to use PABRON DM SYRUP (KIDS)

### -How much to use

#### **Children 6 to 12 years old:**

Take 5 ml of this medicine after meals, 3 to 4 times a day or every 6 to 8 hours.

#### **Children over 12 years old:**

Take 10 ml of this medicine after meals, 3 to 4 times a day or every 6 to 8 hours.

Warning (when used for the treatment of cough and cold):

1. Not to be used in children less than 2 years of age.
2. To be used with caution & doctor's advice in children 2 to 6 years of age.

### -When to use it

Use the medication after meal.

### -How long to use it

Continue taking PABRON DM SYRUP (KIDS) for as long as your doctor recommends.

### -If you forget to use it

Take the missed dose as soon as you remember.

If it is almost time for your next dose, wait until then to take the medicine and skip the missed dose.

Do not take a double dose to make up for the missed dose.

# PABRON DM SYRUP (KIDS)

Dextromethorphan HBr 7.5mg/5ml and Bromhexine HCl 4.0mg/5ml

---

## -If you use too much (overdose)

Taking too much of this medicine may cause the following symptoms:

In mild overdose, tachycardia (increase heart rate), high blood pressure, vomiting, mydriasis (dilation of the pupil), diaphoresis (perspiring profusely), nystagmus (involuntary eye movement), euphoria (over excited), loss of motor coordination, and giggling.

In moderate intoxication, in addition to those listed above, hallucinations and a plodding ataxic gait (an unsteady, staggering, or irregular style of walking).

In severe intoxication, other symptoms are agitation or somnolence (sleepiness).

## **While you are using PABRON DM SYRUP (KIDS)**

### -Things you must do

Take your medicine exactly as your doctor has told you.

Tell all the doctors, dentists and pharmacists treating you that you are taking PABRON DM SYRUP (KIDS).

During treatment, do not take any anti-cough medicine.

### -Things you must not do

Do not drink alcohol while taking PABRON DM SYRUP (KIDS).

The sedation effects of alcohol may be increased.

Do not take more than the recommended dose unless your doctor tells you to.

Do not give PABRON DM SYRUP (KIDS) to anyone else, even if they have the same symptoms or condition as you.

### -Things to be careful of

This medication may cause drowsiness and impair performance in test of auditory vigilance in some people. If this

occurs do not drive or operate machinery.

Should not be used for persistent or chronic cough.

## **Side effects**

PABRON DM SYRUP (KIDS) may have unwanted side effects in some people. All medicines can have side effects. Sometimes they are serious, most of the time they are not.

Stop taking the medicine if you have experience: headache, dizziness, sweating, skin rash, constipation, stomach discomfort, vomiting, nausea and gastrointestinal disturbances.

If you get any side effects, even those not mentioned in this leaflet, tell your doctor or pharmacist.

You may report any side effects or adverse drug reactions directly to the National Centre for Adverse Drug Reaction Monitoring by calling Tel: 03-78835550, or visiting the website [portal.bpfk.gov.my](http://portal.bpfk.gov.my) (Consumers → Reporting)

## **Storage and Disposal of PABRON DM SYRUP (KIDS)**

### -Storage

Store below 30°C.

Keep in a cool dry place, protected from light.

Keep out of reach of children.

### -Disposal

Dispose the medicine after the expiry date. Medicines should not be disposed of via wastewater or household waste, Ask your pharmacist how to dispose of medicines no longer required. These measures will help to protect the environment.

## **Product description**

### -What it looks like

PABRON DM SYRUP (KIDS) is a clear, pink viscous liquid with mixed fruits and strawberry odor.

PABRON DM SYRUP (KIDS) is available in 60 ml bottle.

## -Ingredients

-Active ingredient(s):

Each 5 ml syrup contains 7.5 mg of Dextromethorphan HBr and 4.0 mg of Bromhexine HCl.

-Inactive ingredients:

Glycerin, Sodium Benzoate, CMC Sodium 7HOF, Color Ponceau 4R, Mixed Fruits Liquid Flavor, Purified Water, Acesulfame potassium salt, Sorbitol Solution (70% NC), Propylene Glycol, Aspartame, Trisodium Citrate Dihydrate, Strawberry Flavor, Disodium edetate dehydrate, Citric Acid Monohydrate

## -MAL number:

MAL12090003AC

## **Manufacturer**

OLIC (Thailand) Limited  
166 Moo 16,  
Bangpa-In Industrial Estate,  
Udomsorasuyuth Road,  
Bangkrason, Bangpa-in,  
Ayutthaya, Province 13160,  
Thailand.

## **Product Registration Holder:**

HOE Pharmaceuticals Sdn. Bhd.  
Lot 10, Jalan Sultan Mohamed 6,  
Bandar Sultan Suleiman,  
42000 Port Klang, Selangor,  
Malaysia.

## **Date of revision**

-09/03/2015

## **Serial number**

BPFK(R4/1)090315/00038