

DYNAVATE CREAM

Betamethasone Valerate (0.1% w/w)

What is in this leaflet

1. What DYNAVATE CREAM is used for
2. How DYNAVATE CREAM works
3. Before you use DYNAVATE CREAM
4. How to use DYNAVATE CREAM
5. While you are using it
6. Side effects
7. Storage and Disposal of DYNAVATE CREAM
8. Product Description
9. Manufacturer and Product Registration Holder
10. Date of revision

What DYNAVATE CREAM is used for

For the relief of skin inflammation that is responsive to corticosteroid.

How DYNAVATE CREAM works

It is responsible for anti-inflammatory effects to topical application of corticosteroid.

Before you use DYNAVATE CREAM

- When you must not use it

Do not use it if you have:

- Facial redness
- Acne
- Small bumps around the mouth
- Viral infections
- Bacterial and fungal infections
- Hypersensitivity to Betamethasone or the preparation.
- Diabetes
- Increased pressure in the eye

Pregnancy and lactation

Do not use DYNAVATE CREAM if you are pregnant, trying to get pregnant or think you may be pregnant.

Do not use DYNAVATE CREAM if you are breast-feeding. Ask your doctor or pharmacist for advice before taking any medicine.

- Before you start use it

Topical applications of corticosteroid should not be made with an occlusive dressing to large areas of the body because of the increased risk of systemic toxicity. Corticosteroid should not be applied to ulcers of the leg.

- Taking other medicines

Tell your doctor if you are taking any other medicines, including any that you buy without a prescription from a

pharmacy, supermarket or health food shop.

How to use DYNAVATE CREAM

- How much to use

Apply thinly to the affected parts.

Follow all directions given to you by your doctor and pharmacist carefully. They may differ from the information contained in this leaflet. If you do not understand the instructions on the label, ask your doctor or pharmacist for help.

- When to use it

Use two to three times daily or as directed by your doctor or pharmacist.

- How long to use it

Continue taking DYNAVATE CREAM for as long as your doctor recommends.

- If you forget to use it

Consult your doctor or pharmacist on what you should do if you forget to use it.

Apply the cream as soon as you remember. If it is almost time for your next dose, wait until then to take the medicine and skip the missed dose. Do not take a double dose to make up for the missed dose.

- If you use too much (overdose)

For topical application of corticosteroids, problems of over dosage are unlikely to be encountered. Consult the doctor immediately if any unusual serious reactions happened.

While you are using it

- Things you must do

Tell all the doctors, dentists and pharmacists treating you that you are using DYNAVATE CREAM.

Tell your doctor immediately if you become pregnant while using this medication.

- Things you must not do

Do not stop taking the medicine unless advised by your doctor or pharmacist.

Do not take any new medicines without consulting your doctor.

Do not give DYNAVATE CREAM to anyone else, even if they have the same symptoms or condition as you.

- Things to be careful of

Do not use this cream in the presence of infection. Occasionally they may be used with the addition of a suitable antimicrobial medicine in the treatment of infected skin but there is a risk of sensitivity reactions occurring. Long-term topical use is best avoided, especially in children.

Side effects

Like all medicines, DYNAVATE CREAM can cause side effects, although not everybody gets them.

The following local adverse reactions are reported infrequently with topical corticosteroids but may occur more frequently with the use of occlusive dressings. These reactions are listed in an approximate decreasing order of occurrence; burning, itching, irritation, dryness, hair follicle inflammation, abnormal hair growth, acne eruptions, lighting of the skin, softening and breaking down of the skin, small bumps around the mouth, allergic skin inflammation, secondary infection, skin thinning, stretch mark and heat rashes.

Visit your doctor or pharmacist immediately if you experience any side effects after taking this medicine. You may report any side effects or adverse drug reactions directly to the National Centre for Adverse Drug Reaction Monitoring by calling Tel: 03-78835549, or visiting the website npra.moh.gov.my (Public → Reporting Side Effects to Medicines (ConSERF) or Vaccines (AEFI))

Storage and Disposal of DYNAVATE CREAM

- Storage

Keep out of reach of children
Keep container tightly closed
Store in a dry place below 30°C
Protect from light
For external use only

- Disposal

Medicines should not be disposed of via wastewater or household waste. Ask your pharmacist how to dispose of medicines no longer required. These measures will help to protect the environment.

Product Description

- What it looks like

White colour cream with Baby Jo fragrance

DYNAVATE CREAM

Betamethasone Valerate (0.1% w/w)

- Ingredients

- Active ingredient

Betamethasone-17-Valerate

- Inactive ingredients

White Soft Paraffin, Liquid Paraffin,

Emulgate 1000 NI, Chlorocresol,

Deionised Water, Glycerin, Baby Jo

Fragrance

- MAL number:

MAL19950042AZ

Manufacturer & Product Registration

Holder

Dynapharm (M) Sdn. Bhd.

2497, Mk 1, Lorong Perusahaan Baru 5,

Kawasan Perusahaan Perai 3, 13600

Perai, Pulau Pinang, Malaysia

Date of revision

16/07/2018

Serial Number:

NPRA(R4/1)100718/00189