AXCEL KAOPEC SUSPENSION

Light Kaolin / Pectin Citrus (1g/5ml / 20mg/5ml)

What is in this leaflet

- 1. What Axcel Kaopec Suspension is used for.
- 2. How Axcel Kaopec Suspension works.
- 3. Before you use Axcel Kaopec Suspension.
- 4. How to use Axcel Kaopec Suspension.
- 5. While you are using it.
- 6. Side effects.
- 7. Storage and disposal of Axcel Kaopec Suspension.
- 8. Product description.
- Manufacturer and Product Registration Holder.
- 10. Date of revision.

What Axcel Kaopec Suspension is used for

Axcel Kaopec Suspension is an antidiarrhoeal used for the relief of occassional diarrhoea.

How Axcel Kaopec Suspension works

This medicine contain Light Kaolin to help remove toxic substances from the digestive system and increase the bulk of solid waste. It also contains Pectin which is a natural fiber.

Before you use Axcel Kaopec Suspension

-When you must not use it

Do not take Axcel Kaopec Suspension if:

- You are having intestinal obstruction.
- It is not prescribed by a qualified doctor or pharmacist for you
- You are pregnant, trying to get pregnant or think you may be pregnant.
- You are breast-feeding. Ask your doctor or pharmacist for advice before taking any medicine.

-Before you start to use it

Tell your doctor if:

- You are allergic to any medicines, food and dyes.
- You are taking any other medicines (including those not prescribed by your doctor).

-Taking other medicines

You should tell your doctor if you are taking the following medicine as it may interact with your Axcel Kaopec Suspension:

• Antibiotics.

It may still be all right for you to take Axcel Kaopec Suspension and your doctor will be able to decide what is suitable for you.

How to use Axcel Kaopec Suspension

-How much to use

To be administered orally.

To be taken 3 times a day, when necessary.

Children:

6-12 years: 20ml.

-When to use it

Your doctor will decide when it is appropriate for you to use Axcel Kaopec Suspension according to the type of condition you have.

-How long to use it

Continue taking Axcel Kaopec Suspension for as long as your doctor recommends.

-If you forget to use it

Consult your doctor or pharmacist on what you should do if you forget to use it.

Take the missed dose as soon as you remember. If it is almost time for your next dose, wait until then to take the medicine and skip the missed dose. Do not take a double dose to make up for the missed dose.

-If you use too much (overdose)

Contact your doctor immediately or go to the Emergency Department of your nearest hospital, if you think you or anyone else may have taken too much of this medicine. Do this even if there are no signs of discomfort or poisoning. You may need urgent medical attention.

While you are using it

-Things you must do

 Keep yourself hydrated at all time.

- Take your medicine exactly as your doctor has told you.
- Tell all the doctors, dentists and pharmacists treating you that you are taking Axcel Kaopec Suspension.
- Tell your doctor immediately if you become pregnant while talking this medication.

-Things you must not do

- Do not stop taking the medicine unless advised by your doctor.
- Do not take any new medicines without consulting your doctor or pharmacist.
- Do not give Axcel Kaopec Suspension to anyone else, even if they have the same symptoms or condition as you.

-Things to be careful of

 Not recommended for children below 6 years old.

Side effects

Like all medicines, Axcel Kaopec Suspension can cause side effects although not everybody gets them. You are advised to consult doctor/pharmacist for further information. The following common side effects have been reported:

 Severe constipation, which may lead to faecal impaction, may rarely occur in children and elderly patients.

You may report any side effects or adverse drug reactions directly to the National Centre for Adverse Drug Reaction Monitoring by calling Tel: 03-78835550, or visiting the website npra.moh.gov.my (Public → Reporting Medicinal Problems / Side Effects / AEFI / Vaccine Safety).

Storage and disposal of Axcel Kaopec Suspension

-Storage

Keep Axcel Kaopec Suspension out of reach of children.

Store below 30°C. Keep container well closed and protect from light.

AXCEL KAOPEC SUSPENSION

Light Kaolin / Pectin Citrus (1g/5ml / 20mg/5ml)

-Disposal

Do not use Axcel Kaopec Suspension after the expiry date which is stated on the carton.

Medicines should not be disposed of via wastewater or household waste. Ask your pharmacist how to dispose of medicines no longer required. These measures will help to protect the environment.

Product description

-What it looks like

Axcel Kaopec Suspension is a cream coloured with cream flavoured suspension.

-Ingredients

- Active Ingredient: Light Kaolin and Pectin (Citrus)
- Inactive Ingredients:
 Refined Sugar, Methylparaben,
 Sorbitol, Purified Water, Cream
 Flavour and Propylene Glycol.

<u>-MAL Number</u> MAL19911416X

Manufacturer and product registration holder
Kotra Pharma (M) Sdn Bhd
No. 1, 2 & 3, Jalan TTC 12,
Cheng Industrial Estate,
75250 Melaka, Malaysia.

Date of revision

24/03/2018

Seriel number

NPRA(R4/1)200318/00075