

PRIVIN ENEMA

Glycerin/ Sodium Chloride (25% w/v / 15% w/v)

What is in this leaflet

1. What Privin Enema is used for
2. How Privin Enema works
3. Before you use Privin Enema
4. How to use Privin Enema
5. While you are using Privin Enema
6. Side effects
7. Storage and Disposal of Privin Enema
8. Product description
9. Manufacturer and Product Registration holder
10. Date of revision

What Privin Enema is used for

For relief of constipation.

How Privin Enema works

Glycerin promotes movement of the intestines and evacuation of the lower bowel by its irritant action.

Before you use Privin Enema

- When you must not use it

- allergic to glycerin or other ingredients of the product.
- do not use this product when you have nausea, vomiting or abdominal pain unless directed by physician.

- Before you start to use it

It should be used cautiously if you are dehydrated.

- Taking other medicines

The medicine is for topical application, usually will have minimal effect on other medications. However, please inform the doctor or pharmacist if taking other medicines.

How to use Privin Enema

- How much to use

Children: use 10ml enema when necessary. Adult: Use 20ml enema when necessary.

- When to use it

Please follow the advice of the doctor or pharmacist.

- How long to use it

You should check with your doctor or pharmacist.

- If you forget to use it

If forget to use this medicine, use it as soon as remember. Then next, use this medicine at the usual time.

- If you use too much (overdose)

If used too much Privin Enema, contact your doctor, an emergency department or a poison center immediately.

While you are using Privin Enema

- Things you must do

Check with the physician if condition does not improve or if it becomes worse. Inform doctor of medications taking.

- Things you must not do

Do not stop using medication although feeling better after a few days, until the prescribed course is finished.

- Things to be careful of

Treatment should be discontinued if severe irritation or allergy develops. If pregnant or become pregnant or breast-feeding during treatment, should not use Privin Enema unless as advised by doctor.

Side effects

Side effects are mainly due to dehydrating effects of glycerin. Glycerin when used rectally can cause irritation.

You may report any side effects or adverse drug reactions directly to the National Centre for Adverse Drug Reaction Monitoring by calling Tel: 03-78835550, or visiting the website npra.moh.gov.my (Public → Reporting Medicinal Problems / Side Effects / AEFI / Vaccine Safety).

Storage and Disposal of Privin Enema

- Storage

Store in cool, dry place, below 30°C. Protect from light. Keep out of reach of children.

- Disposal

Ask your pharmacist how to dispose of medicines no longer required.

Product description

- What it looks like

Colourless solution.

- Ingredients:

- Active ingredient: glycerin and sodium chloride.
- Inactive ingredients: water.

- MAL number:

MAL19962542X

Manufacturer and Product Registration Holder

Prime Pharmaceutical Sdn. Bhd.
1505, Lorong Perusahaan Utama 1
Taman Perindustrian Bukit Tengah
14000 Bukit Mertajam, Penang, Malaysia.

Date of revision

11/08/2017

Serial number

NPRA(R4/1)100817/00205