

# ISOMIN TABLET

Dexchlorpheniramine maleate (2mg)

## What is in this leaflet

1. What *Isomin* is used for
2. How *Isomin* works
3. Before you use *Isomin*
4. How to use *Isomin*
5. While you are using it
6. Side effects
7. Storage and Disposal of *Isomin*
8. Product description
9. Manufacturer and Product Registration holder
10. Date of revision

## What *Isomin* is used for

For treatment of long term and seasonal allergic or nonallergic rhinitis (nose inflammation or congestion), allergic eye inflammation, itchiness, hives and swelling under the skin.

## How *Isomin* works

*Isomin* can help to relieve symptoms of some allergies and itchy skin rashes. It can be used to treat the itchiness, redness, swelling, tenderness and irritation that can be caused by hay fever, hives, prickly heat, reactions to food/medicines, insect bites or chickenpox rashes.

## Before you use *Isomin*

### - When you must not use it

- Do not take *Isomin* if allergic to its active ingredient or other ingredients in it.
- Ask your doctor or pharmacist for advice before taking any medicine.

### Pregnancy and lactation

Do not take *Isomin* if you are pregnant, trying to get pregnant or think you may be pregnant. Do not take *Isomin* if you are breast-feeding. Ask your doctor or pharmacist for advice before taking any medicine.

### - Before you start to use it

Inform the doctor or pharmacist if having asthma and kidney function impairment, increased pressure of the eye, hyperthyroidism (high thyroid hormones in the blood), heart or blood vessels disease, high blood pressure, intestinal ulcer, obstruction of the stomach and intestines, enlargement of the prostate gland and bladder neck obstruction.

### - Taking other medicines

Tell your doctor if you are taking any other medicines e.g. tricyclic antidepressants or monoamine oxidase (MAO) inhibitors that may increase the effects of medicines. Also, including any medicine that you buy without a prescription from a pharmacy, supermarket or health food shop.

## How to use *Isomin*

### - How much to use

Adults: 2mg every 4 – 6 hours as needed.  
Children 6 – 12 years old: 1mg every 4 – 6 hours as needed. Follow all directions given by your doctor and pharmacist carefully. They may differ from the information contained in this leaflet. If you do not understand the instructions on the label, ask your doctor or pharmacist for help.

### - When to use it

Use as directed by your doctor or pharmacist.

### - How long to use it

Continue taking *Isomin* for as long as your doctor recommends.

### - If you forget to use it

Consult your doctor or pharmacist on what you should do if you forget to use it. Take the missed dose as soon as you remember. If it is almost time for your next dose, wait until then to take the medicine and skip the missed dose. Do not take a double dose to make up for the missed dose.

### - If you use too much (overdose)

Contact your doctor immediately or go to the Emergency Department of your nearest hospital, if you think you or anyone else may have taken too much of this medicine. Do this even if there are no signs of discomfort or poisoning. You may need urgent medical attention. Signs of overdose include clumsiness or unsteadiness, severe dryness of mouth, nose or throat, severe drowsiness, hallucinations (hear, smell, taste or feel things that do not exist), seizures, shortness of breath or troubled breathing.

## While you are using *Isomin*

### - Things you must do

Take your medicine exactly as your doctor has told you.  
Tell the doctors, dentists and pharmacists treating you that you are taking *Isomin*.  
Tell your doctor immediately if you are pregnant.

### - Things you must not do

Do not stop taking the medicine unless advised by your doctor.  
Do not take any new medicines without consulting your doctor or pharmacist.  
Do not give *Isomin* to anyone else, even if they have the same symptoms or condition as you.

### - Things to be careful of

Caution when driving or doing jobs requiring alertness because of possible drowsiness. Avoid alcoholic beverages.

## Side effects

*Isomin* can cause side effects, although not everybody gets them. Visit the doctor or pharmacist immediately if you experience any side effects after taking this medicine. Drowsiness, dryness of mouth, nose and throat, gastrointestinal disturbances, and skin rash may occur. Anaphylactic shock (life-threatening allergic reaction), sensitivity to light, tiredness, low blood pressure, dizziness, insomnia (sleeping difficulty), blurred vision, difficult urination, fast heartbeat, increased sweating, restlessness and pounding of heartbeats. You may report any side effects or adverse drug reactions directly to the National Centre for Adverse Drug Reaction Monitoring by calling Tel: 03-78835550, or visiting the website [npra.moh.gov.my](http://npra.moh.gov.my) (Public → Reporting Medicinal Problems / Side Effects / AEFI / Vaccine Safety).

## Storage and Disposal of *Isomin*

### - Storage

Store in cool, dry place below 30°C. Protect from light. Keep out of reach and sight of children.

### - Disposal

Medicines should not be disposed of via wastewater or household waste. Ask your pharmacist how to dispose of medicines no longer required.

## Product description

### - What it looks like

Red, oblong, biconvex, top-scored tablet.

### - Ingredients

- Active ingredient: dexchlorpheniramine maleate.
- Inactive ingredients: cornstarch, lactose monohydrate, carmoisine certolake, dried cornstarch and magnesium stearate.

### - MAL number:

MAL19930041AZ

## Manufacturer and Product Registration Holder

Prime Pharmaceutical Sdn. Bhd.  
1505, Lorong Perusahaan Utama 1  
Taman Perindustrian Bukit Tengah  
14000 Bukit Mertajam, Penang, Malaysia.

## Date of revision

22/06/2017

## Serial Number

BPFK(R4/1)220617/00149