

REXOMTM WINA ANTAMAC TABLET

Magnesium Trisilicate/Aluminium Hydroxide/Simethicone (250mg/125mg/30mg)

What is in this leaflet

1. What REXOMTM Wina Antamac is used for
2. How REXOMTM Wina Antamac works
3. Before you use REXOMTM Wina Antamac
4. How to use REXOMTM Wina Antamac
5. While you are using REXOMTM Wina Antamac
6. Side effects
7. Storage and disposal of REXOMTM Wina Antamac
8. Product description
9. Manufacturer and Product Registration Holder
10. Date of revision

What REXOMTM Wina Antamac is used for

For the symptomatic relief from indigestion, upset stomach, heartburn, flatulence, gastric hyperacidity.

How REXOMTM Wina Antamac works

REXOMTM Wina Antamac contains antacid to reduce excess stomach acid.

Before you use REXOMTM Wina Antamac

-When you must not use it

It should not be used if you had known to be hypersensitive to any of the components of the formulation.

-Before you start to use it

Caution to people with impaired kidney function.

Pregnancy and lactation

Talk to your doctor before taking this medicine if you are pregnant, might become pregnant or are breastfeeding. Risk-benefit must be considered.

-Taking other medicines

Inform your doctor or pharmacist before taking this medicines, including any that you get without a prescription from your pharmacy, supermarket or health food shop.

REXOMTM Wina Antamac may interfere with the absorption of:

- Tetracycline (as antibiotic)
- Anticholinergics

- Barbiturates (to treat seizure)
- Digoxin (to treat heart problem)
- Warfarin (to prevent blood clot)
- Vitamins (as health supplement)

An interval of 3 hours is required to prevent this interference.

How to use REXOMTM Wina Antamac

-How much to use

Always use REXOMTM Wina Antamac exactly as your doctor has prescribed. You should check with your doctor or pharmacist if in doubt.

The usual dose:

2 tablets.

Tablets should be chewed and swallowed with water.

-When to use it

To be used 4 times daily, after or between meals and at bedtime. Use as directed by your doctor or pharmacist.

-How long to use it

Continue taking REXOMTM Wina Antamac as your doctor recommended. If symptoms persist, consult your doctor.

-If you forget to use it

Consult your doctor or pharmacist on what you should do if you forget to use it.

Take the missed dose as soon as you remember. If it is almost time for your next dose, wait until then to take the medicine and skip the missed dose. Do not take a double dose to make up for the missed dose.

-If you use too much (overdose)

Contact your doctor immediately or go to the Emergency Department of your nearest hospital, if you think you or anyone else may have taken too much of this medicine. Do this even if there are no signs of discomfort or poisoning. You may need urgent medical attention.

Large doses of REXOMTM Wina Antamac may cause hypermagnesemia (high level of magnesium in the blood).

While you are using REXOMTM Wina Antamac

-Things you must do

- Tell all the doctors, dentists and pharmacists treating you that you are taking REXOMTM Wina Antamac.
- Tell your doctor immediately if you become pregnant while taking this medication.

-Things you must not do

- Do not take more than the recommended dose unless your pharmacist or doctor tells you.
- Do not take any new medicines without consulting your doctor or pharmacist.
- Do not give REXOMTM Wina Antamac to anyone else, even if they have the same symptoms or condition as you.

-Things to be careful of

Be careful driving or operating machinery until you know how REXOMTM Wina Antamac affects you.

Side effects

Like all medicines, REXOMTM Wina Antamac can cause side effects, although not everybody gets them.

If side effects persist, do not take the medicine and refer to the doctor.

Tell your doctor if you notice any of the following side effects or notice any other effects not listed:

- Mild constipation or diarrhea
- Nausea
- Vomiting
- Stomach cramps
- Unusual increased in thirst.

You may report any side effects or adverse drug reactions directly to the National Centre for Adverse Drug Reaction Monitoring by calling Tel: 03-78835550, or visiting the website npra.moh.gov.my (Public → Reporting Medicinal Problems / Side Effects / AEFI / Vaccine Safety).

Storage and disposal of REXOMTM Wina Antamac

-Storage

Keep container tightly closed. Protect from heat and light. Store in a cool, dry place.

Keep out of reach of children.

REXOMTM WINA ANTAMAC TABLET

Magnesium Trisilicate/Aluminium Hydroxide/Simethicone (250mg/125mg/30mg)

-Disposal

Medicines should not be disposed of via wastewater or household waste. Ask your pharmacist how to dispose of medicines no longer required. These measures will help to protect the environment.

Product description

-What it looks like

Round with top embossed 'ANTAMAC' and bottom 'KCK', white colour tablet.

-Ingredients

-Active ingredient

250mg Magnesium Trisilicate, 125mg Aluminium Hydroxide and 30mg Simethicone.

-Inactive ingredients

Lactose, Mannitol, Peppermint oil, Menthol, Sodium saccharin, Magnesium stearate, Corn starch, Starch and Deionized water.

-MAL number

MAL19911378X

Manufacturer and Product Registration Holder

KCK Pharmaceutical Industries Sdn. Bhd.
Plot 61, Bayan Lepas Industrial Park
Phase 4, 11900 Penang,
Malaysia.

Date of revision

30/10/2017

Serial Number

NPRA(R4/1)301017/00298