

NEUROBINE FILM COATED TABLET

Thiamine mononitrate/Pyridoxine HCL/Cyanocobalamin (100mg/200mg/0.2mg)

What is in this leaflet

1. What Neurobine F.C. Tablet is used for
2. How Neurobine F.C. Tablet works
3. Before you use Neurobine F.C. Tablet
4. How to use Neurobine F.C. Tablet
5. While you are using it
6. Side Effects
7. Storage and Disposal of Neurobine F.C. Tablet
8. Product Description
9. Manufacturer and Product Registration Holder
10. Date of revision

What Neurobine F.C. Tablet is used for

Neurobine F.C. Tablet is used for disorder of the nervous system, involving damages that affect nerves or group of nerves and multiple nerves. This includes

- nerve damage caused by long term high blood sugar,
- nerve damage results from excessive drinking of alcohol,
- nerve damage caused by drug ingestion, drug or chemical abuse,
- inflammation of a nerve and pain in the one or more nerves especially low back pain, numbness, and/or weakness that radiate from lower back to the buttocks and leg, etc.

It is also used for other disorders associated with deficiency or raised requirement of vitamin B1, B6 and B12.

How Neurobine F.C. Tablet works

Thiamine is an essential nutrient required for carbohydrate metabolism and is involved in nerve transmission.

Pyridoxine, stored mainly in the liver, is involved in the metabolism of amino acid, carbohydrate and fat. It is required for the formation of red blood cell.

Cyanocobalamin is involved in formation and metabolism of amino acids.

Before you use Neurobine F.C. Tablet

- When you must not use it

Do Not take Neurobine F.C. Tablet if:

- You are hypersensitive to vitamins B₁, B₆ or B₁₂.
- You are undergoing Levodopa therapy.

If you are pregnant, trying to get pregnant or think you may be pregnant, consult your doctor before taking this product.

If you breastfeeding, consult your doctor before taking this product.

Cyanocobalamin not to be given to individuals with suspected Vitamin B12 deficiency (without first confirming the diagnosis), individuals with Leber's disease, tobacco amblyopia (decrease in vision's ability due to heavy tobacco consumption) and megaloblastic anaemia of pregnancy.

- Before you start to use it

Tell your doctor or pharmacist if you are allergic to any other medicines or you have any other health related problems.

- Taking other medicines

High vitamin B6 dose decrease action of L-dopa. Seek for your doctor or pharmacist advice if you intended to use Neurobine F.C. Tablet together with other medication. Besides, you must also inform them about all the medicines you are taking, including those without a prescription.

How to use Neurobine F.C. Tablet

- How much to use

1-3 tablets daily as initial, supportive or follow up measure to parenteral therapy.

- When to use it

Neurobine F.C. Tablet should be taken according to the time advised by your doctor or pharmacist. It can be taken with or without food.

- How long to use it

Continue taking Neurobine F.C. Tablet for as long as your doctor recommends.

- If you forgot to use it

Consult your doctor or pharmacist on what you should do if you forget to use it. Take the missed dose as soon as you remember. If it is almost time for your next dose, wait until then to take the dose and skip the missed dose. Do not take a double dose to make up for the missed dose.

- If you use too much (overdose)

Contact your doctor or pharmacist for immediate advice. Do this even if there are no signs of discomfort or poisoning. You may need urgent medical attention.

Individuals in whom intentional overdose is confirmed or suspected should be referred for psychiatric consultation.

While you are using it

- Things you must do

Follow your doctor's instruction when you are taking Neurobine F.C. Tablet.

- Things you must not do

Do not use Neurobine F.C. Tablet after the expiry date which is stated on the carton and blister.

Do not stop taking the medicine unless advised by your doctor.

Do not take any new medicines without consulting your doctor or pharmacist.

Do not give Neurobine F.C. Tablet to anyone else, even if they have the same symptoms or condition as you.

- Things to be careful of

Avoid alcohol consumption during the course of treatment with Neurobine F.C. Tablet. Avoid driving a car or operating potentially dangerous machinery after taking Neurobine F.C. Tablet.

Side effects

Incidence rare; Thiamine can cause anaphylactic reaction (coughing, difficulty in swallowing, hives, itching of skin, swelling of face, lips, or eyelids, wheezing or difficulty in breathing) after large dose is administered through veins. Pyridoxine in high dose taken for several months can cause severe damage of body's sensory nerves. Cyanocobalamin can cause anaphylactic reaction (skin rash, itching, wheezing).

You may report any side effects or adverse drug reactions directly to the National Centre for Adverse Drug Reaction Monitoring by calling Tel: 03-78835550, or visiting the website npra.moh.gov.my (Public → Reporting Side Effects to Medicines (ConSERF) or Vaccines (AEFI)).

Storage and Disposal of Neurobine**F.C. Tablet****- Storage**

Store below 30°C, protect from light & moisture.


- Disposal

Neurobine F.C. Tablet should be disposed when your doctor or pharmacist tells you to stop taking this tablet or it has passed the expiry date.

Ask your pharmacist how to dispose medicines that are no longer required.

Product Description**- What it looks like**

A reddish pink colour film coated round tablet, one side impressed with a

mark ,

- Ingredients

- Active ingredient(s)
Thiamine Mononitrate
Pyridoxine HCl
Cyanocobalamin

- Inactive ingredients

Lactose
Cellulose
Magnesium Stearate
Magnesium Aluminium Silicate
Sodium Starch Glycolate
Colorant

MAL number

MAL07082983XZ

Manufacturer

Y.S.P. Industries (M) Sdn. Bhd.
Lot 3, 5 & 7, Jalan P/7, Section 13,
Kawasan Perindustrian Bandar Baru
Bangi, 43000 Kajang, Selangor Darul
Ehsan, Malaysia.

Product Registration Holder

Y.S.P. Industries (M) Sdn. Bhd.
Lot 3, 5 & 7, Jalan P/7, Section 13,
Kawasan Perindustrian Bandar Baru
Bangi, 43000 Kajang, Selangor Darul
Ehsan, Malaysia.

Date of revision:

23/07/2018

Serial Number

NPRA(R4/1)121217/00355