

AXCEL ERYTHROMYCIN ES-200 SUSPENSION

Erythromycin (200mg/5ml)

What is in this leaflet

1. What Axcel Erythromycin ES-200 Suspension is used for.
2. How Axcel Erythromycin ES-200 Suspension works.
3. Before you use Axcel Erythromycin ES-200 Suspension.
4. How to use Axcel Erythromycin ES-200 Suspension.
5. While you are using Axcel Erythromycin ES-200 suspension.
6. Side effects.
7. Storage and disposal of Axcel Erythromycin ES-200 Suspension.
8. Product description.
9. Manufacturer and Product Registration Holder.
10. Date of revision.

What Axcel Erythromycin ES-200 Suspension is used for

Axcel Erythromycin ES-200 Suspension is indicated in the treatment of infections caused by susceptible organisms in the diseases listed below: upper and lower respiratory tract infections, pertussis (*whooping cough*), skin and soft tissue infections, otitis media, erythrasma (skin disease), listeriosis, diphtheria, primary syphilis (sexually transmitted infection) and prophylaxis of infection of the heart.

How Axcel Erythromycin ES-200 Suspension works

Axcel Erythromycin ES-200 Suspension contains a substance called Erythromycin. Erythromycin is an antibiotic that belongs to the group of medicines called macrolides. These medicines work by killing or stopping the growth of bacteria, which cause infections.

Before you use Axcel Erythromycin ES-200 Suspension

-When you must not use it

Do not take Axcel Erythromycin ES-200 Suspension if:

- You have ever had an allergic reaction to medicines containing erythromycin or to other antibiotics from the macrolide family. Some of the symptoms

of an allergic reaction may include rash, itching or hives, swelling of face, lips, tongue or other parts of the body, shortness of breath, wheezing or difficulty in breathing.

- You are allergic to any of the ingredients listed at the end of this leaflet.
- You have liver disease or liver impairment.

-Before you start to use it

Tell your doctor if:

- You are allergic to any medicines.
- You have impaired liver function.
- You have myasthenia gravis.
- You have history of arrhythmias.
- If you have diarrhea.
- You suffer from rhabdomyolysis (muscle injury) or kidney disease.
- Please consult your doctor before start to use Axcel Erythromycin ES-200 Suspension if you are pregnant, plan to become pregnant, breast-feeding or intend to breast-feed. Your doctor will have weighed the risks of you taking Axcel Erythromycin ES-200 Suspension against the benefits they expect it will have for you.

-Taking other medicines

Tell your doctor if you are taking any other medicines, including any that you buy without a prescription from a pharmacy, supermarket or health food shop as it may interact with your Axcel Erythromycin ES-200 Suspension such as:

- Medications causing liver or hearing injury
- Cimetidine, medicine for gastric
- Carbamazepine, phenytoin, Valproate, medicines for seizure
- Cyclosporine, medicine used for after organ transplant
- Alfentanil, pain killer
- Cisapride, medicine for stomach
- Lovastatin, medicine to lower the cholesterol
- Terfenadine, for allergic reaction
- Theophylline, for asthma

- Digoxin, medicine for heart
- Ergotamine, medicine for migraine
- Midazolam, medicine for anxiety

Your doctor or pharmacist may have more information on medicines to be careful with or to avoid while taking this medication.

How to use Axcel Erythromycin ES-200 Suspension

-How much to use

Adults: 400mg (10ml), 6 hourly or 800mg (20ml), 12 hourly.

Dosage may be increased up to 4g per day according to the severity of the infection.

Children 2 to 8 years: 1g (25ml) daily in divided doses.

Infant and children up to 2 years: 500 mg (12.5ml) daily in divided doses.

Direction for mixing:

Shake bottle well to loosen granules. Add some water to mix with the granules first and shake, later top up to the mark and replace with cap. Shake bottle well to mix medicine properly.

Follow all directions given to you by your doctor and pharmacist carefully. They may differ from the information contained in this leaflet. If you do not understand the instructions on the label, ask your doctor or pharmacist for help.

-When to use it

Your doctor will decide when it is appropriate for you to take Axcel Erythromycin ES-200 Suspension according to the type of infection you have.

-How long to use it

Continue taking Axcel Erythromycin ES-200 Suspension for as long as doctor recommends. Do not stop taking Axcel Erythromycin ES-200 Suspension, even if you feel better after a few days, unless advised by your doctor. Your infection may not clear completely if you stop taking your medicine too soon.

AXCEL ERYTHROMYCIN ES-200 SUSPENSION

Erythromycin (200mg/5ml)

-If you forget to use it

Consult your doctor or pharmacist on what you should do if you forget to use it.

Take the missed dose as soon as you remember. If it is almost time for your next dose, wait until then take the medicine and skip the missed dose. Do not take a double dose to make up for the missed dose.

-If you use too much (overdose)

- In case of overdosage, erythromycin should be discontinued.
- Contact your doctor immediately or go to the Emergency Department of your nearest hospital, if you think you or anyone else may have taken too much of this medicine. Do this even if there are no signs of discomfort or poisoning. You may need urgent medical attention.

While you are using it

-Things you must do

- Take your medicine exactly as your doctor has told you.
- Tell all the doctors, dentists and pharmacists treating you that you are taking Axcel Erythromycin ES-200 Suspension.
- Tell your doctor immediately if you become pregnant while talking this medication.
- If you are taking Axcel Erythromycin ES-200 Suspension for an infection and your symptoms do not improve within a few days, or if they become worse, tell your doctor.
- If you get severe diarrhea tell your doctor or pharmacist immediately.

-Things you must not do

- Do not stop taking the medicine unless advised by your doctor.
- Do not take any new medicines without consulting your doctor.
- Do not give Axcel Erythromycin ES-200 Suspension to anyone else, even if they have the same symptoms or condition as you.

-Things to be careful of

- Prolonged or repeated use of erythromycin may result in an overgrowth of nonsusceptible bacteria or fungi.
- Be careful driving or operating machinery until you know how Axcel Erythromycin ES-200 Suspension affects you.

Side effects

Like all medicines, Axcel Erythromycin ES-200 Suspension can cause side effects although not everybody gets them. The following side effects have been reported:

- Gastro-intestinal disturbances such as: stomach discomfort and cramp, nausea, vomiting, and diarrhea
- Pseudomembranous colitis (inflammation of large intestine), hepatotoxicity (liver damage), ototoxicity (ear poisoning).
- Allergic reactions ranging from urticaria (hives) to anaphylaxis (a severe, life-threatening allergic reactions to a foreign substance that affects various parts of the body).
- Ventricular arrhythmias (abnormal rapid heart rhythm) including ventricular tachycardia (rapid heartbeat).
- Agranulocytosis (severe reduction in the number of white blood cells), central neurotoxicity (damage to nervous tissue) including psychotic reactions (such as hallucination) and nightmares, myasthenia-like syndrome (weakness and fatigue of muscle) and pancreatitis (inflammation of pancreas).

Tell your doctor or pharmacist if you notice anything else that is making you feel unwell. Other side effects not listed above may also occur in some people.

You may report any side effects or adverse drug reactions directly to the National Centre for Adverse Drug Reaction Monitoring by calling Tel: 03-78835550, or visiting the website

portal.bpfk.gov.my (Consumers -> Reporting).

Storage and disposal of Axcel Erythromycin ES-200 Suspension

-Storage

Keep Axcel Erythromycin ES-200 Suspension out of reach of children. Keep container well closed. Store at room temperature not exceeding 30°C until reconstitution. After reconstituted, store below 25°C and use within 35 days. Refrigeration is not required.

-Disposal

Medicines should not be disposed of via wastewater or household waste. Ask your pharmacist how to dispose of medicines no longer required. These measures will help to protect the environment.

Product description

-What it looks like

Axcel Erythromycin ES-200 Suspension is a white colored granules which gives a yellow colour suspension with fruit flavour when reconstituted.

-Ingredients

Active Ingredient:

Erythromycin Ethylsuccinate

Inactive Ingredients:

Icing Sugar, Dibasic Sodium Phosphate, Sodium Saccharin, PVP K30, Dimethicone, Xanthan Gum, Polysorbate, Aerosil, Mixed Fruit Flavor, Quinoline Yellow.

-MAL number

MAL20000365AZ

Manufacturer and product registration holder

KOTRA PHARMA (M) SDN BHD
No. 1, 2 & 3, Jalan TTC 12,
Cheng Industrial Estate,
75250 Melaka, Malaysia.

Date of revision

19/11/2015

Serial Number

BPFK(R4/1)021115/00392