TRILON-E CREAM

Econazole Nitrate / Triamcinolone Acetonide (1% w/w / 0.1% w/w)

What is in this leaflet:

- 1. What Trilon-E Cream is used for
- 2. How Trilon-E Cream works
- 3. Before you use Trilon-E Cream
- 4. How to use Trilon-E Cream
- 5. While you are using Trilon-E Cream
- 6. Side Effects
- 7. Storage and Disposal of Trilon-E Cream
- 8. Product description
- 9. Manufacturer and Product Registration Holder
- 10. Date of Revision

What Trilon-E Cream is used for

It is a combination of Econazole Nitrate and Triamcinolone. The combination of the cream is indicated for the relief of the inflammatory manifestation of corticosteroid-responsive skin disease when complicated by secondary infection caused by fungi or organisms sensitive to econazole nitrate. The cream is indicated in cases of diaper rash, inflammation or infection of skin, groin or hair follicles caused by fungus.

Econazole nitrate is effective in the treatment of fungal infections such as: cutaneous candidiasis (fungal infection of the skin), pityriasis versicolor (causing depigmentation in affected skin), athlete's foot, jock itch and tinea corporis (ringworm of the body).

How Trilon-E Cream works

Trilon-E Cream helps to reduce inflammation (swelling, pain, redness, heat) and itchiness from actions of triamcinolone acetonide and helps to treat infections caused by fungal and gram-positive bacteria from the actions of econazole nitrate.

Before you use Trilon-E Cream

- When you must not use it
- If you are allergic (hypersensitive) to any ingredients of Trilon-E Cream.
- ♦ If you have viral diseases of the skins such as varicella and vaccina.

- Before you start to use it

You must tell your doctor if:

- you have allergies to any medicine which you have taken previously to treat your current condition.
- If you are pregnant, trying to get pregnant or breast feeding.

- Taking other medicines

Always inform your doctor and pharmacist if you are taking any other medicines, including herbal tonics, supplements and medicines that you buy without prescription.

How to use Trilon-E Cream

Always take this medicine exactly as your doctor has told you. You should check with your doctor or pharmacist if you are not sure. Do not take this medicine by mouth. This medicine is only for topical application.

- How much to use it

A thin film of the cream should be applied to cover completely the affected and surrounding skin areas twice daily, in the morning and at night.

- When to use it

The cream is used topically. The cream should be applied regularly for treatment to be effective.

- How long to use it

Duration of therapy varies and depends upon the extent and location of the disease.

- If you forget to use it

- If you forget a dose, take it as soon as you remember it.
- However, if it is nearly time for the next dose, skip the missed dose.
- ◆ Do not take a double dose to make up for a forgotten dose.

- If you take too much (Overdose)

Symptoms of toxicity are associated with the use of topical corticosteroids. Symptoms are described under 'side effects'. Simple withdrawal of the medicine is usually all that is necessary.

While you are using Trilon-E Cream

- Things you must do

Avoid applying the cream with an occlusive dressing to large areas of the body because of the increased risk of systemic toxicity.

Avoid using the cream in the presence of infection as there is a risk of sensitivity reaction occurring.

If you do become pregnant whilst taking Trilon-E Cream, tell your doctor.

- Things you must not do

Do not give Trilon-E Cream to anyone else, even if their symptoms seem similar or they have same condition as you. Your doctor has prescribed Trilon-E Cream for you and your condition.

- Things to be careful of

The cream should not be applied to ulcers of the legs.

Long term topical use is best avoided especially in children and during pregnancy.

Side Effects

Like all medicines, Trilon-E Cream can have side effects, although not everybody gets them. The following effects may occur:

- Signs of infection such as pain, redness or pus-containing blisters
- Signs of irritation such as burning, itching, blistering or peeling not present before therapy.

Prolonged usage may cause side effects such as:

- ♦ Moon face due to systemic toxicity
- Reddish purple lines on arms, legs, trunk or groin
- ♦ Thinning of skin with easy bruising
- Unusal increase in hair growth
- ♦ Unusual loss of hair

If any of the side effects gets serious, or if you notice any side effects not listed in this leaflet, please tell your doctor or pharmacist.

Consult your doctor or pharmacist if you are in doubt or for any further information.

You may report any side effects or adverse drug reactions directly to the National Centre for Adverse Drug Reaction Monitoring by calling Tel: 03-78835550, or visiting the website portal.bpfk.gov.my

(Consumers→Reporting)

Storage and Disposal of Trilon-E Cream

- Storage

Do not use after the expiry date stated on the tube end. The expiry date refers to the last day of that month.

Store below 30°C Protect from heat and freezing

Keep out of the reach and sight of children.

- <u>Disposal</u>

Medicines should not be disposed of via wastewater or household waste. Ask your pharmacist how to dispose of medicines no longer required. These measures will help to protect the environment.

Product Description

- What Trilon-E Cream looks like

A white cream base in printed aluminium tubes with white cap.

They are supplied in 15g collapsible

- Ingredients

- Active ingredients

aluminium tubes.

Each gram contains Econazole Nitrate 10mg and Triamcinolone Acetonide 1mg

- Inactive ingredients

The other ingredients are propylene glycol, mineral oil heavy, polyoxyl 40 stearate, stearic acid, cetyl alcohol and ethylparaben

- MAL number: MAL19910845A

Manufacturer and Product Registration Holder

Malaysian Pharmaceutical Industries Sdn. Bhd.

Plot 14, Lebuhraya Kampung Jawa, 11900 Bayan Lepas, Pulau Pinang, Malaysia.

Date of Revision

10/09/2015

Serial Number:

BPFK(R4/1)030915/00298