

BISOL F.C. TABLET

Each tablet contains: Bisoprolol Fumarate 5mg

What is in this leaflet:

1. What Bisol F.C. Tablet is used for
 2. How Bisol F.C. Tablet works
 3. Before you use Bisol F.C. Tablet
 4. How to take Bisol F.C. Tablet
 5. While you are using Bisol F.C. Tablet
 6. Side effects
 7. After using Bisol F.C. Tablet
 8. Product Description
 9. Manufacturer
-

What Bisol F.C. Tablet is used for

Bisol F.C. Tablet is used for the treatment of high blood pressure, coronary heart disease, heart failure in adjunction with other drugs.

How Bisol F.C. Tablet Works

The active ingredient in the medicine is 'Bisoprolol Fumarate'. This belongs to a group of medicines called 'beta blocker'. It primarily acts on the beta-1 receptor found in the heart, resulting in slower heart rate with little change observed in force of heart muscle contraction, thereby lowering blood pressure.

Before you use Bisol F.C. Tablet

When you must not take it

- If you are allergic to Bisoprolol Fumarate or any other ingredients in the product
- If you have heart block
- If you have markedly slow heart rate

Before you start to take it

You should check with your doctor:-

- If you are or think you may be pregnant
 - If you are planning to become pregnant
 - If you have kidney disease
 - If you have liver disease
 - If you have diabetes mellitus
 - If you have asthma
 - If you have history of heart failure and other cardiovascular diseases
 - If you plan or are undergoing surgery
 - If you have thyroid disease
-

Taking Other Medicines

There are a few medicines which may not mix with Bisol F.C. Tablet. It is important to tell your doctor or pharmacist about all the medicines that you are taking, including those obtained without doctor's prescription. You should talk to your doctor or pharmacist if you are on the following:-

- Other beta blockers
 - Other blood pressure lowering agents
 - Other medications for heart disease treatment
 - Antibiotics
 - Digoxin
 - Ulcer healing drugs
-

How to take Bisol F.C. Tablet

How much to Take

You should take Bisol F.C. Tablet exactly as advised by your doctor or pharmacist. The amount you take each day will depend on your condition. Do not take more or less than what your doctor prescribes.

The usual initial dose for adults is 2.5 to 5mg daily, gradually increased to maximum 20mg daily if required.

When to take

Bisol F.C. Tablet is to be taken in the morning after breakfast. Swallow the tablets with some water

How long to take it

It is important to take Bisol F.C. Tablet as long as your doctor prescribes.

If you forget to take it

Do not take extra tablets to make up for a missed dose. Just take your next dose at usual the time.

If you take too much (overdose)

If you take too many tablets by mistake, contact your doctor immediately or seek medical attention.

While you are using Bisol F.C. Tablet

Things you must do

Take Bisol F.C. Tablet exactly the way as prescribed.

Things you must not do

Do not share your medicines with others even if they have the same diagnosis as you

Things to be careful of

If you experience any side effects, consult your doctor or pharmacist for advice

Side Effects

The possible side effects are dizziness, headache, vivid dreams, gastrointestinal discomfort, skin rashes, decreased libido, general tiredness.

Please refer to your doctor or pharmacist if you experience any other symptoms while taking this medication.

After using Bisol F.C. Tablet

Storage

Store below 30°C and away from excess Heat and moisture.

Disposal

If you have any unwanted tablets, do not put them in waste water or household rubbish. Ask your pharmacist how to dispose of the tablets.

Product Description

What it looks like

Yellow, biconvex, heart-shaped, film-coated tablet with a ST on one side, score on another.
411

Ingredients

Each tablet contains Bisoprolol Fumarate 5mg as active ingredient.

Other non-active ingredients are:

Magnesium stearate, corn starch, sodium starch glycolate, polyethylene glycol 6000, hydroxypropyl methyl cellulose, dibasic calcium phosphate anhydrous, titanium dioxide, iron oxide yellow.

Product Registration Number

MAL10012459A

Manufacturer

Taiwan Biotech Co., Ltd.
22, Chieh-Shou Road,
Taoyuan,
Taiwan R.O.C.

Date of Revision
16 August 2011

Market Authorization Holder
Sunward Pharmaceutical Sdn. Bhd.

BISOL F.C. TABLET

Setiap tablet mengandungi: Bisoprolol fumarat
5mg

Kandungan

1. Kegunaan Bisol F.C. Tablet.
 2. Cara Bisol F.C. Tablet berfungsi.
 3. Perhatian sebelum mengambil Bisol F.C. Tablet.
 4. Cara pengambilan Bisol F.C. Tablet.
 5. Semasa pengambilan Bisol F.C. Tablet.
 6. Kesan sampingan.
 7. Selepas mengambil Bisol F.C. Tablet.
 8. Gambaran produk.
 9. Pengilang.
-

Kegunaan Bisol F.C. Tablet

Bisol F.C. Tablet diindikasikan untuk rawatan tekanan darah tinggi, penyakit jantung koronari, lemah jantung bersama ubatan jantung yang lain.

Cara Bisol F.C. Tablet Berfungsi

Bisoprolol fumarat tergolong dalam kumpulan ubatan penyekat beta. Ia bertindak pada reseptor beta-1 di jantung untuk menurunkan kadar/ beban jantung sekaligus menurunkan tekanan darah.

Perhatian Sebelum Mengambil Bisol F.C. Tablet

Jangan ambil jika

- Anda alergi kepada Bisoprolol fumarat atau bahan-bahan lain yang terkandung di dalam produk ini.
- Mengalami serangan jantung.
- Mempunyai kadar jantung yang rendah.

Sebelum mengambil

Dapatkan nasihat doktor jika anda:-

- Mengandung/ Disyaki mengandung.
- Merancang untuk mengandung.
- Menghidapi penyakit buah pinggang.
- Menghidapi masalah hati.
- Menghidapi penyakit kencing manis.
- Menghidap penyakit asma.
- Mempunyai sejarah kegagalan jantung dan lain-lain penyakit jantung.
- Merancang untuk menjalani pembedahan.
- Menghidapi penyakit tiroid.

Mengambil ubat lain

Terdapat beberapa jenis ubatan yang tidak digalakkan diambil bersama dengan Bisol F.C. Tablet. Anda harus memaklumkan doktor dan ahli farmasi anda mengenai ubatan yang sedang diambil termasuk ubatan yang dibeli di farmasi tanpa preskripsi doktor. Maklumkan doktor dan ahli farmasi jika anda sedang mengambil ubatan berikut:-

- Penyekat-beta yang lain
 - Ubat penurunan tekanan darah yang lain.
 - Ubat-ubatan lain untuk rawatan jantung
 - Antibiotik
 - Digoxin
 - Ubat rawatan ulser
-

Cara Mengambil Bisol F.C. Tablet

Arahan pengambilan

Ikut arahan doktor atau ahli farmasi anda. Jangan ambil lebih atau kurang daripada amaun yang disyorkan.

Dos permulaan biasa bagi dewasa adalah 2.5-5mg sehari. Dos harian maksima adalah 20mg.

Waktu pengambilan

Bisol F.C. Tablet perlu diambil selepas sarapan pagi.

Jangkamasa pengambilan

Ikut arahan yang telah ditetapkan oleh doktor atau ahli farmasi anda.

Jika terlupa mengambil

Jangan gandakan dos anda. Ambil dos seperti biasa bila tiba waktu untuk dos seterusnya.

Jika terlebih dos

Hubungi doktor anda atau dapatkan bantuan kecemasan.

Semasa Pengambilan Bisol F.C. Tablet

Perkara yang perlu dilakukan

Ambil Bisol F.C. Tablet mengikut arahan doktor.

Larangan

Jangan kongsi ubatan anda dengan orang lain walaupun mereka mengalami masalah yang sama.

Perhatian

Rujuk doktor atau ahli farmasi anda jika anda mengalami mana-mana kesan sampingan yang dinyatakan.

Kesan Sampingan

Kesan sampingan yang mungkin berlaku adalah pening, sakit kepala, gangguan tidur, gangguan gastrointestinal, masalah kulit, penurunan libido, penat.

Rujuk doktor atau ahli farmasi anda jika anda mengalami kesan-kesan lain semasa pengambilan ubat ini.

Selepas Mengambil Bisol F.C. Tablet

Penyimpanan

Simpan pada suhu dibawah 30°C. dijauhkan daripada suhu dan kelembapan yang berlebihan.

Pelupusan

Jangan buang ubatan yang berlebihan secara sebarangan. Dapatkan nasihat ahli farmasi anda mengenai cara-cara pelupusan ubatan..

Gambaran Produk

Rupa produk

Kuning, dwicembung, bentuk hati, tablet bersalut filem bertanda **ST** pada satu muka dan skor pada 144

muka yang bertentangan.

Kandungan

Setiap tablet mengandungi Bisoprolol fumarat 5mg sebagai bahan aktif.

Lain-lain kandungan bahan tidak aktif: Magnesium stearate, corn starch, sodium starch glycolate, polyethylene glycol 6000, hydroxypropyl methyl cellulose, dibasic calcium phosphate anhydrous, titanium dioxide, iron oxide yellow.

Market Authorization Holder
Sunward Pharmaceutical Sdn. Bhd.

Date of Revision
16 August 2011

Nombor Pendaftaran Produk

MAL10012459A

Pengilang

Taiwan Biotech Co., Ltd.
22, Chieh-Shou Road,
Taoyuan,
Taiwan R.O.C.

Date of Revision
16 August 2011

Market Authorization Holder
Sunward Pharmaceutical Sdn. Bhd.