

MELOMET CREAM

Mometasone Furoate (1mg)

What is in this leaflet

1. What Melomet Cream is used for
2. How Melomet Cream works
3. Before you use Melomet Cream
4. How to use Melomet Cream
5. While you are using it
6. Side effects
7. Storage and disposal of Melomet Cream
8. Product description
9. Manufacturer and Product Registration Holder
10. Date of revision

What Melomet Cream is used for

Melomet Cream is used for relief of inflammatory and itching manifestations of corticosteroid-responsive skin disorders.

How Melomet Cream works

Mometasone furoate, a synthetic corticosteroid, exhibits marked anti-inflammatory, anti-itchiness and blood vessel narrowing properties. It control the production of potent mediators of inflammation.

Before you use Melomet Cream

-When you must not use it

Do not use if you have known hypersensitivity to any of the components.

-Before you start to use it

Please refer to doctor or pharmacist for further information.

-Taking other medicines

No related information has been reported.

How to use Melomet Cream

-How much to use

Adults (including elderly) and Children :
Apply a thin film to the affected skin areas.

-When to use it

Apply once daily. Melomet Cream may be used in children 2 years of age or older.

-How long to use it

Apply until the lesion heals or for duration of 3 weeks whichever is sooner or refer to doctor or pharmacist.

-If you forget to use it

If you miss a dose of Melomet Cream, take it as soon as possible. If it is approaching time for the next dose, skip the missed dose and go back to your regular dosing schedule. Do not double doses.

-If you use it too much (overdose)

Excessive, prolonged use of topical corticosteroids can suppress adrenal gland function resulting in secondary hormone insufficiency.

While you are using Melomet Cream

-Things you must do

If irritation or sensitization develops with the use of Melomet Cream, treatment should be withdrawn and appropriate therapy instituted.

Should an infection develop, use of an appropriate antifungal or antibacterial agent should be instituted. If a favorable response does not occur promptly, the corticosteroid should be discontinued until the infection is adequately controlled.

Do not use Melomet Cream at the large surface area or using occlusive technique. It will increased the systemic absorption of topical corticosteroid.

Use of topical corticosteroids in children should be limited to at least amount compatible with an effective therapeutic regimen and suitable precaution should be taken. Child have a larger skin surface area to body weight ration as compared to adults. Chronic corticosteroid therapy may interfere with growth and development of children.

-Things you must not do

Local and systemic toxicity is common especially following long continued use on large areas of damaged skin. Long term continuous therapy should be avoided in all individuals irrespective of age.

Melomet Cream is not for use in the eyes.

-Things to be careful of

Medicine of this class should not be used on pregnant women in large amounts or for prolonged periods of time.

It only should be used during pregnancy only if the potential benefit justifies potential risk to the fetus. Caution should also be exercised when it is administered to a nursing mother as it is not known that whether topical administration of corticosteroids could produce detectable quantities in breast milk.

Side effects

Very rarely local adverse reactions are reported include tingling sensation, itching and signs of skin thinning. Use of occlusive dressings causes infrequent local

adverse reactions include irritation, dryness, hair follicles inflammation, excessive hair growth, acne eruptions, lightening of the skin, tiny rashes around the mouth, secondary infection, stretch mark and heat rashes.

You may report any side effects or adverse drug reactions directly to the National Centre for Adverse Drug Reaction Monitoring by calling Tel: 03-78835550, or visiting the website portal.bpfk.gov.my

(Consumers→Reporting)

Storage and Disposal of MELOMET CREAM

-Storage

Keep in airtight container. Store at temperature below 30°C, protect from light & moisture.

-Disposal

If your doctor or pharmacist tells you to stop using Melomet Cream or the batch has passed the expiry date, ask your doctor or pharmacist what to do with any that is left over.

Product description

-What it looks like

A white colour cream.

-Ingredients

-Active ingredient:

Mometasone Furoate

-Inactive ingredient:

Propylene Glycol

Stearic Acid

Cetareth-6

Cetaryl Alcohol

Stearyl Alcohol

Sodium Lauryl Sulphate

White Petrolatum

Potassium Phosphate Monobasic

Titanium Dioxide

Mineral Oil

Phosphoric Acid

Methyl paraben

Propyl paraben

-MAL number

MAL07122846AZ

Manufacturer

Y.S.P. Industries (M) Sdn. Bhd.
Lot 3, 5 & 7, Jalan P/7, Section 13,
Kawasan Perindustrian Bandar Baru
Bangi, 43000 Kajang, Selangor Darul
Ehsan, Malaysia.

Product Registration Holder

Y.S.P. Industries (M) Sdn. Bhd.
Lot 3, 5 & 7, Jalan P/7, Section 13, Kawasan
Perindustrian Bandar Baru Bangi, 43000
Kajang, Selangor Darul Ehsan, Malaysia.

Date of revision

18/3/2016

Serial Number

BPFK(R4/1)180316/00080