

# MOTIGUT SUSPENSION

Domperidone (1 mg/ml)

## What is in this leaflet

1. What *Motigut Suspension* is used for
2. How *Motigut Suspension* works
3. Before you use *Motigut Suspension*
4. How to use *Motigut Suspension*
5. While you are using it
6. Side effects
7. Storage and Disposal of *Motigut Suspension*
8. Product Description
9. Manufacturer and Product Registration Holder
10. Date of revision

## What *Motigut Suspension* is used for

Indigestion associated with gastric reflux or inflammation of oesophagus, stomach fullness, abdominal distension, abdominal pain, bloating, flatulence, nausea and vomiting, or heart burn.

## How *Motigut Suspension* works

*Motigut suspension* contains domperidone with anti-emetic (stop vomiting) properties. It also increase the gastric emptying.

## Before you use *Motigut Suspension*

### - When you must not use it

Do not use *Motigut suspension* if:

- You are allergic to domperidone or any of the ingredients in this medication.
- You have gastric bleeding, blocking or holes.
- You have prolactin-releasing pituitary tumour.
- You have sorbitol intolerance

### *Pregnancy and lactation*

Do not take *Motigut Suspension* if you are pregnant, trying to get pregnant or think you may be pregnant.

Do not take *Motigut Suspension* if you are breast-feeding. Ask your doctor or pharmacist for advice before taking any medicine.

### - Before you start to use it

Tell your doctor or pharmacist if:

- It is for infant or small children
- You have liver problems
- You have kidney problems
- You have heart disease
- You have low or high potassium level or low magnesium level.

### - Taking other medicines

Tell your doctor if you are taking any other medicines, including any that you buy without a prescription from a pharmacy, supermarket or health food shop.

It is especially important to tell your doctor or pharmacist if you are taking any of the following:

- CYP3A4 inhibitors (eg: ketoconazole (antifungal), ritonavir (antiviral) and erythromycin (antibiotic))
- Medicines that affect the ECG results

## How to use *Motigut Suspension*

### - How much to use

Follow all directions given to you by your doctor and pharmacist carefully. They may differ from the information contained in this leaflet. If you do not understand the instructions on the label, ask your doctor or pharmacist for help.

Adults and adolescents  $\geq 12$  years and children weighing  $\geq 35$  kg: 10ml

Neonates, Infants and children  $<12$  years of age, adults and adolescents weighing  $<35$  kg: 0.25 mg/kg

### - When to use it

To be taken 3-4 times a day or as directed by your doctor or pharmacist.

For people with kidney problems, the dosing frequency of *Motigut* should be reduced to once or twice daily.

It is recommended to take *Motigut* 15-30 minutes before meals.

### - How long to use it

Continue taking *Motigut Suspension* for as long as your doctor recommends.

### - If you forget to use it

Consult your doctor or pharmacist on what you should do if you forget to use it.

Take the missed dose as soon as you remember. If it is almost time for your next dose, wait until then to take the medicine and skip the missed dose. Do not take a double dose to make up for the missed dose.

### - If you use too much (overdose)

Contact your doctor immediately or go to the Emergency Department of your nearest hospital, if you think you or anyone else may have taken too much of this medicine. Do this even if there are no signs of discomfort or poisoning. You may need urgent medical attention.

Taking too many tablets may cause drowsiness, disorientation and extrapyramidal reaction (involuntary muscle movements, loss of movement especially in children).

## While you are using it

### - Things you must do

Take your medicine exactly as your doctor has told you.

Tell all the doctors, dentists and pharmacists treating you that you are taking *Motigut Suspension*.

Tell your doctor immediately if you become pregnant while taking this medication.

### - Things you must not do

Do not stop taking the medicine unless advised by your doctor.

Do not take any new medicines without consulting your doctor or pharmacist.

Do not give *Motigut Suspension* to anyone else, even if they have the same symptoms or condition as you.

### - Things to be careful of

#### Driving and using machines

This medicine may affect your ability to drive or use machines. If the tablets make you feel sick, dizzy or tired, or give you a headache, do not drive or use machines and contact your doctor immediately.

The oral suspension contains sorbitol and may be unsuitable for people with sorbitol intolerance.

Stop taking domperidone and consult your physician if you have irregular heart beat

### Side effects

Like all medicines, *Motigut Suspension* can cause side effects, although not everybody gets them.

Visit your doctor or pharmacist immediately if you experience any side effects after taking this medicine.

Side effects may include allergic reaction, increase milk production, extrapyramidal effects, transient intestinal cramps, hives, breast enlargement, absence of menstrual period and others.

Very rare side effects include abnormal heartbeat, ECG changes, and sudden loss of heart function

You may report any side effects or adverse drug reactions directly to the National Centre for Adverse Drug Reaction Monitoring by calling Tel: 03-78835550, or visiting the website <http://npra.moh.gov.my/> (Public → Reporting Side Effects to Medicines (ConSERF) or Vaccines (AEFI))

### Storage and Disposal of *Motigut*

#### *Suspension*

#### - Storage

Keep out of the reach and sight of children.

Store below 30 °C.

#### - Disposal

Medicines should not be disposed of via wastewater or household waste. Ask your pharmacist how to dispose of medicines no longer required. These measures will help to protect the environment.

### Product Description

#### - What it looks like

A white raspberry-banana flavored suspension having sweet taste, filled and sealed in amber PET bottle with white color SQUARE printed cap.

#### - Ingredients

- Active ingredient  
Domperidone

#### - Inactive ingredients

Sorbitol liquid non crystallizable  
Dispersible cellulose (contains microcrystalline cellulose and carmellose sodium)  
Methyl p-hydroxybenzoate  
Propyl hydroxybenzoate  
Glycerol  
Sodium saccharin  
Polyxyl 40 hydrogenated castor oil  
Banana liquid  
Raspberry liquid  
Purified water

#### - MAL number:

MAL10120032AZ

#### **Manufacturer**

Square Pharmaceuticals Ltd.  
Square Road, Salgaria,  
Pabna, Bangladesh

#### **Product Registration Holder**

Medispec (M) Sdn Bhd  
55 & 57, Lorong Sempadan 2  
11400 Ayer Itam, Penang, Malaysia

#### **Date of revision**

22.2.2018

#### **Serial number**

NPRA(R4/1)240118/00022