

COVERAM

Perindopril arginine / amlodipine

COVERAM 5mg / 5mg tablet
COVERAM 10mg / 5mg tablet
COVERAM 5mg / 10mg tablet
COVERAM 10mg / 10mg tablet

What is in this leaflet

1. What COVERAM is used for
2. How COVERAM works
3. Before you use COVERAM
4. How to take COVERAM
5. While you are using COVERAM
6. Possible side effects
7. After using COVERAM
8. Product description

What COVERAM is and what it is used for

COVERAM is prescribed for treatment of high blood pressure (hypertension) and/or treatment of stable coronary artery disease (a condition where the blood supply to the heart is reduced or blocked).

Patients already taking perindopril and amlodipine from separate tablets may instead receive one tablet of COVERAM which contains both ingredients.

How COVERAM works

COVERAM is a combination of two active ingredients, perindopril and amlodipine.

Perindopril is an ACE (angiotensin converting enzyme) inhibitor. Amlodipine is a calcium antagonist (which belongs to a class of medicines called dihydropyridines). Together they work to widen and relax the blood vessels so that blood passes through them more easily and makes it easier for your heart to maintain a good blood flow.

Before you use COVERAM

Do not take COVERAM

- if you are allergic (hypersensitive) to perindopril or any other ACE inhibitor, or to amlodipine or any other dihydropyridines, or any of

the other ingredients of COVERAM,

- if you are more than 3 months pregnant (It is also better to avoid COVERAM in early pregnancy – see pregnancy section),
- if you have experienced symptoms such as wheezing, swelling of the face or tongue, intense itching or severe skin rashes with previous ACE inhibitor treatment or if you or a member of your family have had these symptoms in any other circumstances (a condition called angioedema),
- if you have cardiogenic shock (when the heart is unable to supply sufficient blood to the body), aortic stenosis (narrowing of the main blood vessels leading from the heart) or unstable angina (chest pain that may occur when resting),
- if you have severe hypotension (very low blood pressure),
- if you have heart failure after acute myocardial infarction, during first 28 days.

Take special care with COVERAM

If you have any of the following please talk to your doctor before taking COVERAM:

- hypertrophic cardiomyopathy (cardiac muscle disease) or renal artery stenosis (narrowing of the artery which supplies the kidney with blood),
- any other heart problems,
- liver problems,
- kidney problems or if you are receiving dialysis,
- collagen vascular disease (disease of the connective tissue) such as systemic lupus erythematosus or scleroderma,

- diabetes,
- if you are on a salt restricted diet or use salt substitutes which contain potassium (a well balanced potassium blood level is essential).

You must tell your doctor if you think you are (or might become) pregnant. COVERAM is not recommended in early pregnancy, and must not be taken if you are more than 3 months pregnant, as it may cause serious harm to your baby if used at that stage (see pregnancy section).

When you are taking COVERAM, you should also inform your doctor or the medical staff if you:

- are going to have a general anaesthetic and/or major surgery,
- have recently suffered from diarrhoea or vomiting (being sick),
- are to undergo LDL apheresis (the removal of cholesterol from your blood by a machine),
- are going to have desensitisation treatment to reduce the effects of an allergy to bee or wasp stings.

COVERAM is not recommended for use in children and adolescents.

Taking other medicines

Please tell your doctor or pharmacist if you are taking or have recently taken any other medicines, including medicines obtained without a prescription.

You should avoid COVERAM with:

- lithium (used to treat mania or depression),

- estramustine (used in cancer therapy),
- potassium-sparing diuretics (spironolactone, triamterene), potassium supplements or potassium-containing salt substitutes.

Treatment with COVERAM can be affected by other medicines. Make sure to tell your doctor if you are taking any of the following medicines as special care may be required:

- other medicines for high blood pressure, including diuretics (medicines which increase the amount of urine produced by the kidneys),
- non-steroidal anti-inflammatory drugs (e.g. ibuprofen) for pain relief or high dose aspirin,
- medicines to treat diabetes (such as insulin),
- medicines to treat mental disorders such as depression, anxiety, schizophrenia etc (e.g. tricyclic antidepressants, antipsychotics, imipramine-like antidepressants, neuroleptics),
- immunosuppressants (medicines which reduce the defence mechanism of the body) used for the treatment of auto-immune disorders or following transplant surgery (e.g. ciclosporin),
- allopurinol (for the treatment of gout),
- procainamide (for the treatment of an irregular heart beat),
- vasodilators including nitrates (products that widen the blood vessels),
- heparin (medicines used to thin blood),
- ephedrine, noradrenaline or adrenaline (medicines used to treat low blood pressure, shock or asthma),
- baclofen or dantrolene (infusion) both used to treat muscle stiffness in diseases such as multiple sclerosis; dantrolene is also used to treat malignant hyperthermia during anaesthesia (symptoms including very

- high fever and muscle stiffness),
- some antibiotics such as rifampicin,
- antiepileptic agents such as carbamazepine, phenobarbital, phenytoin, fosphenytoin, primidone,
- itraconazole, ketoconazole (medicines used for treatment of fungal infections),
- alpha-blockers used for the treatment of enlarged prostate such as prazosin, alfuzosin, doxazosin, tamsulosin, terazosin,
- amifostine (used to prevent or reduce side effects caused by other medicines or radiation therapy that are used to treat cancer),
- corticosteroids (used to treat various conditions including severe asthma and rheumatoid arthritis),
- gold salts, especially with intravenous administration (used to treat symptoms of rheumatoid arthritis).

How to take COVERAM

Always take COVERAM exactly as your doctor has told you. You should check with your doctor or pharmacist if you are not sure. Swallow your tablet with a glass of water, preferably at the same time each day, in the morning, before a meal. Your doctor will decide on the correct dose for you. This will normally be one tablet per day.

COVERAM will usually be prescribed for patients already taking perindopril and amlodipine from separate tablets.

If you take more COVERAM than you should

If you take too many tablets, contact your nearest accident and emergency department or tell your doctor immediately. The most likely effect in case of overdose is low blood pressure which can make you feel dizzy or faint. If this happens, lying down with your legs raised can help.

If you forget to take COVERAM

It is important to take your medicine every day as regular treatment works better. However, if you forget to take a dose of COVERAM, take the next dose at the usual time. Do not take a double dose to make up for a forgotten dose.

If you stop taking COVERAM

As the treatment with COVERAM is usually life-long, you should discuss with your doctor before you stop taking your tablets.

If you have any further questions on the use of this product, ask your doctor or pharmacist.

While you are using COVERAM

Taking COVERAM with food and drink

COVERAM should be taken before a meal.

Pregnancy

You must tell your doctor if you think you are (or might become) pregnant. Your doctor will normally advise you to stop taking COVERAM before you become pregnant or as soon as you know you are pregnant and will advise you to take another medicine instead of COVERAM. COVERAM is not recommended in early pregnancy, and must not be taken when more than 3 months pregnant, as it may cause serious harm to your baby if used after the third month of pregnancy.

Breastfeeding

Tell your doctor if you are breast-feeding or about to start breast-feeding. COVERAM is not recommended for mothers who are breast-feeding, and your doctor may choose another treatment for you if you wish to breast-feed, especially if your baby is newborn, or was born prematurely.

Driving and using machines

COVERAM does not affect alertness but you might experience dizziness or weakness due to low blood pressure which could affect your ability to drive or operate machinery.

Important information about some of the ingredients of COVERAM

COVERAM contains lactose monohydrate (a type of sugar). If you have been told by your doctor that you have an intolerance to some sugars, contact your doctor before taking this medicinal product.

Possible side effects

Like all medicines, COVERAM can cause side effects, although not everybody gets them.

If you experience any of the following, stop taking the medicinal product at once and tell your doctor immediately:

- swelling of the face, lips, mouth, tongue or throat, difficulty in breathing,
- severe dizziness or fainting,
- unusual fast or irregular heart beat.

Other side effects include:

- Common side effects (occur in less than 1 in 10 users but in more than 1 in 100 users): headache, dizziness, vertigo, pins and needles, somnolence (sleepiness), vision disturbances, tinnitus (sensation of noises in the ears), palpitations (very fast heartbeat), flushing (hot or warm feeling in your face), light-headedness due to low blood pressure, cough, shortness of breath, nausea (feeling sick), vomiting (being sick), abdominal pain, taste disturbances, dyspepsia or difficulty of digestion, diarrhoea, constipation, allergic reactions (such as skin rashes, itching), muscle cramps, feeling of tiredness, oedema (swelling of your legs or ankles),

- Uncommon side effects (occur in less than 1 in 100 users but in more than 1 in 1000 users): mood swings, sleep disturbances, trembling, syncope (temporary loss of consciousness), loss of pain sensation, rhinitis (blocked up or runny nose), changed bowel habits, hair loss, red or discoloured patches on skin, back, muscle or joint pain, chest pain, increased need to urinate especially during the night, malaise, bronchospasm (tightening of the chest, wheezing and shortness of breath), dry mouth, angioedema (symptoms such as wheezing, swelling of the face or tongue), kidney problems, impotence, increased sweating, breast enlargement in men, weight increase or decrease,
- Very rare side effects (occur in less than 1 in 10,000 users): confusion, cardiovascular disorders (irregular heart beat, angina, heart attack and stroke), eosinophilic pneumonia (a rare type of pneumonia), erythema multiforme (a skin rash which often starts with red itchy patches on your face, arms or legs), disorders of the blood, pancreas, stomach or liver, peripheral neuropathy (disease that produces loss of sensations, pain, inability to control muscles), hypertonia (abnormal increase in muscle tension), swelling of the gums, high blood sugar.
- The following side effects have also been reported by patients taking COVERAM: hypoglycaemia (very low blood sugar level), vasculitis (inflammation of blood vessels), tremor and rigidity.

If any of the side effects get serious, or if you notice any side effects not listed in this leaflet, please tell your doctor or pharmacist.

After using COVERAM

Storage

Keep out of the reach and sight of children.

Do not use COVERAM after the expiry date which is stated on the carton and bottle. The expiry date refers to the last day of that month.

Keep the bottle tightly closed in order to protect from moisture. Store in the original package.

Disposal

Medicines should not be disposed of via wastewater or household waste. Ask your pharmacist how to dispose of medicines no longer required. These measures will help to protect the environment.

Product description

What COVERAM contains

The active substances are perindopril arginine and amlodipine.

COVERAM 5mg / 5mg: one tablet contains 5 mg perindopril arginine and 5 mg amlodipine.


COVERAM 10mg / 5mg: one tablet contains 10 mg perindopril arginine and 5 mg amlodipine.


COVERAM 5mg / 10mg: one tablet contains 5 mg perindopril arginine and 10 mg amlodipine.

COVERAM 10mg / 10mg: one tablet contains 10 mg perindopril arginine and 10 mg amlodipine.

The other ingredients in the tablet are: lactose monohydrate, magnesium stearate (E470B), cellulose microcrystalline (E460), colloidal anhydrous silica (E551).

What COVERAM looks like and contents of the pack

COVERAM 5mg / 5mg tablets are white, rod-shaped tablets engraved with 5/5 on one face and  on the other face.

COVERAM 10mg / 5mg tablets are white, triangular-shaped tablets engraved with 10/5 on one face and  on the other.

COVERAM 5mg / 10mg tablets are white, square-shaped tablets

engraved with 5/10 on one face and *☞ on the other.

COVERAM 10mg / 10mg tablets are white, round tablets engraved with 10/10 on one face and *☞ on the other.

Registration numbers

COVERAM 5mg / 5mg:
MAL20091910A
COVERAM 10mg / 5mg:
MAL20091911A
COVERAM 5mg / 10mg:
MAL20091912A
COVERAM 10mg / 10mg:
MAL20091913A

The tablets are available in containers of 30 tablets.

Manufacturer

Servier (Ireland) Industries Ltd
Gorey Road
Arklow - Co. Wicklow – Ireland

Product Licence Holder

Servier Malaysia Sdn Bhd
1301 Level 13, Uptown 2
No 2 Jalan SS 21/37,
Damansara Uptown
47400 Petaling Jaya
Selangor Darul Ehsan

Date of revision: 07.07.2010

COVERAM

Perindopril arginine / amlodipine

COVERAM 5mg / 5mg tablets

COVERAM 10mg / 5mg tablets

COVERAM 5mg / 10mg tablets

COVERAM 10mg / 10mg tablets

Dalam sisip kandungan ini

1. Apa itu COVERAM dan kegunaannya
2. Bagaimana COVERAM bertindak
3. Sebelum mengambil COVERAM
4. Cara pengambilan COVERAM
5. Semasa mengambil COVERAM
6. Kesan sampingan yang mungkin berlaku
7. Selepas menggunakan COVERAM
8. Maklumat produk

Apa itu COVERAM dan kegunaannya

COVERAM digunakan untuk rawatan tekanan darah tinggi dan / atau rawatan penyakit arteri koronari stabil (di mana bekalan darah ke jantung telah berkurangan atau terhalang). Pesakit yang sedang mengambil tablet perindopril dan amlodipine berasingan boleh mengambil sebiji tablet COVERAM yang mengandungi dua bahan tersebut.

Bagaimana COVERAM bertindak

COVERAM adalah kombinasi dua bahan aktif iaitu perindopril dan amlodipine. Perindopril adalah angiotensin pengubah enzim (ACE) inhibitor. Amlodipine ialah calcium antagonist (dimana ia tergolong dalam kelas ubat yang dipanggil dihydropyridines). Kedua-dua bahan ini bertindak dengan membesarkan saluran darah, seterusnya memudahkan jantung anda mengepam darah didalamnya.

Sebelum mengambil COVERAM

Jangan mengambil COVERAM

- jika anda alah (hipersensitif) terhadap perindopril atau

ACE inhibitor yang lain atau amlodipine atau mana-mana dihydropyridines lain atau sebarang kandungan di dalam COVERAM,

- Jika anda telah mengandungi lebih daripada 3 bulan. (Adalah lebih baik mengelak COVERAM dalam awal kehamilan – lihat seksyen kehamilan),
- jika anda telah mengalami simptom seperti kesukaran bernafas, bengkak muka, lidah atau kerongkong, gatal yang teruk atau alahan kulit yang teruk dengan rawatan ACE inhibitor yang sebelumnya atau ahli keluarga anda mempunyai simptom tersebut (keadaan yang dipanggil 'angioedema'),
- jika anda mempunyai renjatan kardiogenik (apabila jantung tidak larat membekalkan darah yang secukupnya untuk badan), stenosis aorta (penguncupan saluran darah utama yang keluar dari jantung) atau angina yang tidak stabil (sakit dada apabila berehat),
- jika anda mempunyai hypotension teruk (tekanan darah sangat rendah),
- jika anda mengalami kegagalan jantung selepas acute myocardial infarction, ketika 28 hari pertama.

Ambil perhatian yang khusus dengan COVERAM

Sila berbincang dengan doktor anda jika perkara yang berikut berkaitan dengan anda:

- hypertrophic cardiomyopathy (penyakit otot jantung) atau stenosis arteri ginjal (penguncupan salur darah yang menyalurkan darah ke ginjal),

- jika anda ada masalah jantung lain,
- jika anda ada masalah hati,
- jika anda ada masalah buah pinggang atau menerima rawatan dialisis,
- jika anda mengalami penyakit kolagen vaskular (penyakit tisu penghubung) seperti systemic lupus erythematosus atau scleroderma,
- jika anda mempunyai penyakit kencing manis,
- jika anda sedang dalam diet garam terkawal atau menggunakan pengganti garam yang mengandungi potassium.

Anda mesti memberitahu doktor anda jika anda rasa (atau mungkin) mengandungi COVERAM tidak disarankan semasa mengandung trimester pertama dan ia boleh menyebabkan kemudaratan yang serius selepas 3 bulan mengandung.

Anda juga perlu beritahu doktor anda:

- jika anda akan melalui proses anaesthesia atau pembedahan,
- jika anda akan melalui proses 'LDL apheresis' (pembuangan kolesterol dari saluran darah menggunakan mesin),
- jika anda akan mendapat rawatan untuk menjadikan anda kurang sensitif terhadap kesan alahan lebah atau sengatan tebu,
- jika anda mengalami cirit-birit atau muntah baru-baru ini atau dinyahhidratkan,
- jika anda telah diberitahu oleh doktor anda yang anda tidak mempunyai toleransi keatas sesetengah gula,

COVERAM tidak disarankan untuk kegunaan dalam kanak-kanak dan remaja.

Pengambilan ubat lain

Sila beritahu doktor atau ahli farmasi anda jika anda akan atau telah mengambil mana-mana ubat-ubatan lain termasuk ubat yang tidak memerlukan preskripsi doktor.

Anda harus mengelak COVERAM dengan:

- lithium (digunakan untuk merawat mania atau kemurungan),
- estramustine (digunakan dalam terapi kanser),
- diuretik jenis 'potassium-sparing' (spironolactone, triamterene), potassium suplemen atau potassium yang mengandungi pengganti garam,

Rawatan dengan COVERAM boleh diganggu oleh ubat-ubatan lain.

Sila beritahu doktor jika anda akan atau telah mengambil ubat-ubatan lain yang kerana perhatian yang khusus mungkin diperlukan:

- ubat lain termasuk diuretik untuk penyakit tekanan darah tinggi (ubat yang meningkat jumlah air kencing yang dihasilkan oleh buah pinggang),
- ubat 'non-steroid' anti-radang (contohnya ibuprofen) untuk melegakan kesakitan atau dos aspirin yang tinggi,
- ubat untuk rawatan penyakit kencing manis (seperti insulin),
- ubat untuk rawatan penyakit mental seperti kemurungan, keresahan, schizophrenia (contohnya tricyclic anti-depressants, antipsychotics dan anti-depressants seperti imipramina, neuroleptics),
- ubat immunosuppressant (ubat yang mengurangkan mekanisme penghalang badan) yang digunakan untuk rawatan penyakit auto imuns atau selepas pembedahan transplant (contohnya ciclosporin),
- allopurinol (untuk rawatan gout),

- procainamide (untuk rawatan degupan jantung yang tidak normal),
- vasodilators termasuk nitrates (produk yang melebarkan saluran darah),
- heparin (ubat yang digunakan untuk mencairkan darah),
- ubat yang digunakan untuk rawatan tekanan darah rendah, kejutan atau asma (contohnya ephedrine, noradrenaline atau adrenaline).
- baclofen atau dantrolene (infusion) kedua-dua ubat ini digunakan untuk merawat kekakuan otot seperti penyakit multiple sclerosis; dantrolene juga digunakan untuk merawat malignant hyperthermia semasa anaesthesia (termasuk simptom demam yang sangat tinggi dan kekakuan otot),
- beberapa jenis antibiotik seperti rifampisin,
- ejen antiepileptic seperti carbamazepine, phenobarbital, phenytoin, fosphenytoin, primidone,
- itraconazole, ketoconazole (ubat yang digunakan untuk rawatan jangkitan kulat),
- alfa-blockers seperti prazosin, alfuzosin, doxazosin, tamsulosin, terazosin yang digunakan untuk rawatan prostate yang membesar,
- amifostine (digunakan untuk mencegah atau mengurangkan kesan sampingan yang disebabkan oleh ubat-ubatan lain atau terapi radiasi bagi rawatan kanser),
- corticosteroids (digunakan untuk merawat pelbagai keadaan termasuk asma yang teruk dan rheumatoid arthritis),
- gold salts, melalui pemberian intravena (digunakan untuk merawat simptom rheumatoid arthritis).

Cara pengambilan COVERAM

Ambil COVERAM seperti yang diarahkan oleh doktor anda. Anda perlu hubungi mereka jika anda tidak pasti. Telah tablet dengan segelas air sebaiknya pada sebelah

pagi sebelum makan. Doktor anda akan menetapkan dos yang betul bagi anda. Dos biasa adalah 1 tablet sehari.

COVERAM biasanya akan disyorkan kepada pesakit sudah pun mengambil perindopril dan amlodipine dari tablet berasingan.

Jika anda mengambil terlebih dos COVERAM daripada yang perlu

Jika anda mengambil terlalu banyak, hubungi bahagian kecemasan atau beritahu doktor anda secepat mungkin. Kesan terlebih dos biasanya adalah tekanan darah rendah dimana anda akan berasa pening atau lemah. Jika ini berlaku, baring dengan menaikkan kaki anda dapat membantu.

Jika anda terlupa mengambil COVERAM

Adalah penting anda mengambil ubat ini setiap hari seperti yang diarahkan untuk rawatan yang berkesan. Walau bagaimanapun, jika anda terlupa, ambil dos yang berikutnya pada masa yang sama. Jangan menggandakan dos untuk menggantikan dos yang terlupa itu.

Jika anda berhenti mengambil COVERAM

Oleh kerana rawatan dengan COVERAM adalah untuk sepanjang hayat, anda perlu berbincang dengan doktor sebelum berhenti mengambil ubat ini.

Jika anda ada sebarang pertanyaan berkaitan penggunaan ubat ini, sila bertanya doktor atau ahli farmasi anda.

Semasa mengambil COVERAM

Pengambilan COVERAM dengan makanan dan minuman

COVERAM harus diambil sebelum makan.

Mengandung

Anda mesti beritahu doktor anda jika anda rasa anda mengandung (atau akan mengandung). Doktor anda akan menasihatkan anda supaya berhenti mengambil COVERAM sebaik saja

mengetahui yang anda hamil. Ubat hipertensi yang sesuai mesti digantikan sebelum anda mula hamil. COVERAM tidak disarankan sewaktu awal kehamilan dan boleh menyebabkan kesan mudarat yang serius kepada kandungan jika diambil selepas 3 bulan kehamilan.

Menyusukan anak

Sila beritahu doktor anda jika anda menyusukan bayi atau ingin mula menyusukan bayi. Jangan ambil COVERAM jika anda menyusukan bayi. Doktor anda akan memberi rawatan lain yang bersesuaian sekiranya anda ingin menyusukan bayi anda terutamanya jika bayi yang baru lahir atau pra-matang.

Pemanduan dan penggunaan mesin

COVERAM tidak mempunyai kesan ke atas keterjagaan tetapi anda mungkin akan rasa pening dan lemah disebabkan tekanan darah rendah yang boleh terjadi ke atas sesetengah pesakit. Jika anda mendapat kesan ini, kebolehan anda untuk memandu dan mengendalikan mesin mungkin terjejas.

Informasi penting tentang kandungan COVERAM

COVERAM mengandungi monohidrat laktosa (sejenis gula). Jika anda telah diberitahu oleh doktor anda yang anda tidak mempunyai toleransi keatas sesetengah gula, hubungi doktor anda sebelum mengambil ubat ini.

Kesan sampingan yang mungkin berlaku

Seperti ubat-ubatan lain, COVERAM boleh menyebabkan kesan sampingan, walaupun tidak semua pesakit akan merasainya.

Jika anda mengalami sebarang kesan seperti dibawah ini, sila berhenti mengambil ubat dan maklumkan kepada doktor anda secepat mungkin:

- bengkak di bahagian muka, bibir-bibir, mulut, lidah atau tekak, kesukaran bernafas
- pening yang teruk atau pingsan,

- degupan jantung yang luar biasa cepat atau tidak tetap.

Diantara kesan sampingan yang mungkin terjadi (urutan menurun), kesan yang:

- Selalu berlaku (kurang dari 1 dalam setiap 10 pesakit tetapi lebih dari 1 dalam 100 pesakit): sakit kepala, pening, vertigo, pin & needles, gangguan penglihatan, tinnitus, kepala rasa ringan disebabkan tekanan darah yang rendah, batuk, sesak nafas, gangguan perut-usus (loya, muntah, sakit abdomen, gangguan rasa, mulut kering, dispepsia atau kesukaran penghadaman, cirit-birit, sembelit), reaksi alergi (seperti ruam kulit, gatal), kejang otot, rasa penat.
- Jarang berlaku (terjadi pada kurang dari 1 dalam 100 pesakit tetapi lebih dari 1 di dalam 1000 pesakit): mood swings, gangguan tidur, menggigil, syncope (hilang kesedaran sementara), kehilangan sensasi kesakitan, rhinitis (hidung tersumbat atau selsema), perubahan tabiat usus, keguguran rambut, kulit menjadi merah atau kelunturan warna pada kulit badan, sakit otot atau sendi, sakit dada, kencing yang kerap terutama pada waktu malam, dedar, bronkospasm (dada rasa ketat, wheezing, kesukaran bernafas), mulut kering, angioedema (simptom seperti wheezing, bengkak muka, lidah atau tekak, ruam kulit atau gatal yang teruk), masalah ginjal, impoten, berpeluh, pembesaran buah dada pada lelaki, penambahan atau pengurangan berat badan,
- Sangat jarang (terjadi pada kurang dari 1 dalam 10,000 pesakit): kecelaruan minda, gangguan kardiovaskular (degupan jantung tidak tetap, angina, serangan jantung), eosinophilic pneumonia (satu jenis pneumonia luar biasa), erythema multiforme (kulit ruam yang sering bermula

dengan tompok-tompok gatal merah pada muka, lengan atau kaki), gangguan darah, pankreas, perut atau hati, peripheral neuropathy (penyakit yang menyebabkan kehilangan sensasi, kesakitan, ketidakupayaan mengawal otot), hipertonia (peningkatan dalam ketegangan otot yang luar biasa), bengkak gusi, gula darah yang tinggi.

- Kesan sampingan berikut juga dilaporkan oleh pesakit-pesakit yang mengambil COVERAM: hypoglicaemia (paras gula darah yang sangat rendah), vaskulitis (keradangan pada saluran darah), gegaran dan ketegaran.

Jika sebarang kesan sampingan menjadi serius atau anda mengalami kesan sampingan yang tidak disenaraikan seperti yang diatas, sila beritahu doktor anda atau ahli farmasi anda.

Selepas menggunakan COVERAM

Bagaimana menyimpan COVERAM

Jauhi dari dicapai atau dilihat oleh kanak-kanak.

Jangan mengambil COVERAM selepas tarikh luput yang tercetak pada kotak dan botol.

Tarikh luput adalah merujuk hari terakhir untuk bulan tersebut.

Bekas ini perlu ditutup rapi untuk mengelakkan daripada penyerapan lembapan dari luar.

Cara pembuangan

Ubat ini tidak sepatutnya dibuang melalui system pembuangan air rumah atau bersama barang buangan lain. Sila bertanya ahli farmasi anda cara-cara membuang ubat yang tidak diperlukan. Langkah ini boleh membantu menjaga persekitaran kita.

Maklumat Produk

Apa yang terkandung di dalam COVERAM

Bahan-bahan aktif ialah arginine perindopril dan amlodipine.

COVERAM 5mg / 5mg: sebiji tablet mengandungi 5 mg arginine perindopril dan 5 mg amlodipine.


COVERAM 10mg / 5mg: sebiji tablet mengandungi 10 mg arginine perindopril dan 5 mg amlodipine.


COVERAM 5mg / 10mg: sebiji tablet mengandungi 5 mg arginine perindopril dan 10 mg amlodipine.


COVERAM 10mg / 10mg: sebiji tablet mengandungi 10 mg arginine perindopril dan 10 mg amlodipine.


Bahan-bahan lain dalam tablet: lactose monohydrate, magnesium stearate (E470B), cellulose microcrystalline (E460), colloidal anhydrous silica (E551).

Bagaimana rupa bentuk dan pembungkusan COVERAM

COVERAM 5mg / 5mg ialah tablet yang berwarna putih, berbentuk rod yang diukir dengan 5 / 5 pada satu muka dan  di muka yang lain.

COVERAM 10mg / 5mg ialah tablet yang berwarna putih, berbentuk tiga segi yang diukir dengan 10 / 5 pada satu muka dan  di muka yang lain.

COVERAM 5mg / 10mg ialah tablet yang berwarna putih, berbentuk empat segi yang diukir dengan 5 / 10 di satu muka dan  di yang lain.

COVERAM 10mg / 10mg ialah tablet yang berwarna putih, berbentuk bulat, yang diukir dengan 10 / 10 di satu muka dan  di yang lain.

Terdapat di dalam botol yang mengandungi 30 tablet.

No. pendaftaran PKBD:

COVERAM 5mg / 5mg:

MAL20091910A

COVERAM 5mg / 10mg:

MAL20091911A

COVERAM 10mg / 5mg:

MAL20091912A

COVERAM 10mg / 10mg:

MAL20091913A

Pengilang:

Servier (Ireland) Industries Ltd
Gorey Road
Arklow - Co. Wicklow – Ireland

Pemegang Pendaftaran:

Servier Malaysia Sdn Bhd
1301 Level 13, Uptown 2
No.2 Jalan SS21/37,
Damansara Uptown
46400 Petaling Jaya
Selangor