

AXCEL EVILINE TABLET

Magnesium Hydroxide/ Aluminium Hydroxide/ Simethicone (200mg/ 200mg/ 20mg)

What is in this leaflet

1. What Axcel Eviline Tablet is used for
2. How Axcel Eviline Tablet works
3. Before you use Axcel Eviline Tablet
4. How to use Axcel Eviline Tablet
5. While you are using it
6. Side effects
7. Storage and Disposal of Axcel Eviline Tablet
8. Product Description
9. Manufacturer and Product Registration Holder
10. Date of revision

What Axcel Eviline Tablet is used for

Axcel Eviline Tablet is an antacid / deflatulent for symptomatic treatment of:

- Stomach ulcer.
- Indigestion of functional or organic origin.
- Heartburn in hiatus hernia or pregnancy.
- Flatulence and abdominal distension.
- Gastritis oesophagitis (inflammation of the esophagus lining).
- Other conditions where hyperacidity or flatulence may be present

How Axcel Eviline Tablet work

Axcel Eviline Tablet is an antacid used to prevent and relief pain in the symptomatic management of gastric and duodenal ulcers and gastrointestinal reflux by neutralising hydrochloric acid in the gastric secretions.

Before you use Axcel Eviline Tablet

- When you must not use it

Do not take Axcel Eviline Tablet if:

- It is not prescribed by a qualified doctor or pharmacist for you.
- You are pregnant, trying to get pregnant or think you may be pregnant.
- You are breast-feeding. Ask your doctor or pharmacist for

advice before taking any medicine.

- Before you start to use it

Tell your doctor if:

- You are allergic to any medicines, food and dyes.
- You are taking any other medicines (including those not prescribed by your doctor).
- You are having hypophosphataemia (low level of phosphate in blood) and impaired kidney function.

- Taking other medicines

You should tell your doctor if you are taking the following medicine as it may interact with your Axcel Eviline Tablet:

- Tetracycline and vitamins.

It may still be all right for you to take Axcel Eviline Tablet and your doctor will be able to decide what is suitable for you.

How to use Axcel Eviline Tablet

- How much to use

1-2 tablets to be chewed or sucked 3 times daily or as required, preferably after meals.

- When to use it

Your doctor will decide when it is appropriate for you to use Axcel Eviline Tablet according to the type of condition you have.

- How long to use it

Continue taking Axcel Eviline Tablet for as long as doctor recommends.

- If you forget to use it

Consult your doctor or pharmacist on what you should do if you forget to use it.

Take the missed dose as soon as you remember. If it is almost time for your next dose, wait until then to take the medicine and skip the missed dose. Do not take a double dose to make up for the missed dose.

- If you use too much (overdose)

Contact your doctor immediately or go to the Emergency Department of your nearest hospital, if you think you or anyone else may have taken too much of this medicine. Do this even if there are no signs of discomfort or poisoning. You may need urgent medical attention.

While you are using it

- Things you must do

- Take your medicine exactly as your doctor has told you.
- Tell all the doctors, dentists and pharmacists treating you that you are taking Axcel Eviline Tablets.
- Tell your doctor immediately if you become pregnant while talking this medication.

- Things you must not do

- Do not stop taking the medicine unless advised by your doctor.
- Do not take any new medicines without consulting your doctor.
- Do not give Axcel Eviline Tablets to anyone else, even if they have the same symptoms or condition as you.

- Things to be careful of

- Avoid prolonged use with high dosage.
- Avoid administration during the first trimester of pregnancy.

Side effects

Like all medicines, Axcel Eviline Tablets can cause side effects although not everybody gets them. You are advised to consult doctor/pharmacist for further information. The following common side effects have been reported:

- In kidney failure, the administration of aluminium hydroxide can elevate the plasma aluminium concentration.
- Low level of phosphate in the blood and urine.
- Phosphate depletion syndrome.
- High level of calcium in the blood.
- Low level of magnesium in the blood.

AXCEL EVILINE TABLET

Magnesium Hydroxide/ Aluminium Hydroxide/ Simethicone (200mg/ 200mg/ 20mg)

- Nausea, vomiting and constipation.
- Intestinal obstruction and hard stool.
- Cathartic effect (accelerates defecation).
- Neurological, neuromuscular and cardiovascular impairment and even death in persons with kidney impairment.
- Alkalinization of the urine decreases the excretion of various weakly basic drugs eg. Quinidine.
- Magnesium hydroxide increases the absorption of warfarin.

- Inactive Ingredients:
PVP K90, Corn Starch,
Magnesium Stearate, Purified
water, Saccharin Sodium,
Peppermint Oil, Lactose, Icing
Sugar, Methylparaben and
Mannitol.

- MAL Number
MAL19911458X

**Manufacturer and Product
Registration Holder**

Kotra Pharma (M) Sdn Bhd
No. 1, 2 & 3, Jalan TTC 12,
Cheng Industrial Estate,
75250 Melaka, Malaysia.

Date of revision
21/02/2018

Serial number
NPRA(R4/1)221217/00372

You may report any side effects or adverse drug reactions directly to the National Centre for Adverse Drug Reaction Monitoring by calling Tel: 03-78835550, or visiting the website npra.moh.gov.my (Public → Reporting Medicinal Problems / Side Effects / AEFI / Vaccine Safety).

Storage and Disposal of Axcel Eviline Tablet

- Storage

Keep Axcel Eviline Tablet out of reach of children.

Store below 30°C. Keep container well closed and protect from light.

- Disposal

Do not use Axcel Eviline Tablets after the expiry date which is stated on the carton.

Medicines should not be disposed of via wastewater or household waste. Ask your pharmacist how to dispose of medicines no longer required. These measures will help to protect the environment.

Product Description

- What it looks like

Axcel Eviline Tablets is a white, flat, round tablets, 12.9mm in diameter, and marked KOTRA logo on one side.

- Ingredients

- Active Ingredients:

Magnesium Hydroxide,
Aluminium Hydroxide and
Simethicone.