

CLAMENTIN POWDER FOR SYRUP

Amoxicillin/Clavulanic acid (31.25mg/ml)

What is in this leaflet

1. What CLAMENTIN POWDER SYRUP is used for
2. How CLAMENTIN POWDER SYRUP works.
3. Before you use CLAMENTIN POWDER SYRUP.
4. How to use CLAMENTIN POWDER SYRUP
5. While you are using CLAMENTIN POWDER SYRUP.
6. Side effects.
7. Storage and disposal of CLAMENTIN POWDER SYRUP
8. Product description
9. Manufacturer and Product Registration Holder
10. Date of revision

What CLAMENTIN POWDER SYRUP is used for

It is used for short-term treatment in upper respiratory tract infection, lower respiratory tract infections, skin and soft tissue infections and genital urinary tract infections.

Your doctor may have prescribed CLAMENTIN POWDER SYRUP for another reason. Ask your doctor if you have any questions about why CLAMENTIN POWDER SYRUP has been prescribed for you.

How CLAMENTIN POWDER SYRUP works

CLAMENTIN POWDER SYRUP consists of active substances Amoxicillin/Clavulanic Acid. These medicine work by killing bacteria, which cause infections.

Before you use CLAMENTIN POWDER SYRUP

-When you must not use it

Do not use CLAMENTIN POWDER SYRUP if:

-you have had an allergic reaction to penicillin or similar types of antibiotics (such as cephalosporins) or any of the ingredients contained in CLAMENTIN POWDER SYRUP. The ingredients are listed at the end of this leaflet. Some of the symptoms of an allergic reactions may include skin rash, itching, difficulty in breathing and swelling of the face or tongue.

-you have previously experienced liver problems after using CLAMENTIN POWDER SYRUP or any other penicillin antibiotic.

-Before you start to use it

Tell your doctor if:

-you have ever had an allergic reaction (such as a rash) to antibiotics such as penicillin or other substances in the past.

-you are allergic to foods, dyes, preservatives or any other medicines.

-you have experienced liver problems after using CLAMENTIN POWDER SYRUP or any other medicines.

-you have glandular fever (mononucleosis).

-you have kidney or liver problem.

-you have phenylketonurics (metabolism disorder)

-Taking other medicines

Tell your doctor if you are taking any other medicines, including any that you buy without a prescription from a pharmacy, supermarket or health food shop.

Some medicines when taken with CLAMENTIN POWDER SYRUP may interfere with each other. These include:

-oral contraceptives.

-probenecid, medicine for gout.

-allopurinol, medicine for gout.

Your doctor or pharmacist may have more information on medicines to be careful with or to avoid while taking medication.

How to use CLAMENTIN POWDER SYRUP

-How much to use

Prepare syrup at time of dispensing. Tap bottle until all powder flows freely. Add approximately 2/3 of the total amount of water for reconstitution and shake vigorously to suspend powder. Add remaining water to 100mL and again shake vigorously.

7-12 years: 10mL, 3 times daily

2-7 years: 5mL 3 times daily

Children 9 months -2 years:

25mg/kg/day in 3 divided doses

Severe infections: Increase frequency to 6 hourly intervals.

Follow all directions given to you by your doctor and pharmacist carefully. They may differ from the information contained in this leaflet. If you do not understand the instructions on the label, ask your doctor or pharmacist for help.

-When to use it

Consult your doctor on the time of consumption.

-How long to use it

Consult your doctor on your duration of consumption of this medication.

Do not stop taking medication, even if you feel better after a few days, unless advised by your doctor. Your infection may not clear completely if you stop taking your medicine too soon.

-If you forget to use it

If you miss a dose of CLAMENTIN POWDER SYRUP and you are taking it regularly, take it as soon as possible. If several hours have passed or if it is approaching time for the next dose, do not double the dose to make up for the forgotten dose, unless advised by your health care provider. Do not take 2 doses at once.

-If you use too much (overdose)

Quickly seek for medical attention if you think that you or anyone else has taken too much CLAMENTIN POWDER SYRUP

While you are using CLAMENTIN POWDER SYRUP

-Things you must do

Follow your doctor's instruction when you are taking CLAMENTIN POWDER SYRUP.

Tell all the doctors, dentists and pharmacists treating you that you are taking CLAMENTIN POWDER SYRUP.

If you get severe diarrhoea, tell your doctor or pharmacist immediately.

-Things you must not do

Do not stop taking this medicine or change the dosing regimen without instruction from your doctor. Do not give CLAMENTIN POWDER SYRUP to anyone else, even if they have the same condition as you.

-Things to be careful of

If your child develops itching, swelling or a skin rash while using CLAMENTIN POWDER SYRUP, do not give any more CLAMENTIN POWDER SYRUP and tell your doctor immediately.

Side effects

CLAMENTIN POWDER SYRUP may cause gastrointestinal side effect such as diarrhea, nausea and vomiting. It may give liver effects and cause hypersensitivity reactions.

It may cause blood disorder and central nervous system effects such as dizziness and headache.

Like all medicines, CLAMENTIN POWDER SYRUP can cause side effects, although not everybody gets them.

Visit your doctor or pharmacist immediately if you are experience any other side effects after taking this medicine.

You may report any side effects or adverse drug reactions directly to the National Centre for Adverse Drug Reaction Monitoring by calling Tel: 03-78835550, or visiting the website portal.bpfk.gov.my
(Consumers → Reporting)

Storage and Disposal of CLAMENTIN POWDER SYRUP

-Storage

Store below 30°C, protect from light & moisture.

Reconstituted syrup must be stored under refrigeration and discarded after 7 days.

-Disposal

CLAMENTIN POWDER SYRUP should be disposed when your doctor or pharmacist tells you to stop taking this tablet or it has passed the expiry date. It should not be disposed via wastewater or household waste. Ask your pharmacist how to dispose medicines that are no longer required. These measures will help to protect the environment.

Product description

What it looks like

A white to off-white powder.

Appearance after reconstitution:

An off-white suspension

-Ingredients

-Active ingredient:

Amoxicillin Trihydrate

Clavunate Salt

-Inactive ingredients:

Silicon Dioxide

Aspartame

Mannitol

Orange Powder

Xanthan Gum

-MAL number

MAL09032000AZ

Manufacturer

Yung Shin Pharmaceutical Ind. Co., Ltd.
(Taichung Youth Factory III)
No. 26 Kung Liu Rd. Youth Industrial District, Taichia, Taichung, Taiwan, R.O.C

Product Registration Holder

Y.S.P Industries (M) Sdn. Bhd.
Lot 3,5 & 7, Jalan P/7, Section 13, Kawasan Perindustrian Bandar Baru Bangi, 43000 Kajang, Selangor Darul Ehsan, Malaysia.

Date of revision

21/4/2016

Serial Number

BPFK(R4/1)200416/00117