

# **ANASMA EXPECTORANT®**

**SALBUTAMOL SULPHATE 1.2 mg/5 mL & GUAIPHENESIN 50 mg/5 mL**

<b>What is in this leaflet:</b>	<b>How to take Anasma Expectorant</b>	<b>Side Effects</b>
<ol style="list-style-type: none"> <li>1. What Anasma Expectorant is used for</li> <li>2. How Anasma Expectorant works</li> <li>3. Before you use Anasma Expectorant</li> <li>4. How to take Anasma Expectorant</li> <li>5. While you are using Anasma Expectorant</li> <li>6. Side Effects</li> <li>7. How to store and dispose</li> <li>8. Product description</li> </ol>	<p><b>How much to take</b> Always take Anasma Expectorant exactly as your doctor's advice. If you are unsure, do check with your doctor or pharmacist.</p> <p><b>Doses:</b> Adults : 10-20 mL two or three times a day. Children (6-12 years) : 10 mL two or three times a day. Children (2-6 years) : 5-10 mL two or three times a day.</p>	<p>Frequent incidents are, urge to vomit, nervousness, trembling, and unusually fast or pounding heartbeat. Urination, dizziness or lightheadedness, dryness or irritation of the mouth and throat, headache, heartburn, increase or decrease in blood pressure, restlessness, trouble in sleeping, unusual taste in mouth and vomiting may occur less frequently.</p>
<p><b>What Anasma Expectorant is used for</b> Anasma Expectorant is indicated for the relief of productive and wheezing cough associated with airway infections.</p>	<p><b>When to take it</b> Take 1 hour before or 2 hours after meal.</p>	<p>Usage during pregnancy and lactation should be considered based on risk benefit.</p>
<p><b>How Anasma Expectorant works</b> It contains salbutamol and guaiphenesin. Salbutamol works by relaxing the muscles of the airways of the lungs. Guaiphenesin promotes or facilitates the removal of excretions from the airway. The use of these two compounds maintain adequate ventilation by making lungs' secretion less viscous, thus, facilitating expulsion and relieving airway constriction.</p>	<p><b>How long to take it</b> Do not stop taking this medicine even when you feel better, unless being advised by your doctor.</p>	<p><b>How to store and dispose</b> <b>Storage</b> Store below 25°C, away from direct light. Keep out of the reach of children.</p>
<p><b>Before you use Anasma Expectorant</b> Anasma Expectorant should not be used in patients with known allergy to salbutamol.</p>	<p><b>If you forget to take it</b> If you miss a dose of Anasma Expectorant, take it as soon as possible. However, if its due for the next dose, skip the missed dose and continue with your regular dosing schedule. Do not double up the doses.</p>	<p><b>Product Description</b> <b>What it looks like</b> Anasma Expectorant is a clear solution without any foreign particles with raspberry flavour.</p>
<p>This medicine should be used carefully in patients with medical problems such as:</p> <ol style="list-style-type: none"> <li>1. Heart disease</li> <li>2. Diabetes mellitus</li> <li>3. Enlarged prostate</li> <li>4. Excessive thyroid hormone</li> </ol>	<p><b>If you take too much (overdose)</b> If you have taken more than the prescribed dose, contact your doctor immediately. Signs of an overdose are chest pain, severe dizziness, prolonged headache, severe increase in blood pressure, continuous nausea or vomiting, unusual nervousness or restlessness and muscle twitching.</p>	<p><b>Ingredients</b> The active substances of Anasma Expectorant are salbutamol sulphate and guaiphenesin. The other ingredients of Anasma Expectorant are sugar, sodium saccharine, citric acid monohydrate, propylene glycol, sodium chloride, sodium citrate, methyl paraben, propyl paraben and raspberry flavour.</p>
<p>Please inform your doctor if you are taking other medicines, especially:</p> <ol style="list-style-type: none"> <li>1. Guanethidine, reserpine or methyldopa (for high blood pressure)</li> <li>2. Monoamine oxidase inhibitors e.g. tranylcypromine (for depression)</li> <li>3. Tricyclic antidepressants eg. amitriptyline (for depression)</li> <li>4. Corticosteroid</li> <li>5. Theophylline (for breathing problem)</li> </ol>	<p><b>While you are using Anasma Expectorant</b> Check with your doctor at once:</p> <ol style="list-style-type: none"> <li>1. If you still have trouble in breathing 1 hour after taking this medicine.</li> <li>2. If your symptoms return within 4 hours after using this medicine.</li> <li>3. If your condition gets worse.</li> </ol>	<p>Packed in 100 and 200 mL bottle. Packed in 3.6 L bottle (for tender only).</p> <p><b>Registration Numbers:</b> Anasma Expectorant: MAL20014345A</p>
		<p><b>Manufactured by:</b> Idaman Pharma Manufacturing Sdn Bhd (661901-P) Lot 24, Bakar Arang Industrial Estate, Sungai Petani, Kedah, Malaysia.</p>

Date of revision:

Marketing Authorisation Holder

June 10<sup>th</sup>, 2012

IDAMAN PHARMA MANUFACTURING SDN BHD

# **ANASMA EXPECTORANT®**

**SALBUTAMOL SULPHATE 1.2 mg/5 mL & GUAIPHENESIN 50 mg/5 mL**

<p><b>Apakah yang ada pada risalah ini:</b></p> <ol style="list-style-type: none"> <li>1. Apa kegunaan Expectorant Anasma</li> <li>2. Bagaimana Expectorant Anasma</li> <li>3. Sebelum mengambil Expectorant Anasma</li> <li>4. Cara pengambilan Expectorant Anasma</li> <li>5. Semasa menggunakan Expectorant Anasma</li> <li>6. Kesan-kesan sampingan</li> <li>7. Cara penyimpanan dan pelupusan</li> <li>8. Diskripsi produk</li> </ol> <p><b>Apa kegunaan Expectorant Anasma</b> Anasma Expectorant digunakan untuk batuk berkahak dan nafas berdehit disebabkan oleh jangkitan salur pernafasan.</p> <p><b>Bagaimana Expectorant Anasma berfungsi</b> Ubat ini mengandungi salbutamol dan guaiphenesin. Salbutamol berfungsi merehatkan otot paru-paru dan memudahkan pernafasan. Guaiphenesin membantu mengeluarkan kahak dari salur pernafasan. Kehadiran kedua-dua bahan ini akan mengekalkan ventilasi paru-paru dengan mencairkan kahak, memudahkan penyingkirannya dan melegakan salur pernafasan.</p> <p><b>Sebelum mengambil Expectorant Anasma</b> Ubat ini tidak seharusnya digunakan pada pesakit yang mempunyai alahan terhadap salbutamol.</p> <p>Ubat ini harus digunakan berhati-hati pada pesakit yang mempunyai masalah perubatan seperti:</p> <ol style="list-style-type: none"> <li>1. Penyakit jantung</li> <li>2. Kencing manis</li> <li>3. Pembesaran kelenjar prostat</li> <li>4. Hormon tiroid berlebihan</li> </ol> <p>Sila maklumkan kepada doktor anda jika anda mengambil ubat-ubatan lain, terutamanya:</p> <ol style="list-style-type: none"> <li>1. Guanethidine, reserpine atau methyldopa (untuk tekanan darah tinggi)</li> <li>2. 'Monoamine oxidase inhibitors' contohnya tranylcypromine (untuk kemurungan)</li> <li>3. 'Tricyclic antidepressants' contohnya amitriptyline (untuk kemurungan)</li> <li>4. Corticosteroid</li> <li>5. Theophylline (untuk masalah pernafasan)</li> </ol>	<p><b>Cara pengambilan Expectorant Anasma</b></p> <p><b>Berapa banyak untuk diambil</b> Sentiasa mengambil ubat ini tepat mengikut arahan doktor anda. Jika anda was-was, sila rujuk kepada doktor atau ahli farmasi anda</p> <p><b>Dos:</b> Dewasa : 10-20 mL dua atau tiga kali sehari Kanak-kanak (6 tahun ke atas): 10 mL dua atau tiga kali sehari Kanak-kanak (2-6 tahun): 5-10 mL dua atau tiga kali sehari</p> <p><b>Bila ubat diambil</b> Minum ubat ini 1 jam sebelum atau 2 jam selepas makan.</p> <p><b>Berapa lama ubat diambil</b> Jangan berhenti minum ubat ini walaupun anda berasa lebih sihat melainkan atas arahan doktor anda.</p> <p><b>Jika anda lupa mengambil ubat</b> Jika anda terlewat mengambil dos, ambil ubat secepat mungkin. Jika hampir masa untuk dos berikutnya, abaikan dos tersebut dan teruskan dos rutin anda. Jangan mengambil 2 dos sekaligus.</p> <p><b>Jika anda terambil ubat terlalu banyak</b> Jika anda terambil dos lebih daripada yang ditetapkan, hubungi doktor serta merta. Tanda-tanda akibat dos berlebihan adalah sakit dada, pening yang teruk, sakit kepala yang berterusan, peningkatan tekanan darah yang teruk, loya atau muntah-muntah, rasa cemas yang luar biasa, keresahan dan sentakan otot.</p> <p><b>Semasa menggunakan Expectorant Anasma</b></p> <p>Periksa dengan doktor anda serta merta:</p> <ol style="list-style-type: none"> <li>1. Jika anda masih mempunyai masalah pernafasan 1 jam selepas mengambil ubat ini.</li> <li>2. Jika gejala anda kembali dalam masa 4 jam selepas mengambil ubat ini.</li> <li>3. Jika keadaan anda menjadi lebih teruk.</li> </ol>	<p><b>Kesan-kesan sampingan</b> Insiden yang kerap terjadi adalah loya, rasa cemas, menggeletar, dan denyutan jantung yang cepat atau berdebar-debar. Kencing, pening atau seperti rasa ringan kepala, kekeringan atau kerengsaan pada mulut dan tekak, sakit kepala, pedih ulu hati, peningkatan atau penurunan tekanan darah, keresahan, masalah untuk tidur, rasa luar biasa di dalam mulut dan muntah mungkin jarang berlaku.</p> <p>Penggunaan ubat ini semasa mengandung dan penyusuan perlu berdasarkan risiko dan manfaatnya.</p> <p><b>Cara penyimpanan dan pelupusan</b></p> <p><b>Penyimpanan</b> Simpan di bawah suhu 25°C. Jauhi daripada cahaya dan kanak-kanak.</p> <p><b>Pelupusan</b> Jangan buang ubat ke dalam longkang atau tong sampah. Sila bertanya kepada ahli farmasi anda berkenaan cara pelupusan ubat.</p> <p><b>Diskripsi produk</b></p> <p><b>Rupa produk</b> Expectorant Anasma ialah cecair tidak berwarna tanpa sebarang bahan asing dan berperisa rasberi.</p> <p><b>Bahan kandungan</b> Bahan aktif ubat ini ialah 'salbutamol sulphate' dan 'guaiphenesin'. Kandungan lain Expectorant Anasma ialah gula, 'sodium saccharine', 'citric acid monohydrate', 'propylene glycol', 'sodium chloride', 'sodium citrate', 'methyl paraben', 'propyl paraben' dan perisa rasberi.</p> <p>Saiz pek ialah 100 dan 200 mL botol. Pek 3.6 L botol (untuk tender sahaja).</p> <p><b>Nombor Pendaftaran:</b> Expectorant Anasma: MAL20014345A</p> <p><b>Dikilangkan oleh:</b> Idaman Pharma Manufacturing Sdn Bhd (661901-P) Lot 24, Bakar Arang Industrial Estate, Sungai Petani, Kedah, Malaysia.</p>
---	--	---

Tarikh semakan:

Pemegang Otorisasi Pasaran

Jun 10, 2012

IDAMAN PHARMA MANUFACTURING SDN BHD