

REXOMTM BROMHEXINE ELIXIR

Bromhexine Hydrochloride (4mg/5ml)

What is in this leaflet

1. What REXOMTM Bromhexine is used for
2. How REXOMTM Bromhexine works
3. Before you use REXOMTM Bromhexine
4. How to use REXOMTM Bromhexine
5. While you are using REXOMTM Bromhexine
6. Side effects
7. Storage and disposal of REXOMTM Bromhexine
8. Product description
9. Manufacturer and Product Registration Holder
10. Date of revision

What REXOMTM Bromhexine is used for

It is used to loosen the mucus, making it easier to cough up the mucus for airway diseases, accompanied by excessive mucus and impaired mucus clearance.

How REXOMTM Bromhexine works

REXOMTM Bromhexine helps to clear the chest by thinning the mucus in your air passages.

Before you use REXOMTM Bromhexine

-When you must not use it

It should not be used if you had known to be hypersensitive to Bromhexine or other components of the formulation.

-Before you start to use it

- There will be an expected increase in the flow of secretions.
- Should be used with care in people with a history of ulcer in digestive tract and asthma.

Pregnancy and lactation

Talk to your doctor before taking this medicine if you are pregnant, might become pregnant or are breastfeeding. Bromhexine should only be used if the maternal condition justifies the potential risk to the fetus or baby.

-Taking other medicines

Inform your doctor or pharmacist before taking this medicines, including any that you get without a prescription from your

pharmacy, supermarket or health food shop.

Some medicines and REXOMTM Bromhexine may interfere with each other. They may be affected by REXOMTM Bromhexine or may affect how well it works.

- It may increase the concentration of concurrently used antibiotics.

How to use REXOMTM Bromhexine

-How much to use

Always use REXOMTM Bromhexine exactly as your doctor has prescribed. You should check with your doctor or pharmacist if in doubt.

The usual dose:

Under 2 years: 1.25 ml

2 to 6 years: 2.5 ml

6 to 12 years: 5 ml

-When to use it

To be taken 3 times a day. Use as directed by your doctor or pharmacist.

-How long to use it

Continue taking REXOMTM Bromhexine as your doctor recommended. If symptoms persist, consult your doctor.

-If you forget to use it

Consult your doctor or pharmacist on what you should do if you forget to use it.

Take the missed dose as soon as you remember. If it is almost time for your next dose, wait until then to take the medicine and skip the missed dose. Do not take a double dose to make up for the missed dose.

-If you use too much (overdose)

Contact your doctor immediately or go to the Emergency Department of your nearest hospital, if you think you or anyone else may have taken too much of this medicine. Do this even if there are no signs of discomfort or poisoning. You may need urgent medical attention.

While you are using REXOMTM Bromhexine

-Things you must do

- Tell all the doctors, dentists and pharmacists treating you that you are taking REXOMTM Bromhexine.

- Tell your doctor immediately if you become pregnant while taking this medication.

- Things you must not do

- Do not take more than the recommended dose unless your pharmacist or doctor tells you.
- Do not take any new medicines without consulting your doctor or pharmacist.
- Do not give REXOMTM Bromhexine to anyone else, even if they have the same symptoms or condition as you.

-Things to be careful of

Be careful driving or operating machinery until you know how REXOMTM Bromhexine affects you.

Side effects

Like all medicines, REXOMTM Bromhexine can cause side effects, although not everybody gets them.

If side effects persist, do not take the medicine and refer to the doctor.

Tell your doctor if you notice any of the following side effects or notice any other effects not listed:

- Gastrointestinal side effects
- Transient rise in blood test aminotransferase
- Headache
- Dizziness
- Sweating
- Skin rashes

You may report any side effects or adverse drug reactions directly to the National Centre for Adverse Drug Reaction Monitoring by calling Tel: 03-78835550, or visiting the website npra.moh.gov.my (Public Reporting Medicinal Problems / Side Effects / AEFI / Vaccine Safety).

Storage and disposal of REXOMTM Bromhexine

-Storage

Keep container tightly closed. Store below 30°C. Protect from heat and light. Keep medicine out of reach of children.

REXOM™ BROMHEXINE ELIXIR

Bromhexine Hydrochloride (4mg/5ml)

-Disposal

Medicines should not be disposed of via wastewater or household waste. Ask your pharmacist how to dispose of medicines no longer required. These measures will help to protect the environment.

Product description

-What it looks like

Colourless liquid with strawberry flavor.

-Ingredients

-Active ingredient

Each 5ml contains

Bromhexine Hydrochloride (4mg)

-Inactive ingredients

Strawberry flavour, Menthol Crystals, Sodium Benzoate, Sodium Citrate, Saccharin Sodium, Citric Acid Anhydrous, Propylene Glycol, Hydroxypropyl Methylcellulose and Purified Water.

-MAL number

MAL19962479XZ

**Manufacturer and Product
Registration Holder**

KCK Pharmaceutical Industries Sdn.
Bhd.

Plot 61, Bayan Lepas Industrial Park
Phase 4, 11900 Penang,
Malaysia.

Date of revision

17/10/2017

Serial Number

NPRA(R4/1)161017/00273