

# AXCEL DEXXON LOZENGES

Dequalinium Chloride (0.25mg)

## What is in this leaflet

1. What Axcel Dexion Lozenges is used for.
2. How Axcel Dexion Lozenges works
3. Before you use Axcel Dexion Lozenges.
4. How to use Axcel Dexion Lozenges.
5. While you are using it.
6. Side effects.
7. Storage and disposal of Axcel Dexion Lozenges.
8. Product description.
9. Manufacturer and Product Registration Holder.
10. Date of revision.

## What Axcel Dexion Lozenges is used for

Axcel Dexion Lozenges is used in the treatment of infection such as:

- Minor infection of the mouth and throat.
- Anginous infection (bacterial infection of mouth).
- Catarrhal infection (inflammation of mucous membrane especially of the nose and throat).
- Inflammation of the mucous membrane and gums.

## What Axcel Dexion Lozenges is used for

Dequalinium Chloride is an antiseptic medicine. It works against a wide range of bacteria, yeasts and fungi.

## Before you use Axcel Dexion Lozenges

### -When you must not use it

Do not take Axcel Dexion Lozenges if:

- You are allergy or hypersensitive to dequalinium.
- Your child is under 3 years of age.
- You are being treated with sulphonamides (antibiotic).
- It is not prescribed by a qualified doctor or pharmacist for you.
- You are pregnant, trying to get pregnant or think you may be pregnant.

- You are breast-feeding. Ask your doctor or pharmacist for advice before taking any medicine.

### -Before you start to use it

Tell your doctor if:

- You are allergic to medicines, food and dyes.
- You are taking any other medicines (including those not prescribed by your doctor).

### -Taking other medicines

Tell your doctor if you are taking any other medicines, including any that you buy without a prescription from a pharmacy, supermarket or health food shop.

It may still be all right for you to use Axcel Dexion Lozenges and your doctor will be able to decide what is suitable for you.

## How to use Axcel Dexion Lozenges

### -How much to use

Adults and children (above 3 years): Allow 1 lozenge to be dissolved slowly in the mouth every 2 hours. After the inflammatory symptoms have subsided, suck 1 lozenge every 4 hours.

### -When to use it

Your doctor will decide when it is appropriate for you to use Axcel Dexion Lozenges according to the type of condition you have.

### -How long to use it

Continue taking Axcel Dexion Lozenges for as long as doctor recommends.

### -If you forget to use it

Consult your doctor or pharmacist on what you should do if you forget to use it.

Take the missed dose as soon as you remember. If it is almost time for your next dose, wait until then to take the medicine and skip the missed dose. Do not take a double dose to make up for the missed dose.

### -If you use too much (overdose)

Contact your doctor immediately or go to the Emergency Department of your nearest hospital, if you think you or anyone else may have taken too much of this medicine. Do this even if there are no signs of discomfort or poisoning. You may need urgent medical attention.

## While you are using it

### -Things you must do

- Take your medicine exactly as your doctor has told you.
- Tell all the doctors, dentists and pharmacists treating you that you are taking Axcel Dexion Lozenges.
- Tell your doctor immediately if you become pregnant while talking this medication.

### -Things you must not do

- Do not stop taking the medicine unless advised by your doctor.
- Do not take any new medicines without consulting your doctor or pharmacist.
- Do not give Axcel Dexion Lozenges to anyone else, even if they have the same symptoms or condition as you.

### -Things to be careful of

- Severe sore throat accompanied by high fever or headache, nausea and vomiting, any sore throat persisting more than 2 days may be serious, and need to be under medical attention.

## Side effects

Side effects and undesirable accompanied symptoms of Axcel Dexion Lozenges have not been observed. You are advised to consult doctor/ pharmacist for further information.

You may report any side effects or adverse drug reactions directly to the National Centre for Adverse Drug Reaction Monitoring by calling Tel: 03-78835550, or visiting the website [npra.moh.gov.my](http://npra.moh.gov.my) (Public → Reporting Medicinal Problems / Side Effects / AEFI / Vaccine Safety).

# AXCEL DEXXON LOZENGES

Dequalinium Chloride (0.25mg)

---

## **Storage and disposal of Axcel Dexton Lozenges**

### -Storage

Keep Axcel Dexton Lozenges out of reach of children.

Store below 30°C. Keep container well closed and protect from light.

### -Disposal

Do not use Axcel Dexton Lozenges after the expiry date which is stated on the carton.

Medicines should not be disposed of via wastewater or household waste. Ask your pharmacist how to dispose of medicines no longer required. These measures will help to protect the environment.

## **Product description**

### -What it looks like

Axcel Dexton Lozenges is orange, flat, round lozenges orange flavoured, 12.9mm in diameter and marked Kotra logo on one side.

### -Ingredients

- Active Ingredient:  
Dequalinium Chloride

- Inactive Ingredients:  
Icing Sugar, Mannitol, Sodium Saccharin, Isopropyl Alcohol, P.V.P. K90, Sunset Yellow, Orange powder, Magnesium Stearate, Talcum Powder and Purified Water.

### -MAL Number

MAL19911425X

## **Manufacturer and product registration holder**

Kotra Pharma (M) Sdn Bhd  
No. 1, 2 & 3, Jalan TTC 12,  
Cheng Industrial Estate,  
75250 Melaka, Malaysia.

## **Date of revision**

29/03/2018

## **Serial number**

NPRA(R4/1)200318/00074