

REDAMOL TABLET

Paracetamol (500mg)

What is in this leaflet

1. What Redamol Tablet is used for
2. How Redamol Tablet works
3. Before you use Redamol Tablet
4. How to take Redamol Tablet
5. While you are using Redamol Tablet
6. Side effects
7. Storage and disposal of Redamol Tablet
8. Product description
9. Manufacturer and Product registration holder
10. Date of revision

What Redamol Tablet is used for

Redamol Tablet is used for the relief of mild to moderate pain and to reduce fever.

How Redamol Tablet works

Paracetamol helps to relieve pain by blocking the pain messengers called prostaglandins in the central nervous system.

Before you use Redamol Tablet

- When you must not use it:

Do not take this medicine if:

- You are allergic to paracetamol or any of the ingredients in this medicine.
- You have alcohol addiction.
- You have liver disease.
- You have viral infection.
- You have kidney function impairment.
- You have heart and blood vessels disease.

Ask your doctor or pharmacist for advice before taking any medicine.

- Before you start to use it:

Please tell your doctor or pharmacist if you are suffering or have ever suffered from any of the following conditions or illnesses:

- You are pregnant.
- You are a nursing mother.

- Taking other medicines:

Please tell your doctor or pharmacist if you are taking or have recently taken any other medicines, including medicines obtained without a prescription. Your doctor may need to change the dose of the other medications or take other precautions. In some cases you may have to stop taking one of the medicines. This applies especially to the medicines

listed below taken at the same time with Redamol Tablet:

- Paracetamol (Do not take other paracetamol-containing medicine at the same time).
- Cholestyramine (used to lower the cholesterol level in blood).
- Metoclopramide or Domperidone (used to treat nausea and vomiting).
- Warfarin (anticoagulant used to prevent heart attacks, strokes and blood clots).
- Chloramphenicol (antibiotic used for the treatment of bacterial infections).
- Busulfan (cancer medicine).

How to take Redamol Tablet

- How much to use:

Your doctor or pharmacist can advise you how many doses are needed each day and for how long you may need to take it. This depends on your condition and whether or not you are taking any other medicines.

The recommended dosages are as follows:

Adult: 1 to 2 tablets.

Children: 7 to 12 years – ½ to 1 tablet.

Not recommended for children under 6 years.

- When to use it:

To be taken 3 or 4 times a day.

Your doctor will decide when it is appropriate for you to use Redamol Tablet according to the type of condition you have.

- How long to use it:

Continue taking Redamol Tablet for as long as your doctor recommends.

- If you forget to use it:

Consult your doctor or pharmacist on what you should do if you forget to use it. Take the missed dose as soon as you remember. If it is almost time for your next dose, wait until then to take the medicine and skip the missed dose. Do not take a double dose to make up for the missed dose.

- If you use too much (overdose):

Contact your doctor immediately or go to the Emergency Department of your nearest hospital, if you think you or anyone else may have taken too much

of this medicine. Do this even if there are no signs of discomfort or poisoning. You may need urgent medical attention.

Symptoms of overdose include diarrhea, vomiting or nausea, loss of appetite, stomach cramps or pain, unusual increase in sweating.

While you are using Redamol Tablet

- Things you must do

- Take your medicine exactly as your doctor has told you.
- Tell all the doctors, dentists and pharmacists treating you that you are taking Redamol Tablet.
- Tell your doctor immediately if you become pregnant while taking this medication.

- Things you must not do

- Do not stop taking the medicine unless advised by your doctor.
- Do not take any new medicines without consulting your doctor or pharmacist.
- Do not give Redamol Tablet to anyone else, even if they have the same symptoms or condition as you.

- Things to be careful of

- Special care is recommended during pregnancy. Take the medicine during pregnancy if the benefits against possible effects on the child had been assessed by the doctor.
- May cause kidney impairment. Should be given with care to people with impaired kidney or liver function.

Side effects

Like all medicines, Redamol Tablet can cause side effects, although not everybody gets them. Visit your doctor or pharmacist immediately if you experience any of the side effects after taking this medicine:

- Bloody or cloudy urine.
- Difficult or painful urination.
- Frequent urge to urinate.
- Sudden decrease in amount of urine.
- Unusual bleeding or bruising.
- Unusual tiredness or weakness.
- Yellowing of eyes or skin.
- Skin hypersensitivity reactions including skin rashes, swelling under

REDAMOL TABLET

Paracetamol (500mg)

the skin, Steven Johnson Syndrome/
Toxic Epidermal Necrolysis.

- Severe skin reactions. Symptoms may include skin reddening, blisters or rash. These could be signs of a serious condition. If these reactions occur, stop use and seek medical assistance right away.

You may report any side effects or adverse drug reactions directly to the National Centre for Adverse Drug Reaction Monitoring by calling Tel:03-78835550 or visiting the website npra.moh.gov.my
(Public → Reporting Side Effects to Medicines (ConSERF) or Vaccines (AEFI).

Storage and disposal of Redamol Tablet

-Storage:

Store in a dry place below 30°C. Protect from light.

Keep out of reach of children.

-Disposal:

Medicines should not be disposed of via wastewater or household waste. Ask your pharmacist how to dispose the medicines you no longer required. These measures will help to protect the environment.

Product description

-What it looks like:

White, round tablet, 12.7mm in diameter, bevel-edge with breaking line one side and plain on the other side.

-Ingredients:

-Active Ingredient:

Paracetamol.

-Inactive Ingredients (in premix):

Pregelatinized starch, povidone and stearic acid.

-MAL No.:

MAL07091034XZ

Manufacturer and Product Registration Holder

Royce Pharma Manufacturing Sdn.Bhd.
PT 1663, Nilai Industrial Estate,
71800 Nilai, Negeri Sembilan,
Malaysia.

Date of revision

03/05/2018
