

U-MOLIN TABLETS

Paracetamol (500mg)

What is in this leaflet:

1. What U- Molin Tablets is used for
2. How U-Molin Tablets works
3. Before you use U- Molin Tablets
4. How to use U- Molin Tablets
5. While you are using it
6. Side Effects
7. Storage and Disposal of U- Molin Tablets
8. Product description
9. Manufacturer and Product Registration Holder
10. Date of Revision

What U- Molin Tablets is used for

It is indicated for the relief of mild to moderate pain and fever such as headache, toothache, colds, and influenza.

How U- Molin Tablets works

Paracetamol is the active ingredient of U- Molin Tablets. Paracetamol belongs to a group of medicines known as analgesics, which are used to relieve pain and to reduce fever.

Before you use U- Molin Tablets

- When you must not use it

- If you are allergic (hypersensitive) to paracetamol or any of the other ingredients of U- Molin Tablets .

- Before you start to use it

You must tell your doctor if:

- you have allergies to any medicine which you have taken previously to treat your current condition.
- you have kidney problems
- you have liver problems
- you have alcohol abuse.

Pregnancy and breastfeeding

You should inform your doctor if you are pregnant or wish to become pregnant, or wish to breast-feed, before taking this medicine.

There are usually no ill effects to paracetamol being used in the recommended dosage during pregnancy, however you should follow the advice of your doctor regarding its use.

- Taking other medicines

Tell your doctor or pharmacist if you are taking, have recently taken or might take any medicines that affect the liver.

You must not take any other medicines that contain paracetamol while you are taking this medicine.

Always inform your doctor and pharmacist if you are taking any other medicines, including herbal tonics, supplements and medicines that you buy without prescription.

How to use U- Molin Tablets

Always take this medicine exactly as your doctor has told you. You should check with your doctor or pharmacist if you are not sure. The tablet should be swallowed with a glass of water.

- How much to use

Adults and children aged 12 years and over:

1-2 tablets every 4 to 6 hours as required, to a maximum of 8 tablets in 24 hours.

Children (6-11 years):

½ - 1 tablets every 4-6 hours when necessary, with maximum of 4 tablets in 24 hours.

- When to use it

Your doctor will tell you the time when to use it. Always take this medicine exactly as your doctor has told you.

- How long to use it

Take this medicine for as long as your doctor tells you do.

- If you forget to use it

- If you forget a dose, take it as soon as you remember it.
- However, if it is nearly time for the next dose, skip the missed dose.
- Do not take a double dose to make up for a forgotten dose.

- If you use too much (Overdose)

Contact your doctor immediately or go to the Emergency Department of your nearest hospital, if you think you or anyone else may have taken too much of

this medicine. Do this even if there are no signs of discomfort or poisoning. You may need urgent medical attention.

While you are using it

- Things you must do

Contact your doctor if your symptoms worsen or they do not improve.

Be careful driving or operating machinery until you know how the medicine affects you.

- Things you must not do

Do not give U- Molin Tablets to anyone else, even if their symptoms seem similar or they have same condition as you. Your doctor has prescribed U- Molin Tablets for you and your condition.

- Things to be careful of

Always take this medicine exactly as your doctor has told you. Do not exceed the stated dose.

Paracetamol may cause severe skin reactions. Symptoms may include skin reddening, blisters or rash.

These could be signs of a serious condition. If these reaction occurs, stop use and seek medical assistance right away.

Side Effects

Like all medicines, U- Molin Tablets can have side effects, although not everybody gets them. The following side effects may happen with this medicine:

- allergic reactions- skin rash
- cutaneous hypersensitivity reactions including skin rashes, angioedema, Steven Johnson Syndrome / Toxic Epidermal Necrolysis have been reported.

If any of the side effects gets serious, or if you notice any side effects not listed in this leaflet, please tell your doctor or pharmacist.

Consult your doctor or pharmacist if you are in doubt or for any further information.

You may report any side effects or adverse drug reactions directly to the National Centre for Adverse Drug Reaction Monitoring by calling Tel: 03-78835550, or visiting the website npra.moh.gov.my

(Public→Reporting Medicinal Problems/ Side Effects/ AEFI/ Vaccine Safety)

Serial Number:
BPFK(R4/1)240717/00182

Storage and Disposal of U- Molin Tablets

- *Storage*

Do not use after the expiry date stated on the pack. The expiry date refers to the last day of that month.

Plastic Container: Store below 25°C

Blister Pack: Store below 30°C

Protect from light.

Keep the medicine out of reach of children.

- *Disposal*

Medicines should not be disposed of via wastewater or household waste. Ask your pharmacist how to dispose of medicines no longer required. These measures will help to protect the environment.

Product Description

- *What it looks like*

A blue, scored, oblong tablet of size 17 x 6.5mm with 'Life' marking.

- *Ingredients*

- *Active ingredient:*
Paracetamol.

- *Inactive ingredients:*

Corn starch, Magnesium Stearate, Povidone K-30, Sodium Starch Glycolate and Indigocarmine.

MAL number:

MAL19971334XZ

Manufacturer and Product Registration Holder

Malaysian Pharmaceutical Industries Sdn. Bhd.

Plot 14, Lebuhraya Kampung Jawa, 11900 Bayan Lepas, Pulau Pinang, Malaysia.

Date of Revision

26/07/2017