

# MILI SYRUP/SUSPENSION

Paracetamol (120mg/5ml, 250mg/5ml)

## What is in this leaflet

1. What Mili is used for
2. How Mili works
3. Before you use Mili
4. How to use Mili
5. While you are using Mili
6. Side effects
7. Storage and disposal of Mili
8. Product description
9. Manufactured by and Product Registration Holder
10. Date of revision

## What Mili is used for

For fever & relief of mild to moderate pain.

## How Mili works

Mili contains paracetamol, a medicine which works by relieving pain and reducing high temperature and fever.

## Before you use Mili

### -When you must not use it

- Do not take Mili if allergic to paracetamol or to other ingredients in it.
- Do not take any other Paracetamol containing medicine at the same time.
- Allergy alert: Paracetamol may cause severe skin reactions. Symptoms may include skin reddening, blisters or rash. These could be signs of a serious condition. If these reactions occur, stop use and seek medical assistance right away.

### -Before you start to use it

Caution if you have impaired kidney or liver function. It should also be given with care to individuals with alcohol dependence.

### Pregnancy and lactation

Talk to your doctor before taking this medicine if you are pregnant, might become pregnant or are breastfeeding.

### -Taking other medicines

Inform your doctor or pharmacist before taking these medicines, including any that you get without a prescription from your pharmacy, supermarket or health food shop.

Some medicines and Mili may interfere with each other. They may be affected by Mili or may affect how well it works especially:

- Enhance anticoagulant effect of warfarin (prevent heart attacks, strokes, and blood clots) and other coumarins with increased risk of bleeding.

## How to use Mili

### -How much to use

Always use Mili exactly as your doctor has prescribed. You should check with your doctor or pharmacist if in doubt.

### The usual dose:

#### Mili Syrup 120mg/5ml:

##### **Adults:**

10 to 15 mg/kg/dose orally every 4 to 6 hours.

##### **Children:**

**3 months to 1 year:** 2.5 to 5 ml/dose.

**1 year to 5 years:** 5 to 10 ml/dose.

These doses may be given every 4 to 6 hours if necessary up to a maximum of 4 doses in 24 hours

#### Mili Syrup 250mg/5ml:

##### **Adults:**

500mg every 4 hours as needed; or 1g every 4-6 hours up to 4 times a day.

##### **Children:**

**4-6 years:** 250mg every 4-6 hours as needed.

**6-9 years:** 375mg every 4-6 hours as needed

**9-12 years:** 500mg every 4-6 hours as needed.

#### Mili Suspension 250mg/5ml

**Above 12 years & adult:** 10 - 20 ml/dose

**1 to 5 years:** 2.5 - 5 ml/dose

**6 to 12 years:** 5 - 10 ml/dose

These doses may be repeated every 4-6 hours when necessary. Maximum of 4 doses in 24 hours.

### -When to use it

Use as directed by your doctor or pharmacist.

### -How long to use it

Continue taking Mili as your doctor recommended. If symptoms persist, consult your doctor.

### -If you forget to use it

Consult your doctor or pharmacist on what you should do if you forget to use it.

Take the missed dose as soon as you remember. If it is almost time for your

next dose, wait until then to take the medicine and skip the missed dose. Do not take a double dose to make up for the missed dose.

### -If you use too much (overdose)

Contact your doctor immediately or go to the Emergency Department of your nearest hospital, if you think you or anyone else may have taken too much of this medicine. Do this even if there are no signs of discomfort or poisoning. You may need urgent medical attention.

Taking too much Mili may cause:

- Vomiting, abdominal pain, low blood pressure and sweating. The most serious adverse effect of acute overdose of paracetamol is a dose-dependent, potentially fatal liver failure.

## While you are using Mili

### -Things you must do

- Inform your doctor or pharmacist if symptoms persist after taking this medication.
- Take your medicine exactly as your doctor has told you.
- Tell all the doctors, dentists and pharmacists treating you that you are taking Mili.
- Tell your doctor immediately if you become pregnant while taking this medication.

### -Things you must not do

- This preparation contains paracetamol. Do not take any other paracetamol containing medicines at the same time.
- Do not take more than the recommended dose unless your pharmacist or doctor tells you.
- Do not take any new medicines without consulting your doctor.
- Do not give Mili to anyone else, even if they have the same symptoms or condition as you.

### -Things to be careful of

- Do not take alcoholic beverages with this medicine.

## Side effects

Like all medicines, Mili can cause side effects, although not everybody gets them. If symptoms persist, do not take the medicine and refer to the doctor. Side effects of paracetamol are rare and usually mild.


# MILI SYRUP/SUSPENSION

Paracetamol (120mg/5ml, 250mg/5ml)

---

Tell your doctor if you notice any of the following side effects or notice any other effects not listed:

- Blood disease reactions.
- Cutaneous hypersensitivity reactions including skin rashes, angioedema (swelling beneath the skin's surface), Stevens Johnson Syndrome/ Toxic Epidermal Necrolysis have been reported.

You may report any side effects or adverse drug reactions directly to the National Centre for Adverse Drug Reaction Monitoring by calling Tel: 03-78835550, or visiting the website [npra.moh.gov.my](http://npra.moh.gov.my) (Public  Reporting Medicinal Problems / Side Effects / AEFI / Vaccine Safety).

## Storage and disposal of Mili

### -Storage

Keep container tightly closed. Protect from light.  
Keep out of reach of children.

### Mili Syrup 120mg/5ml:

Store below 30°C

### Mili Syrup 250mg/5ml:

Store between 15 -30°C

### Mili Suspension 250mg/5ml

Store below 30°C

### -Disposal

Medicines should not be disposed of via wastewater or household waste. Ask your pharmacist how to dispose of medicines no longer required. These measures will help to protect the environment.

## Product description

### -What it looks like

### Mili Syrup 120mg/5ml:

Red colour syrup with raspberry and strawberry flavour

### Mili Syrup 250mg/5ml:

Red colour syrup with raspberry and strawberry flavour.

### Mili Suspension 250mg/5ml

Pink suspension with raspberry flavour

## -Ingredients

-Active ingredient

Each 5ml contains

Paracetamol

-Inactive ingredients

Purified water, sugar syrup, saccharin sodium, raspberry flavour.

### Mili Syrup 120mg/5ml:

Strawberry flavour, glycerine, menthol crystal, propylene glycol, raspberry red, sodium benzoate, citric acid anhydrous, sorbitol solution 70%.

### Mili Syrup 250mg/5ml:

Strawberry flavour, glycerine, menthol crystal, propylene glycol, raspberry red, sodium benzoate, citric acid anhydrous, spirit chloroform.

### Mili Suspension 250mg/5ml

Spirit chloroform, methyl paraben, propyl paraben, gum tragacanth, sodium chloride, xanthan gum, guar gum, carmosine supra.

## -MAL number:

### Mili Syrup 120mg/5ml:

MAL19962480XZ

### Mili Syrup 250mg/5ml:

MAL19962481XZ

### Mili Suspension 250mg/5ml

MAL19962482XZ

## **Manufactured by and Product Registration Holder**

KCK Pharmaceutical Industries Sdn. Bhd.

Plot 61, Bayan Lepas Industrial Park Phase 4, 11900 Penang, Malaysia.

## **Date of revision**

13/09/2017

## **Serial Number**

NPRA(R4/1)130917/00248