

BECAME TABLET

Carbinoxamine Maleate/Pseudoephedrine Hydrochloride (4mg/60mg)

What is in this leaflet

1. What BECAME TABLET is used for
2. How BECAME TABLET works
3. Before you use BECAME TABLET
4. How to take BECAME TABLET
5. While you are using BECAME TABLET
6. Side effects
7. Storage and disposal of BECAME TABLET
8. Product description
9. Manufacturer and product registration holder
10. Date of revision

What BECAME TABLET is used for

BECAME TABLET is used to relief seasonal and allergic rhinitis (inflammation of the mucous membrane inside of the nose), and vasomotor rhinitis (non-allergic rhinitis) symptoms such as sneezing, stuff nose and running nose.

How BECAME TABLET works

Carbinoxamine Maleate is a type of antihistamines that works by blocking the action of histamines, biological substances in the body induced by specific allergens, Pseudoephedrine is a sympathomimetic agent that works by constricting blood vessels, leading to a reduction in mucosal edema (swelling) and nasal congestion.

Before you use BECAME TABLET

When you must not take it

If you are allergic towards any content of this medication or have lower respiratory tract diseases (eg, asthma), severe hypertension (high blood pressure), severe coronary artery (the vessels that supply the heart muscle with blood rich in oxygen) disease, urinary retention and peptic ulcer. You should avoid this medicine if you are taking monoamine oxidase inhibitors (MAOIs).

Before you start to take it

Consult your doctor and pharmacist regarding the consumption of BECAME TABLET if you are pregnant or breastfeeding.

Tell your doctor if you are allergic to any other medicines or you have any other health related problems such as hyperthyroidism (overactive thyroid), cardiovascular disease (heart disease), diabetes mellitus and closed-angle glaucoma (increased pressure inside the eye).

Taking other medicines

Avoid concurrent use of BECAME TABLET with sympathomimetic agent and monoamine oxidase inhibitors (MAOI). You must inform your doctor or pharmacist about all the medicines you are taking, including those without a prescription.

How to take BECAME TABLET

How much to take

Adult and children > 6 years:
1 tablet, three to four times daily
Children 18 months to 6 years:
½ tablet, three to four times daily

When to take it

BECAME TABLET should be taken according to the time advised by your doctor or pharmacist. It can be taken with or without food.

How long to take it

BECAME TABLET should be taken as your doctor advised you to do so.

If you forget to take it

If you miss a dose of BECAME TABLET and you are taking it regularly, take it as soon as possible. If it is approaching time for the next dose, wait until then to take the medicine and skip the missed dose. Do not take 2 doses at once.

If you take too much (overdose)

Contact your doctor immediately or go to the Emergency Department of your nearest hospital, if you think you or anyone else may have taken too much of this medicine. Do this even if there are no signs of discomfort or poisoning. You may need urgent medical attention.

While you are using BECAME TABLET

Things you must do

Follow your doctor's instruction when you are taking BECAME TABLET.

Things you must not do

Do not stop taking this medicine or change the dosing regimen without instruction from your doctor.

Do not give BECAME TABLET to anyone else, even if they have the same condition as you.

Do not give BECAME TABLET to premature infants.

Things to be careful of

Avoid alcohol consumption during the course of treatment with BECAME TABLET.

Avoid driving a car or operating potentially dangerous machinery after taking BECAME TABLET.

Side effects

BECAME TABLET may cause sedation, dizziness, diplopia (double vision), vomiting, diarrhea, dry mouth, headache, nervousness, nausea, anorexia (eating disorder), heartburn, weakness, polyuria (production of abnormally large volumes of dilute urine), dysuria (painful or difficult urination), convulsion, central nervous system stimulations, cardiac arrhythmias (an abnormal heart rhythm), respiratory difficulty (difficulty in breathing), increased heart rate or blood pressure, hallucinations, tremors, insomnia and pallor (an unhealthy pale appearance).

Please tell your doctor or pharmacist if you experience any other side effects.

You may report any side effects or adverse drug reactions directly to the National Centre for Adverse Drug Reaction Monitoring by calling Tel: 03-78835550, or visiting the website portal.bpfk.gov.my (Consumers→Reporting)

Storage and Disposal of BECAME TABLET

Storage



Store below 30°C, protect from light & moisture. Keep in a tight container.

Disposal

BECAME TABLET should be disposed when your doctor or pharmacist tells you to stop taking this tablet or it has passed the expiry date. Ask your pharmacist how to dispose medicines that are no longer required. Medicines should not be disposed of via wastewater or household waste. These measures will help to protect the environment.

Product description

What it looks like

BECAME TABLET is a white round tablet, with the diameter of 7.4mm one side impressed with the mark “” and the other side with a score “”.

Ingredients

Active ingredients-

Carbinoxamine Maleate
Pseudoephedrine Hydrochloride

Inactive ingredients-

Colloidal silicon dioxide
Sodium starch glycolate
Magnesium aluminometasilicate
Hydroxypropyl cellulose
Talc
Magnesium stearate
Ethanol

MAL number

MAL 19930600A

Manufacturer

Y.S.P. Industries (M) Sdn. Bhd.
Lot 3,5 & 7, Jalan P/7, Section 13, Kawasan
Perindustrian Bandar Baru Bangi, 43000
Kajang, Selangor Darul Ehsan, Malaysia.

Product Registration Holder

Y.S.P Industries (M) Sdn. Bhd.
Lot 3,5 & 7, Jalan P/7, Section 13,
Kawasan Perindustrian Bandar Baru Bangi,
43000 Kajang, Selangor Darul Ehsan,
Malaysia.

Date of revision :

30 October 2015

Serial Number

BPFK(R4/1)301015/00389