

PEACE SYRUP

Triprolidine hydrochloride/Pseudoephedrine hydrochloride (1.25mg/5mL / 30mg/5mL)

What is in this leaflet

1. What PEACE SYRUP is used for
2. How PEACE SYRUP works
3. Before you use PEACE SYRUP
4. How to use PEACE SYRUP
5. While you are using it
6. Side effects
7. Storage and disposal of PEACE SYRUP
8. Product description
9. Manufacturer and Product Registration Holder
10. Date of revision

What PEACE SYRUP is used for

PEACE SYRUP is used for symptomatic relief of sneezing, stuffy nose and runny nose caused by seasonal allergic and other unknown causes.

How PEACE SYRUP works

Triprolidine belongs to a class of medications known as antihistamine. It works by blocking the action of histamines, biological substances in the body that produces symptoms of sneezing, itching, watery eyes and runny nose. Pseudoephedrine is a sympathomimetic agent that works by constricting blood vessels, leading to a reduction in mucosal swelling and nasal congestion.

Before you use PEACE SYRUP

-When you must not use it

You must not take PEACE SYRUP if you have disease relating to heart or blood vessels, endocrine disorder (hormone) or sensitive towards this preparation of medication. People on monoamine oxidase inhibitor (MAO) treatment or within 2 weeks of stopping it should avoid this medication.

-Before you start to use it

Consult your doctor and pharmacist regarding the consumption of PEACE SYRUP if you are pregnant or breastfeeding. Tell your doctor or pharmacist if you are allergic to any other medicines or you have any other health related problems such as:

- narrow-angle glaucoma (angle closure eye disease due to increased pressure inside eyes)
- pyloroduodenal obstruction (gastric outflow obstruction)
- prostatic hypertrophy (enlargement of prostate gland)
- diabetes (high blood glucose in blood)
- ischemic heart disease
- hyperthyroidism (overactive thyroid hormone)
- severe impairment of kidney or liver function.

-Taking other medicines

Tell your doctor if you are taking any other medicines, including any that you buy without a prescription from a pharmacy, supermarket or health food.

Avoid concurrent use of PEACE SYRUP with:

- monoamine oxidase inhibitors (MAOIs), medicines to treat depression.
- narcotic analgesic, general anesthetics, medicines that use to relieved pain.
- tranquilizers, sedative hypnotics, medicine that use to treat mental illness.
- alcohol
- surgical skeletal muscle relaxant, medicine to relax the skeletal muscle function during surgery
- other central nervous system depressant.
- sympathomimetic agents.

How to use PEACE SYRUP

-How much to use

Adults and children ≥ 12 years:

10mL, 3 times daily

Children 6 ~ 12 years:

5mL, 3 times daily

Children 2 ~ 5 years:

2.5mL, 3 times daily

-When to use it

PEACE SYRUP should be taken according to the time advised by your doctor or pharmacist. It can be taken with or without food.

-How long to use it

PEACE SYRUP should be taken regularly and as long as your doctor advised you to do so.

-If you forget to use it

If you miss a dose of PEACE SYRUP and you are taking it regularly, take it as soon as possible. If several hours have passed or if it is approaching time for the next dose, do not double the dose to make up for the forgotten dose, unless advised by your health care provider. Do not take 2 doses at once.

-If you use too much (overdose)

Seek for medical attention.

Symptoms of acute toxicity of triprolidine include drowsiness, dizziness, incoordination (unstable in coordination), weakness, convulsion (involuntary muscle contraction) and breathing difficulty.

While for pseudoephedrine overdose are central nervous system stimulation leading to excitement, restlessness, rapid speech, hallucinations (hearing voices or seeing things which are not there), hypertonicity (higher muscle tension) and hyperreflexia (overactive, or overresponsive reflexes), dilated pupils and tachycardia (abnormal rapid heart beat). In severe poisoning the tachycardia may lead to cardiac arrhythmias (irregular heart beat) which may be further potentiated by the development of hypokalemia (deficiency potassium in blood).

While you are using PEACE SYRUP

-Things you must do

Follow your doctor's instruction when you are taking PEACE SYRUP. It should only be given to pregnant or nursing mother under doctor's supervision.

-Things you must not do

Do not stop taking this medicine or change the dosing regimen without instruction from your doctor. Do not give PEACE SYRUP to anyone else, even if they have the same condition as you.

Do not use in children below 2 years old.

-Things to be careful of

Avoid alcohol consumption during the course of treatment with PEACE SYRUP. Be careful if you drive or do anything that requires you to be alert and awake after consuming PEACE SYRUP. It should be used with caution in children between 2-6 years old.

Side effects

PEACE SYRUP may cause dryness of mouth, nose and throat, sedation, sleepiness, dizziness, disturbed co-ordination, hypotension (low blood pressure), headache, palpitation (feeling of fast or irregular heart beat), tachycardia (abnormal rapid heart rate), epigastric distress (pain at the epigastric region), anorexia (excessive weight loss due to eating disorder), nausea, vomiting and thickening of bronchial secretions (phlegm). Please tell your doctor or pharmacist if you experience any other side effects.

You may report any side effects or adverse drug reactions directly to the National Centre for Adverse Drug Reaction Monitoring by calling Tel: 03-78835550, or visiting the website portal.bpfk.gov.my
(Consumers → Reporting)

Storage and Disposal of PEACE SYRUP

-Storage

Store at temperature below 30°C, protect from light & moisture.

-Disposal

PEACE SYRUP should be disposed when your doctor or pharmacist tells you to stop taking this tablet or it has passed the expiry date. Ask your pharmacist how to dispose medicines that are no longer required.

Product description

-What it looks like

PEACE SYRUP is a clear, slight orange solution.

-Ingredients

-Active ingredient:

Triprolidine Hydrochloride
Pseudoephedrine Hydrochloride

-Inactive ingredients:

Sugar
Sodium Citrate
Sodium Benzoate
Sunset Yellow
Veltol Plus
Orange Powder

-MAL number

MAL19985561AZ

Manufacturer

Y.S.P. Industries (M) Sdn. Bhd.
Lot 3,5 & 7, Jalan P/7, Section 13, Kawasan Perindustrian Bandar Baru Bangi, 43000 Kajang, Selangor Darul Ehsan, Malaysia.

Product Registration Holder

Y.S.P Industries (M) Sdn. Bhd.
Lot 3,5 & 7, Jalan P/7, Section 13,
Kawasan Perindustrian Bandar Baru Bangi,
43000 Kajang, Selangor Darul Ehsan,
Malaysia.

Date of revision

21/3/2016

Serial Number

BPFK(R4/1)180316/00081