

# FREEDERM LOTION

Consumer Medication Information Leaflet (RiMUP)

Terbinafine Hydrochloride (1%w/v)

## What is in the leaflet

1. What Freederm Lotion is used for
2. How Freederm Lotion works
3. Before you use Freederm Lotion
4. How to take Freederm Lotion
5. While you are using it
6. Side Effects
7. Storage and Disposal of Freederm Lotion
8. Product Description
9. Manufacturer and Product Registration Holder
10. Date of revision

## What Freederm Lotion is used for

Freederm Lotion is an antifungal treatment and other fungal skin infections – athlete’s foot (tinea pedis), jock itch (tinea crucis), ringworm (tinea corporis), Pityriasis versicolor and fungal nail infection (cutaneous candidiasis).

## How Freederm Lotion works

Freederm Lotion has a broad-spectrum antifungal activity. Terbinafine works by preventing the formation of substances needed by the fungus to grow and develop, which causes weakness and death of the fungus.

## Before you use Freederm Lotion

### - When you must not take it

Do not take if you are hypersensitivity to Terbinafine HCl or any other ingredients in this product.

### - Before you start to use it

For external use only.

Consult doctor or pharmacist before using if you are pregnant or breast-feeding.

### - Taking other medicines

Terbinafine does not generally affect the activity of other medications.

## How to take Freederm Lotion

### - How much to take

Cleanse and dry the infected area thoroughly with warm water before application. Apply a thin layer onto the infected area and its surroundings once daily. Rub in gently. Treatment can be repeated if necessary.

### - When to take it

Take it when required.

### - How long to take it

Duration of treatment is one to two weeks. To prevent recurrence, continue to treat the area for one to two weeks after signs or symptoms of infections have gone.

### - If you forget to take it

If you forget to take Freederm Lotion, take it as soon as you remember.

### - If you take too much (overdose)

Overdose is unlikely occur. In the event of accidental ingestion, the affected person should be send to the nearest hospital for medical advice.

## While you are using Freederm Lotion

### - Things you must do

Check with your doctor if symptoms persist or getting worse.

### - Things you must not do

Do not take Freederm Lotion to treat any other conditions unless your doctor or pharmacist tells you so.

### - Things to be careful of

Avoid contact with eyes, nose and mouth. Not to be taken.

## Side Effects

Occasionally itching or redness may occur.

You may report any side effects or adverse drug reactions directly to the National

Centre for Adverse Drug Reaction Monitoring by calling Tel: 03-78835550, or visiting the website [npra.moh.gov.my](http://npra.moh.gov.my) (Public Reporting Medicinal Problems / Side Effects / AEFI / Vaccine Safety).

## Storage and Disposal of Freederm Lotion

### - Storage

- Store below 30°C.
- Protect from light.
- Keep out of reach of children.

### - Disposal

Do not take this after the expiry date. An expiry date is printed on the label and box of the medicine.

## Product Description

### - What it looks like

Freederm Lotion is a white to off-white gel-like solution.

### - Ingredients

Active ingredient

Terbinafine HCl 1%w/v

Inactive ingredients

Butylated Hydroxytoluene, Polysorbate 80, Isopropyl Alcohol, Benzyl Alcohol, Menthol, Isoprop Myristate, Purified Water, NAOH 1M, N-methyl-2-pyrrolidine, Carbomer 940, Hydrogenated Polyisobutene.

### - MAL Number:

MAL07125186 X

## Manufacturer and Product

### Registration Holder

Winwa Medical Sdn Bhd  
1952, Lorong I.K.S. Bukit Minyak 3,  
Taman I.K.S. Bukit Minyak,  
14000 Bukit Mertajam,  
Pulau Pinang, Malaysia.

## Date of revision

9/03/2018

Serial Number:

NPRA(R4/1)220218/00057

X.EYL-PIL.01