

CHLORPROMAZINE TABLET 25 mg® AND 100 mg®

CHLORPROMAZINE TABLET 25 mg and 100 mg

<p><u>What is in this leaflet:</u></p> <ol style="list-style-type: none">1. What Chlorpromazine Tablet 25 mg and 100 mg is used for2. How Chlorpromazine Tablet 25 mg and 100 mg works3. Before you use Chlorpromazine Tablet 25 mg and 100 mg4. How to take Chlorpromazine Tablet 25 mg and 100 mg5. While you are using Chlorpromazine Tablet 25 mg and 100 mg6. Side Effects7. How to store and dispose8. Product description <p><u>What Chlorpromazine Tablet 25 mg and 100 mg is used for</u></p> <p>Chlorpromazine Tablet 25 mg and 100 mg is used in the treatment of behavioural disturbances. It is also used for the short term treatment of anxiety, agitation and violent or impulsive behaviour. It is used to treat persistent hiccup and nausea and vomiting associated with terminal illness.</p> <p><u>How Chlorpromazine Tablet 25 mg and 100 mg works</u></p> <p>It contains chlorpromazine. Chlorpromazine is thought to work by blocking the action of a chemical called dopamine which acts on the nervous system in the brain.</p> <p><u>Before you use Chlorpromazine Tablet 25 mg and 100 mg</u></p> <p>Do not take Chlorpromazine Tablet 25 mg and 100 mg and tell your doctor if you have:</p> <ol style="list-style-type: none">1. Allergic to chlorpromazine tablet.2. Pregnant or breast-feeding.3. History of blood problem.4. Severe heart disease.5. Dulled senses such as feeling sleepy or uncoordinated, having blurred vision, slurred speech or being less aware of your surroundings. <p><u>How to take Chlorpromazine Tablet 25 mg and 100 mg</u></p> <p><u>How much to take</u></p> <p>Always take Chlorpromazine Tablet 25 mg and 100 mg exactly as your doctor has told you. If you are not sure, check with your doctor or pharmacist.</p>	<p>Doses:</p> <p>In most patients oral treatment may be used from the start, commencing with a dosage of 25 to 50 mg thrice daily and increasing as necessary. The usual daily maintenance dose ranges from 25 to 100 mg thrice daily.</p> <p>Psychotic patients may require daily doses of up to 1 g or more.</p> <p>In the treatment of mild alcohol withdrawal, agitated patients may be given 25 to 50 mg twice daily by mouth. Patients with less severe withdrawal symptoms may be given 30 to 75 mg daily in divided doses by mouth.</p> <p>Chlorpromazine should not be given to stuporous patients.</p> <p>Paediatric use of the tablets is not recommended.</p> <p><u>When to take it</u></p> <p>Swallow the tablets with water. If you are taking antacids, you should take at least 2 hours after taking chlorpromazine.</p> <p><u>If you forget to take it</u></p> <p>If you miss a dose of Chlorpromazine Tablet 25 mg and 100 mg, take it as soon as possible. Unless if nearly time for next dose, skip the missed dose and go back to your regular dosing schedule. Do not take 2 doses at once.</p> <p><u>If you take too much (overdose)</u></p> <p>If you have taken more than the prescribed dose contact your doctor immediately. Signs of an overdose include quick and shallow breaths, low body temperature, low blood pressure, restlessness, twisting of your limbs, fits, unusual heart beats and coma.</p> <p><u>While you are using Chlorpromazine Tablet 25 mg and 100 mg</u></p> <ol style="list-style-type: none">1. Do not take more than the recommended dose.2. You are advised to avoid alcohol.3. Chlorpromazine is incompatible with ampicillin, atrophine sulphate, benzylpenicillin, chloramphenicol, heparin, hydrocortisone hydrogen succinate, metaraminol, methicillin, oxytetracycline, thiopentone sodium, and vitamin B and C injection. <p><u>Side Effects</u></p> <p>Side effects include drowsiness, dryness of mouth, nasal congestion, postural hypotension, lowering of body temperature, rapid heart beat, abnormal</p>	<p>heart rhythm, agitation, insomnia, depression, blurred vision, convulsions, skin become sensitive to light, skin rashes and inhibition of ejaculation.</p> <p><u>How to store and dispose</u></p> <p><u>Storage</u></p> <p>Keep container tightly closed in a dry place, below 25°C. Protect from light. Keep out of the reach of children.</p> <p><u>Disposal</u></p> <p>Do not dispose the medicine into wastewater or household garbage. Ask your pharmacist how to dispose of medicines no longer required.</p> <p><u>Product Description</u></p> <p><u>What it looks like?</u></p> <p>Chlorpromazine Tablet 25 mg is a round and orange sugar coated tablet. The pack sizes are: Plastic Container of 20 tablets and Plastic Container of 500 and 1000 tablets (for Tender or Export only).</p> <p>Chlorpromazine Tablet 100 mg is a round and white film coated tablet. The pack sizes are: Plastic Container of 20 tablets, Plastic Container of 500 and 1000 tablets (for Tender or Export only), blister of 2 strips of 10 tablets and blister of 50 strips of tablets..</p> <p><u>Ingredients</u></p> <p>The active substance is chlorpromazine either 25 or 100 mg per tablet.</p> <p>The other ingredients chlorpromazine 25 mg are lactose, povidone, sodium starch glycolate, magnesium stearate, talc powder, acacia, light calcium carbonate, maize starch, titanium dioxide, light magnesium carbonate, color FDC yellow No. 6 lake, plain syrup, PVP shellac, beeswax and carbowax.</p> <p>The other ingredients chlorpromazine 100 mg are lactose, povidone, ethanol 96%, sodium starch glycolate, magnesium stearate, talc powder, acacia, light calcium carbonate, maize starch, plain syrup, titanium dioxide, colour FDC yellow no. 6 lake.</p> <p><u>Registration Numbers:</u></p> <p>Chlorpromazine Tablet 25 mg: MAL19860201A Chlorpromazine Tablet 100 mg: MAL19860027A</p> <p><u>Manufactured by:</u></p> <p>Idaman Pharma Manufacturing Sdn Bhd (661901-P) Lot 24, Bakar Arang Industrial Estate, Sungai Petani, Kedah, Malaysia.</p> <p>Marketing Authorisation Holder</p>
---	--	---

Date of revision:

December 13th, 2012

IDAMAN PHARMA MANUFACTURING SDN BHD