

# HOEMAL JUNIOR ORAL SUSPENSION

Paracetamol (120mg/ 5ml)

---

## What is in this leaflet?

1. What HOEMAL JUNIOR is used for?
2. How HOEMAL JUNIOR works?
3. Before you use HOEMAL JUNIOR
4. How to use HOEMAL JUNIOR
5. While you are using HOEMAL JUNIOR
6. Side effects
7. Storage and Disposal of HOEMAL JUNIOR
8. Product description
9. Manufacturer and Product Registration Holder
10. Date of Revision

This leaflet answers some common questions about HOEMAL JUNIOR. It does not contain all the available information. It does not take the place of talking to your pharmacist or doctor.

All medicines have risks and benefits. Your pharmacist or doctor has weighed the risks of you taking HOEMAL JUNIOR against the benefits they expect it will have for you.

**If you have any concerns about taking this medicine, ask your pharmacist or doctor.**

**Keep this leaflet with the medicine.**

You may need to read it again.

## What HOEMAL JUNIOR is used for?

HOEMAL JUNIOR is indicated for the relief of most mild to moderate pain including teething pain and for the symptomatic relief of fever.

## How HOEMAL JUNIOR works?

HOEMAL JUNIOR contains active ingredient Paracetamol which has analgesic (painkiller) and antipyretic (relieves fever) effects.

## Before you use HOEMAL JUNIOR

### When you must not use it

This preparation contains PARACETAMOL. Do not take any other PARACETAMOL-containing medicines at the same time.

Do not take HOEMAL JUNIOR if you are allergic to Paracetamol or any components in this preparation.

### Before you start to use it

Tell your pharmacist or doctor if you have:

- Liver problems
- Kidney problems

### **Pregnancy and breast feeding:**

Talk to your doctor before taking this medicine if you are pregnant, might become pregnant or are breast-feeding.

### Taking other medicines

Tell your pharmacist or doctor if you are taking any other products (such as over the counter drug, prescribed drug and herbal products).

If you are undergoing anticoagulant therapy, tell your doctor or pharmacist before taking HOEMAL JUNIOR.

## How to use HOEMAL JUNIOR

### How much to use

10 – 15 mg/kg/dose every 4 to 6 hours

### **3 – 6 months:**

½ teaspoonful (2.5ml), 3-4 times daily

### **6 months – 2 years:**

1 teaspoonful (5ml), 3-4 times daily

### **2 - 4 years:**

1 ½ teaspoonfuls (7.5 ml), 3-4 times daily

### **4 – 6 years:**

2 teaspoonfuls (10ml), 3-4 times daily

Do not take more than 4 times in 24 hours and never exceed the recommended dose unless prescribed by your doctor or pharmacist.

### When to use it

HOEMAL JUNIOR can be taken before or after meal.

### How long to use it

Continue taking HOEMAL JUNIOR for as long as your doctor recommends.

### If you forget to use it

If you forget to take the dose at the right time, use it as soon as you remember, then carry on as before. Do not double the dose to catch up.

If you use too much (overdose)

The symptoms of overdose of HOEMAL JUNIOR include vomiting, abdominal pain, hypotension and sweating. This is because, too much medicine can cause delayed liver damage.

Immediately telephone your doctor for advice or go to Accident and Emergency at the nearest hospital, if you think that you or anyone else may have taken too much HOEMAL JUNIOR.

**While you are using HOEMAL JUNIOR**

Things you must do

Take your medicine exactly as your doctor has told you.

Tell all the doctors and pharmacists treating you that you are taking HOEMAL JUNIOR.

Things you must not do

Do not take HOEMAL JUNIOR to treat any other complaints unless your doctor tells you to.

Things to be careful of

Paracetamol may cause severe skin reactions. Symptoms may include skin reddening, blisters or rash. These could be signs of a serious condition. If these reactions occur, stop use and seek medical assistance right away.

If you have impaired kidney or liver function or undergoing anticoagulant therapy, consult your doctor or pharmacist before taking HOEMAL JUNIOR.

**Side effects**

Like all medicines, HOEMAL JUNIOR can cause side effects, although not everybody gets them.

Any side effects are usually mild. Cutaneous hypersensitivity reactions including skin rashes, angioedema (swelling and itching of the skin), Steven Johnson Syndrome/ Toxic Epidermal Necrolysis (Blistering of the skin) have been reported.

You may report any side effects or adverse drug reactions directly to the National Centre for Adverse Drug Reaction Monitoring by calling Tel: 03-78835550, or visiting the website [npra.moh.gov.my](http://npra.moh.gov.my) (Public Reporting Medicinal Problems / Side Effects / AEFI / Vaccine Safety.)

**Storage and Disposal of HOEMAL JUNIOR**

Storage

**Keep container well-closed.  
Store below 30°C.  
Protect from strong light.  
Keep out of reach of children.**

Disposal

Dispose the medication after expiry date. Consult your pharmacist on the safety disposal of any medicines no longer required.

**Product description**

What it looks like

HOEMAL JUNIOR is a translucent red viscous liquid with strawberry flavour, available in 60ml plastic bottle.

Ingredients

Active ingredient:  
Paracetamol 120mg/5ml

Inactive ingredients:  
Methyl Paraben, Xanthan Gum, Hydroxyethylcellulose, Glycerin, Polysorbate 80, Sodium Citrate, Citric Acid Anhydrous, Ponceau 4R, Monoammonium Glycyrhizinate, Sucralose, Strawberry flavor & Purified water.

MAL number: **MAL12015006XZ**

**Manufacturer and Product Registration Holder**

Manufacturer

HOE Pharmaceuticals Sdn. Bhd.  
Lot 10, Jalan Sultan Mohamed 6,  
Bandar Sultan Suleiman,  
42000 Port Klang, Selangor,  
MALAYSIA.

Product Registration Holder

HOE Pharmaceuticals Sdn. Bhd.  
Lot 10, Jalan Sultan Mohamed 6,  
Bandar Sultan Suleiman,  
42000 Port Klang, Selangor,  
MALAYSIA.

**Date of revision**

06/03/2017

**Serial Number**

BPFK(R4/1)060317/00010