

REXOMTM MIST. MAGNESIUM TRISILICATE

Magnesium Trisilicate/Light Magnesium Carbonate/Sodium Bicarbonate (250mg/250mg/250mg per 5ml)

What is in this leaflet

1. What REXOMTM Mist. Magnesium Trisilicate is used for
2. How REXOMTM Mist. Magnesium Trisilicate works
3. Before you use REXOMTM Mist. Magnesium Trisilicate
4. How to use REXOMTM Mist. Magnesium Trisilicate
5. While you are using REXOMTM Mist. Magnesium Trisilicate
6. Side effects
7. Storage and disposal of REXOMTM Mist. Magnesium Trisilicate
8. Product description
9. Manufacturer and Product Registration Holder
10. Date of revision

What REXOMTM Mist. Magnesium Trisilicate is used for

For the symptomatic relief of upset stomach associated with hyperacidity (heartburn, acid indigestion, and sour stomach); hyperacidity associated with gastric and intestinal ulcers and in the management of esophagus inflammation due to reflux of gastric content.

How REXOMTM Mist. Magnesium Trisilicate works

This medicine contains antacids to reduce excess stomach acid.

Before you use REXOMTM Mist. Magnesium Trisilicate

-When you must not use it

- Should not be taken with tetracycline as it may interfere with its absorption.
- Do not take REXOMTM Mist. Magnesium Trisilicate if allergic to the ingredients in it.

-Before you start to use it

Pregnancy and lactation

Talk to your doctor before taking this medicine if you are pregnant, might become pregnant or are breastfeeding.

-Taking other medicines

Inform your doctor or pharmacist before taking this medicines, including any that you get without a prescription from your pharmacy, supermarket or health food shop.

Some medicines and REXOMTM Mist. Magnesium Trisilicate may interfere

with each other. They may be affected by REXOMTM Mist. Magnesium Trisilicate or may affect how well it works.

- It may interfere with the absorption of tetracycline (antibiotic).

How to use REXOMTM Mist. Magnesium Trisilicate

-How much to use

Always use REXOMTM Mist. Magnesium Trisilicate exactly as your doctor has prescribed. You should check with your doctor or pharmacist if in doubt.

The usual dose:

10 ml orally, after or between meals and bedtime.

-When to use it

To be used 4 times daily. Use as directed by your doctor or pharmacist.

-How long to use it

Continue taking REXOMTM Mist. Magnesium Trisilicate as your doctor recommended. If symptoms persist, consult your doctor.

-If you forget to use it

Consult your doctor or pharmacist on what you should do if you forget to use it.

Take the missed dose as soon as you remember. If it is almost time for your next dose, wait until then to take the medicine and skip the missed dose. Do not take a double dose to make up for the missed dose.

-If you use too much (overdose)

Contact your doctor immediately or go to the Emergency Department of your nearest hospital, if you think you or anyone else may have taken too much of this medicine. Do this even if there are no signs of discomfort or poisoning. You may need urgent medical attention.

Taking too much REXOMTM Mist. Magnesium Trisilicate may cause:

- Hypermagnesemia (high level of magnesium in the blood).
- Hypernatremia (rise in serum sodium concentration)
- High level of sodium and water.

While you are using REXOMTM Mist. Magnesium Trisilicate

-Things you must do

- Tell all the doctors, dentists and pharmacists treating you that you are taking REXOMTM Mist. Magnesium Trisilicate.
- Tell your doctor immediately if you become pregnant while taking this medication.

-Things you must not do

- Do not take more than the recommended dose unless your pharmacist or doctor tells you.
- Do not take any new medicines without consulting your doctor or pharmacist.
- Do not give REXOMTM Mist. Magnesium Trisilicate to anyone else, even if they have the same symptoms or condition as you.

-Things to be careful of

Be careful driving or operating machinery until you know how REXOMTM Mist. Magnesium Trisilicate affects you.

Side effects

Like all medicines, REXOMTM Mist. Magnesium Trisilicate can cause side effects, although not everybody gets them.

If side effects persist, do not take the medicine and refer to the doctor.

Tell your doctor if you notice any of the following side effects or notice any other effects not listed:

- Diarrhea
- Mucosal irritation
- Hypermagnesemia (high level of magnesium in the blood) if kidney function is impaired.

You may report any side effects or adverse drug reactions directly to the National Centre for Adverse Drug Reaction Monitoring by calling Tel: 03-78835550, or visiting the website npra.moh.gov.my (Public → Reporting Medicinal Problems / Side Effects / AEFI / Vaccine Safety).

REXOMTM MIST. MAGNESIUM TRISILICATE

Magnesium Trisilicate/Light Magnesium Carbonate/Sodium Bicarbonate (250mg/250mg/250mg per 5ml)

Storage and disposal of REXOMTM Mist. Magnesium Trisilicate

-Storage

Keep container tightly closed. Store between 15 - 30°C. Protect from light. Keep medicine out of reach of children.

-Disposal

Medicines should not be disposed of via wastewater or household waste. Ask your pharmacist how to dispose of medicines no longer required. These measures will help to protect the environment.

Product description

-What it looks like

White suspension with peppermint flavour.

-Ingredients

-Active ingredient

Each 5ml contains
250mg Magnesium trisilicate, 250mg
Light magnesium carbonate and 250mg
Sodium bicarbonate,

-Inactive ingredients

Chloroform, Peppermint oil, Saccharin sodium, Sodium propyl paraben, Cetylpyridinium chloride, Sodium methyl paraben, Xanthan gum and Purified water.

-MAL number

MAL19962486X

Manufacturer and Product Registration Holder

KCK Pharmaceutical Industries Sdn. Bhd.
Plot 61, Bayan Lepas Industrial Park
Phase 4, 11900 Penang,
Malaysia.

Date of revision

30/10/2017

Serial Number

NPRA(R4/1)301017/00296