

# PABRON COUGH KIDS SYRUP

Carbocisteine (100mg/ 5ml)

---

## What is in this leaflet

1. What PABRON COUGH KIDS is used for
2. How PABRON COUGH KIDS works
3. Before you use PABRON COUGH KIDS
4. How to use PABRON COUGH KIDS
5. While you are using PABRON COUGH KIDS
6. Side effects
7. Storage and Disposal of PABRON COUGH KIDS
8. Product description
9. Manufacturer and Product Registration Holder
10. Date of Revision

This leaflet answers some common questions about PABRON COUGH KIDS. It does not contain all the available information. It does not take the place of talking to your pharmacist or doctor.

All medicines have risks and benefits. Your pharmacist or doctor has weighed the risks of you taking PABRON COUGH KIDS against the benefits they expect it will have for you.

## Keep this leaflet with the medicine.

You may need to read it again.

## What PABRON COUGH KIDS is used for

PABRON COUGH KIDS aids expectoration by reducing mucus thickness, and therefore loosens and clears phlegm and relieve cough.

## How PABRON COUGH KIDS works

PABRON COUGH KIDS preparation is alcohol-free and sugar free.

**Carbocisteine** is a mucolytic. It alters the structure of mucus and decreases its thickness in order to facilitate its removal by expectoration.

Ask your pharmacist or doctor if you have any questions about this medicine.

## Before you use PABRON COUGH KIDS

### -When you must not use it

### **Do not use PABRON COUGH KIDS:**

- if you are allergic to Carbocisteine or other ingredients in this preparation.
- If you have active gastric or duodenal ulcers.

### -Before you start to use it

Tell your pharmacist or doctor:

### *Pregnancy and breast feeding:*

- Safe use of Carbocisteine during pregnancy and lactation has not been established.
- Ask your doctor or pharmacist for advice before taking this medicine if you are pregnant or breast-feeding.

### -Taking other medicines

Tell your pharmacist or doctor if you are taking any other products (such as over the counter product, prescribed medicine and herbal products).

During treatment, do not take any cough medicine or any medicine which dries airway secretion.

## How to use PABRON COUGH KIDS

### -How much to use

**Children above 5 years:**  
5ml

**Children below 5 years:**  
2.5ml

**Do not give this medicine to children under 2 years of age.**

### -When to use it

PABRON COUGH KIDS need to be taken three times daily or as directed by your doctor or pharmacist.

### -How long to use it

Continue taking PABRON COUGH for as long as your doctor recommends.

### -If you forget to use it

If you miss a dose, use it as soon as you remember. If it is near the time of the next dose, skip the missed dose and resume your usual dosing schedule. Do not double the dose to catch up.

### -If you use too much (overdose)

If you take too much of this medicine, consult your doctor or pharmacist. Bring any remaining medicine with you.

**While you are using PABRON COUGH KIDS**

-Things you must do

If you are about to start taking on any other new medicine, tell your pharmacist or doctor that you are taking PABRON COUGH KIDS.

-Things you must not do

Do not take more than the recommended dose unless your doctor tells you to.

Do not give PABRON COUGH KIDS to anyone else, even if they have the same symptoms or condition as you.

-Things to be careful of

This medicine may affect your ability to drive or use machines. If the tablets make you feel sick, dizzy or tired, or give you a headache, do not drive or use machines and contact your doctor immediately.

**Side effects**

PABRON COUGH KIDS may have unwanted side effects in some people. All medicines can have side effects. Sometimes they are serious, most of the time they are not.

Stop using this medication and tell your doctor immediately if any of these side effects occur: nausea, gastric discomfort, gastrointestinal bleeding and skin rash.

This is not a complete list of possible side effects. If you notice other effects not listed above, contact your doctor or pharmacist.

You may report any side effects or adverse drug reactions directly to the National Centre for Adverse Drug Reaction Monitoring by calling Tel: 03-78835549, or visiting the website

[npra.moh.gov.my](http://npra.moh.gov.my) (Public Reporting Side Effects to Medicines (ConSERF) or Vaccines (AEFI)).

**Storage and Disposal of PABRON COUGH KIDS**

-Storage

Store below 30°C and away from heat.  
Keep out of reach of children.

-Disposal

Dispose the medicine after the expiry date.  
Consult your pharmacist on how to safely dispose of any medicines no longer required.

**Product description**

-What it looks like

PABRON COUGH KIDS is a reddish purple syrup with blackcurrant flavor, available in 60ml, 90ml, 100ml and 120ml PET amber bottle.

-Ingredients

Active ingredient:  
Each 5ml syrup contains 100mg of Carbocisteine.

Inactive ingredients:  
Methyl Hydroxybenzoate, Sorbitol 70%, Propylene Glycol, Colour Powder Indigocarmine, Colour Powder Carmoisine, Flavour Blackcurrent, Sodium Hydroxide and Purified Water.

-MAL number: **MAL06041436XC**

**Manufacturer**

Xepa-Soul Pattinson (M) Sdn Bhd  
1-5 Cheng Industrial Estate  
75250 Melaka,  
Malaysia.

**Product Registration Holder**

HOE Pharmaceuticals Sdn. Bhd.  
Lot 10, Jalan Sultan Mohd. 6,  
Bandar Sultan Suleiman,  
42000 Port Klang, Selangor,  
Malaysia.

**Date of revision**

08/05/2018

**Serial Number:**

NPRA(R4/1)070518/00140