

PHARMANIAGA PARACETAMOL TABLET

Paracetamol (500mg, 650mg)

What is in this leaflet

1. What Pharmaniaga Paracetamol Tablet is used for
2. How Pharmaniaga Paracetamol Tablet works
3. Before you use Pharmaniaga Paracetamol Tablet
4. How to use Pharmaniaga Paracetamol Tablet
5. While you are using it
6. Side effects
7. Storage and Disposal of Pharmaniaga Paracetamol Tablet
8. Product Description
9. Manufacturer and Product Registration Holder
10. Date of Revision

What Pharmaniaga Paracetamol Tablet is used for

Pharmaniaga Paracetamol Tablet is used for the relief of minor aches, pains and fever.

How Pharmaniaga Paracetamol Tablet works

Pharmaniaga Paracetamol Tablet has analgesic (pain relief) and antipyretic (anti-fever) properties.

Before you use Pharmaniaga Paracetamol Tablet

-When you must not use it

Do not take Pharmaniaga Paracetamol Tablet if you are allergic to Paracetamol or Acetaminophen and other ingredients.

If you are not sure whether you should start taking this medicine, talk to your doctor or pharmacist.

- Before you start to use it

This product contains Paracetamol. Do not take other Paracetamol containing medicine at the same time.

Your doctor needs to know before you take Pharmaniaga Paracetamol Tablet if you have kidney or liver problem.

Pharmaniaga Paracetamol Tablet should also be taken with care when you take other medicines that affect the liver.

Discuss with your doctor or pharmacist first if you are pregnant or breastfeeding. Problems in human have not been reported when taking Pharmaniaga Paracetamol Tablet during pregnancy or breastfeeding.

-Taking other medicines

Tell your doctor or pharmacist before taking Pharmaniaga Paracetamol Tablet, if you are taking any other medicines, including any that you buy without a prescription from your pharmacy, supermarket or health food shop. The effects of some medicines may be affected if other medicines are used at the same time.

Inform your doctor if you are using Pharmaniaga Paracetamol Tablet in combination with the alcohol, liver enzyme inducers and hepatotoxic medications. Risk of liver toxicity increased with liver enzyme inducers and medicines that are known to be toxic to the liver.

How to use Pharmaniaga Paracetamol Tablet

Always take the medicine exactly as your doctor or pharmacist has told you. Read the direction on the labels carefully.

-How much to use

Pharmaniaga Paracetamol 500mg:

Adults: 1 to 2 tablets every 4 to 6 hours up to a maximum of 8 tablets daily.

Children: 6 to 12 years: 1/2 to 1 tablet 3 to 4 times daily.

Pharmaniaga Paracetamol 650mg :

Adults: 1 to 1_{1/2} tablets three or four times a day up to a maximum of 6 tablets daily.

-When to use it

Follow the direction label carefully and as instructed by your doctor or pharmacist.

-How long to use it

Follow the direction label carefully and as instructed by your doctor or pharmacist.

-If you forget to use it

Take the missed dose as soon as you remember. If it is almost time for your next dose, skip the dose you missed and take your next dose when you are meant to.

Do not take a double dose to make up for the dose that you missed. This may increase the chance of you getting unwanted side effects.

-If you use too much (overdose)

Immediately call your doctor or go to the emergency department at your nearest hospital if you or anyone else has taken a large amount of this medication.

Symptoms of overdosage of Pharmaniaga Paracetamol Tablet can be seen in in the first 24 hours. Among the symptoms are:

- nausea
- vomiting
- loss of appetite
- abdominal pain

Liver damage (with a symptoms of pain, tenderness, and/ or swelling in the upper abdominal area) may become more apparent 12 to 48 hours after the intake of the medicine.

In severe poisoning, liver failure may progress to brain malfunction or damage, coma and death.

Prompt treatment is essential in the management of Pharmaniaga Paracetamol Tablet overdosage.

PHARMANIAGA PARACETAMOL TABLET

Paracetamol (500mg, 650mg)

While you are using Pharmaniaga Paracetamol Tablet

-Things you must do

Take the medication exactly as your doctor prescribed.

- Tell all the doctors, dentists and pharmacists treating you that you are taking Pharmaniaga Paracetamol Tablet.
- Tell your doctor immediately if you become pregnant while taking this medication.

-Things you must not do

Do not give Pharmaniaga Paracetamol Tablet to anyone else, even if their symptoms seem similar to yours.

Do not take your medicine to treat any other complaints unless your doctor or pharmacist tells you to.

-Things to be careful of

Avoid drinking alcohol while taking this medication.

Side Effects

All medicines may have some unwanted side effects. Unwanted side effects do not always occur in every person. Tell your doctor or pharmacist as soon as possible if you do not feel well while you are taking Pharmaniaga Paracetamol Tablet.

- **Allergy alert:** Paracetamol may cause severe skin reactions. Symptoms may include skin reddening, blisters or rash.
- These could be signs of a serious condition. If these reactions occur, stop use and seek medical assistance right away.

You may report any side effects or adverse drug reactions directly to the National Centre for Adverse Drug Reaction Monitoring by calling Tel: 03-78835550 or visiting the website npra.moh.gov.my (Public → Reporting Side Effects to Medicines (ConSERF)/Vaccine (AEFI)).

Storage and Disposal of Pharmaniaga Paracetamol Tablet

-Storage

Keep all medicines out of sight and reach of children.

Do not take this medicine after the expiry date stamped on the pack.

- Store below 25°C.
- Protect from light.

-Disposal

Medicines should not be disposed of via wastewater or household waste. Ask your pharmacist how to dispose of medicines no longer required. These measures will help to protect the environment.

Product Description

-What it looks like

Pharmaniaga Paracetamol 500mg:

White, round, flat bevelled-edge with Pharmaniaga icon on one side and breakline on the other.

Pharmaniaga Paracetamol 650mg:

White, oblong-shaped tablets with Phamaniaga icon on one side and breakline on the other.

-Ingredients

-Active ingredient(s):
Paracetamol

-Inactive ingredients:

Potassium sorbate, Stearic acid, Pregelatinized Starch, Microcrystalline Cellulose, Croscarmellose Sodium, Povidone, Talc, and Purified Water.

-MAL Number(s)

Pharmaniaga Paracetamol 500mg:

MAL20000235XZ

Pharmaniaga Paracetamol 650mg:

MAL19990608XZ

Manufacturer

Pharmaniaga Manufacturing Berhad (60016-D)

11A, Jalan P/1, Kawasan Perusahaan Bangi, 43650, Bandar Baru Bangi, Selangor Darul Ehsan, Malaysia.

Product Registration Holder

Pharmaniaga Manufacturing Berhad (60016-D)

11A, Jalan P/1, Kawasan Perusahaan Bangi, 43650, Bandar Baru Bangi, Selangor Darul Ehsan, Malaysia.

Date of Revision

9/4/2018

Serial Number

NPRA(R4/1)140218/00049