

SOLCOSERYL® DENTAL ADHESIVE PASTE

Dried, Protein-Free, Biologica (2.125mg/g) / Polidocanol (10mg/g)

What is in this leaflet

1. What Solcoseryl Dental Adhesive Paste used for
2. How Solcoseryl Dental Adhesive Paste works
3. Before you use Solcoseryl Dental Adhesive Paste
4. How to use Solcoseryl Dental Adhesive Paste
5. While you are using Solcoseryl Dental Adhesive Paste
6. Side effects
7. Storage and Disposal of Solcoseryl Dental Adhesive Paste
8. Product Description
9. Manufacturer and Product Registration holder
10. Date of revision

What Solcoseryl Dental Adhesive Paste used for

Solcoseryl Dental Adhesive Paste is used for the local treatment of:

- Painful and inflammatory disorders of the mucus membranes of the mouth, of the gums and of the lips: mouth ulcer (aphthae), cold sore (herpes labialis), inflammation of the gums (gingivitis and periodontitis).
- Denture sores
- Problems associated with teething and eruption of wisdom teeth.
- As wound dressing after surgical treatment of the mouth such as: scaling, removing tissue via scraping or scooping (curettage), operations of tissues around the tooth (periodontal), extractions, fitting of immediate dentures, inflammatory of tooth socket (alveolitis).

How Solcoseryl Dental Adhesive Paste works

Solcoseryl® Dental Adhesive Paste contains heamodialysate from calves' blood to promote healing, and polidocanol to relieve pain quickly and lastingly. The paste forms a dressing which adheres for a long time to the

wound and protects it from irritation when eating and drinking.

Before you use Solcoseryl Dental Adhesive Paste

- When you must not use it

Do not use Solcoseryl Dental Adhesive Paste if you are hypersensitive or allergic to parahydroxybenzoates (E210, E216, E218) which are present as preservatives.

- Before you start to use it

Tell your doctor before you start to use Solcoseryl Dental Adhesive Paste if:

- The mucous membrane or the wound is infected
- You are suffering from other illnesses
- You have other allergies
- For acute infection it is advisable to rinse mouth with disinfectant solutions (chlorhexidine digluconate)

- Taking other medicines

Tell your doctor if you are taking any other medicines, including medicines you buy without a prescription from a pharmacy, supermarket or health food shop.

How to use Solcoseryl Dental Adhesive Paste

- How much to use

Unless otherwise prescribed, apply a ribbon of paste about ½ cm long thinly to the wound 3-5 times a day.

- When to use it

It is particularly advisable to be applied before retiring for the night.

- How long to use it

Solcoseryl Dental Adhesive Paste is applied until the wound is completely healed.

- If you forget to use it

Please contact your physicians.

- If you use too much (overdose)

Please contact your physicians.

While you are using Solcoseryl Dental Adhesive Paste

- Things you must do

Use the dosage indicated in the package insert or prescribed by the physician.

Solcoseryl Dental Adhesive Paste is best applied to the wound, which has been previously dried, using a cotton bud or cleaned fingertip. Subsequently either the paste should be smoothed with the moistened finger, or the mouth should be rinsed with a glass of water. The paste should not be massaged in.

For denture sore, the paste is applied to the cleaned and dried denture.

To improve adhesion of the denture, please rinse the mouth with a glass of water before inserting it. Close the tube tightly after use.

Close the tube tightly after use, the tube opening becomes blocked on exposure to moisture.

- Things you must not do

Solcoseryl Dental Adhesive Paste should not be packed into deep wound which are subsequently closed to a surgical suture.

Do not use the product after the date indicated by 'Exp' on the container.

- Things to be careful of

If you are or are attempting to become pregnant, or are breastfeeding, use the paste only if necessary.

Oil may be occasionally deposited at the mouth of the tube and likewise does not affect the quality of the product.

Side effects

To date no side effects have been reported with the use of Solcoseryl Dental Adhesive Paste.

If you think the effect of the medicine is too weak or too strong, speak to your physician or pharmacist.

You may report any side effects or drug reactions directly to the National Centre for Adverse Drug Reaction

Monitoring by calling Tel: 03-78835549, or visiting the website npra.moh.gov.my (Public → Reporting Side Effects to Medicines (ConSERF) or Vaccines).

47301 Petaling Jaya, Selangor Darul Ehsan, Malaysia.

Date of revision
03/07/2018

Storage and Disposal of Solcoseryl Dental Adhesive

Serial Number
NPRA(R4/1)210518/00162

- Storage

Solcoseryl Dental Adhesive Paste should be stored protected from heat at temperature below 30°C, out of the reach of children.

- Disposal

Medicines should not be disposed of via wastewater or household waste. Ask your pharmacist how to dispose of medicines no longer required. These measures will help to protect the environment.

Product description

- What it looks like

Solcoseryl Dental Adhesive Paste is pale beige, grainy, spreadable, homogeneous mass and has the characteristic odor of peppermint.

- Ingredients

Active Ingredients

- Polidocanol (10mg/g)
- Solcoseryl® (protein-free haemodialysate of blood from veal calves, chemically and biologically standardized), Dried, Protein-Free, Biologica (2.125mg/g)

Inactive Ingredients

- Methyl parahydroxybenzoate (1.80mg/g) (E 218)
- Propyl parahydroxybenzoate (0.20mg/g) (E216)
- Menthol, peppermint oil, sodium carboxymethylcellulose, paste base

- MAL number:

MAL19920090XC

Manufacturer

Legacy Pharmaceuticals Switzerland GmbH, Birsfelden, Switzerland.

Product Registration Holder:

A.Menarini Singapore Pte. Ltd.
B-18-2, Level 18, The Ascent
Paradigm,
No.1, Jalan SS7/26A, Kelana Jaya,