

# AXCEL IRIDIN PLUS LINCTUS

Pholcodine (5mg/5ml)

Tripolidine HCl (1.25 mg/5ml)

## WHAT IS IN THIS LEAFLET

1. What Axcel Iridin Plus Linctus is used for.
2. How Axcel Iridin Plus Linctus works.
3. Before you use Axcel Iridin Plus Linctus.
4. How to take Axcel Iridin Plus Linctus.
5. While you are using it.
6. Side effects.
7. Storage and disposal of Axcel Iridin Plus Linctus.
8. Product description.
9. Manufacturer and Product Registration Holder.
10. Date of revision.

## WHAT AXCEL IRIDIN PLUS LINCTUS IS USED FOR

Axcel Iridin Plus Linctus is used for the relief of unproductive or irritating coughs associated with common cold and flu-like symptoms such as runny nose, sneezing and watery eye.

## HOW AXCEL IRIDIN PLUS LINCTUS WORKS

Pholcodine is cough suppressant and has a mild sedative effect which relieves local irritation of the respiratory tract over 8 to 10 hours. Tripolidine is a competitive histamine H1-receptor antagonist. It blocks the effects of the naturally occurring chemical histamine in your body.

## BEFORE YOU USE AXCEL IRIDIN PLUS LINCTUS

### When you must not use it

Do not take Axcel Iridin Plus Linctus if:

- If you are allergic or hypersensitive pholcodine or tripolidine or any other component of Axcel Iridin Plus Linctus.
- Do not use this medicine in children under 2 years of age.

### Before you start to use it

Tell your doctor if:

- If you have respiratory problems.
- If have liver and renal impairment.
- If you are pregnant or lactating.

### Taking other medicines

You should tell your doctor if you are taking the following medicine as it may interact with your Axcel Iridin Plus Linctus:

- Alcohol
- Medicines used to treat depression
- Opioid analgesics
- Antimuscarinic drugs

## HOW TO USE AXCEL IRIDIN PLUS LINCTUS

### How much to use

For oral administration only.

Adult and children over 12 years: 10 ml, 3 times daily.

Children 6-12 years: 5 ml, 3 times daily.

Children 2-5 years: 2.5 ml, 3 times daily.

Please use measuring cup for dosing measurement.

### When to use it

Your doctor will decide when it is appropriate for you to take Axcel Iridin Plus Linctus according to the type of condition you have.

### How long to use it

- Take this medicine as stated in the leaflet without exceeding the dosage.
- Follow any recommendations and advices from your doctor or pharmacist.

### If you forget to use it

- Take the missed dose as soon as you remember. If it is almost time for your next dose, wait until then to take the medicine and skipped the missed dose.
- Do not double up the dosage as Axcel Iridin Plus Linctus may give you unacceptable side effects.

### If you use too much (overdose)

- Drowsiness, restlessness, excitement, ataxia (uncoordinated movement), antimuscarinic symptoms, central nervous system (CNS) depression with coma, convulsion, progressing to respiratory failure and

cardiovascular collapse may be seen following acute overdose with Axcel Iridin Plus Linctus.

- Treatment should be symptomatic.
- Necessary measures should be taken to maintain and support respiration and control convulsions.
- Gastric lavage may be performed if indicated.

It is advisable for you to contact and inform your doctor if you did take Axcel Iridin Plus Linctus too much.

## WHILE YOU ARE USING IT

### Things you must do

- Discuss with your doctor or pharmacist the dos' and don'ts' while taking this medicine.
- Tell your doctor if unexpectedly you need to take any other medicines.
- Take the Axcel Iridin Plus Linctus according to the recommended dosage.
- Tell your doctor or pharmacist if you experience any side effects that are not listed in this leaflet.

### Things you must not do

- Do not take your medicine more than the suggested frequency in a day.
- Do not over dosage the medicine during any single consumption.

### Things to be careful of

- If you have respiratory problems.
- If you have liver impairment.
- If you have kidney problem.
- If you are driving or use machine because it can cause drowsiness.
- If you are alcohol consumer or use centrally-acting sedatives.
- Caution when used for children 2 to 6 years old.
- If you are pregnant or lactating.

## SIDE EFFECTS

Like all medicines, Axcel Iridin Plus Linctus can cause side effects although not everybody gets them. You are advised to consult doctor/

# AXCEL IRIDIN PLUS LINCTUS

Pholcodine (5mg/5ml)

Triprolidine HCl (1.25 mg/5ml)

pharmacist for further information. The following side effects have been reported.

- Nausea
- Vomiting
- Drowsiness
- Skin rashes, with or without irritation.
- Dryness of mouth, nose, throat
- After large doses, restlessness, excitement, ataxia (uncoordinated movement) and respiratory depression may occur.

If any of these side effects become more serious, or you experience other side effects which are not mentioned in this leaflet, please inform your doctor or pharmacist.

You may report any side effects or adverse drug reactions directly to the National Centre for Adverse Drug Reaction Monitoring by calling Tel: 03-78835550, or visiting the website [portal.bpfk.gov.my](http://portal.bpfk.gov.my) (Consumers -> Reporting).

## STORAGE AND DISPOSAL OF AXCEL IRIDIN PLUS LINCTUS

### Storage

Keep Axcel Iridin Plus Linctus out of reach children.

Store below 30°C. Protect from light.

### Disposal

Do not use Axcel Iridin Plus Linctus after the expiry date which is stated on the carton.

Medicines should not be disposed of via wastewater or household waste. Ask your pharmacist how to dispose of medicines no longer required. These measures will help to protect the environment.

## PRODUCT DESCRIPTION

### What it looks like

Axcel Iridin Plus Linctus is purple colour liquid with fruity flavour.

### Ingredients

#### *Active Ingredient (s):*

Pholcodine and Triprolidine Hydrochloride

#### *Inactive Ingredients:*

Saccharin Sodium, Refined Sugar, Methylparaben, Glycerin, Methol Crystals, Citric Acid Anhydrous, Trisodium Citrate Dihydrate, Cherry Red, Brilliant Blue, Mixed Fruit Flavour and Purified water.

#### MAL Number

MAL09051551A

#### **MANUFACTURER AND PRODUCT REGISTRATION HOLDER**

KOTRA PHARMA (M) SDN BHD  
No. 1, 2 & 3, Jalan TTC 12,  
Cheng Industrial Estate,  
75250 Melaka, Malaysia.

#### **DATE OF REVISION**

28/08/2014

#### **SERIAL NUMBER**

BPFK(R4/1)030914/00004