SULBUTIAMINE 200 CAPSULE

Sulbutiamine (200mg)

What is in this leaflet

- 1. What Sulbutiamine is used for
- 2. How Sulbutiamine works
- 3. Before you use Sulbutiamine
- 4. How to use Sulbutiamine
- 5. While you are using it
- 6. Side effects
- 7. Storage and Disposal of Sulbutiamine
- 8. Product Description
- 9. Manufacturer and Product Registration Holder
- 10. Date of revision

What Sulbutiamine is used for

Sulbutiamine is used for treatment of all form of function asthenia (lack of strength but also disease) eg,

- Physical and intellectual overwork;
- Difficulty in concentration;
- Fatigue not overcome by rest;
- Decreased intellectual efficiency and lucidity, especially in the elderly.

How Sulbutiamine works

Sulbutiamine is a thiamine derivative. Sulbutiamine is a lipophilic molecule that crosses the blood brain barrier more easily than vitamin B1 and leads to an increase in the level of Thiamine triphosphate. It gives better physical resistance, improved intellectual performance and psychological balance.

Before you use Sulbutiamine

- When you must not use it Do not take Sulbutiamine if:
- You are allergic (hypersensitive) to Sulbutiamine or any of the other ingredients.

If you are pregnant, trying to get pregnant or think you may be pregnant, consult your doctor before taking this medicine.

If you are breastfeeding, consult your doctor before taking this medicine. This is because Thiamine is widely distributed to most body tissues and appears in breast milk.

- <u>Before you start to use it</u> You must consult a doctor or pharmacist if you:
 - Allergic

 (hypersensitivity) to
 Sulbutiamine or other ingredients in this product.
- <u>Taking other medicines</u> Consult a doctor or pharmacist if you are taking:
 - Loop diuretics medicines such as furosemide, ethacrynic acid and bumetanide.
 - Intake of thiamine with foods that contain sulfites (such as dried fruits,potato chips,jams and grape juice).
 - Intake of thiamin tea, coffee and decaffeinated coffee.

How to use Sulbutiamine

- How much to use

Follow all directions given in the insert carefully. If you do not understand the instructions on the label, ask your doctor or pharmacist for help.

1 to 3 capsules as a single dose or in divided dose.

<u>Adult</u>: The dose is 2 capsules daily with breakfast.

- <u>When to use it</u> Use as directed as in this leaflet.

- <u>How long to use it</u> As per the dosage recommendation
- If you forget to use it
 Take the missed dose as soon
 as you remember. If it is
 almost time for your next dose,
 wait until then to take the
 medicine and skip the missed
 dose. Do not take a double
 dose to make up for the missed
 dose

- <u>If you use too much</u> <u>(overdose)</u>

Contact your doctor or pharmacist for advice immediately.

Signs and symptoms of overdose include nervous excitement (agitation) with intense feelings of well-being (euphoria) and tremor in the extremities.

While you are using it

- <u>Things you must do</u>
Take your dose exactly as given in the leaflet.

Tell all the doctors, dentists and pharmacists that are treating you that you are taking Sulbutiamine.

Consult a doctor immediately if you are become pregnant while taking this medication.

- <u>Things you must not do</u> Do not give Sulbutiamine to anyone else, even if they have the same symptoms or condition as you.
- <u>Things to be careful of</u> Driving and using machines

This product does not affect your ability to drive or use machines.

SULBUTIAMINE 200 CAPSULE

Sulbutiamine (200mg)

Side effects

Like all medicines, Sulbutiamine can cause side effects, although not everybody gets them. The side effects are:

- Mild nervous excitement (agitation) in elderly person.
- Allergic skin reaction
- Unpleasant smell of urine.

Visit your doctor or pharmacist immediately if you experience any side effects after taking this Sulbutiamine.

You may report any side effects or adverse drug reactions directly to the National Centre for Adverse Drug Reaction Monitoring by calling Tel: 03-78835550, or visiting the website npra.moh.gov.my(Public → Reporting Medicinal Problems / Side Effects / AEFI / Vaccine Safety).

Storage and Disposal of Sulbutiamine

- <u>Storage</u> Store in a dry place below 30°C. Protect from light and

- Disposal

moisture.

Products should not be disposed of via wastewater or household waste. Ask your pharmacist how to dispose of Sulbutiamine that is no longer required. These measures will help to protect the environment.

Product Description

- <u>What it looks like</u> Two pieces, size '0' maroon color hard vegetable capsule contain white color powder.

- Ingredients
- Active ingredient(s) Sulbutiamine
- Inactive ingredients Sodium Starch Glycolate Corn Starch Colloidal Silicon Dioxide Microcrystalline Cellulose Magnesium Stearate
- <u>MAL number(s):</u> Sulbutiamine Capsule: MAL06111366X

Manufacturer

Pahang Pharmacy Sdn.Bhd. Lot 5979, Jalan Teratai, 5 ½ Miles Off, Jalan Meru, 41050 Klang, Selangor, Malaysia.

Product Registration Holder

Pahang Pharmacy Sdn.Bhd. Lot 5979, Jalan Teratai, 5 ½ Miles Off, Jalan Meru, 41050 Klang, Selangor, Malaysia.

Date of Revision 19/04/2017

Serial Number BPFK(R4/1)190417/00085