

AXCEL SERRATIOPEPTIDASE-5MG TABLET

Serratiopeptidase (5mg)

What is in this leaflet

1. What Axcel Serratiopeptidase-5mg Tablet is used for.
2. How Axcel Serratiopeptidase-5mg Tablet works.
3. Before you use Axcel Serratiopeptidase-5mg Tablet.
4. How to take Axcel Serratiopeptidase-5mg Tablet.
5. While you are using it.
6. Side effects.
7. Storage and disposal of Axcel Serratiopeptidase-5mg Tablet.
8. Product description.
9. Manufacturer and Product Registration Holder.
10. Date of revision.

What Axcel Serratiopeptidase-5mg Tablet is used for

Axcel Serratiopeptidase-5mg Tablet is used to relief of inflammation in the following diseases:

- After surgery and trauma.
- Engorgement of breast.
- Inflammation of the
 - Bladder.
 - Tube that carries sperm from the testes.
 - Gum tissue of the wisdom tooth.
 - Nasal sinus.
- Tooth abscess (pus).

It is also used in the treatment of difficulty of expectoration in:

- Bronchitis (inflammation of the airways that carry air to lungs).
- Pulmonary tuberculosis (infection of the lungs).
- Asthma.
- Disease after anesthesia.

How Axcel Serratiopeptidase-5mg Tablet works

Serratiopeptidase is a enzymes that helping the body in breakdown of protein. It has the ability to reduce pain by blocking the release of pain-inducing chemicals from inflamed tissues.

Before you use Axcel Serratiopeptidase-5mg Tablet

-When you must not use it

Do not take Axcel Serratiopeptidase-5mg Tablet if:

- You are allergy or hypersensitive to serratiopeptidase or to ingredients in the product.
- You are pregnant, trying to get pregnant or think you may be pregnant.
- You are breast-feeding. Ask your doctor or pharmacist for advice before taking any medicine.

-Before you start to use it

Tell your doctor if:

- You are having history of medicine allergy and blood abnormalities such as bleeding from the nose and bloody mucus.
- You are having severe liver or kidney dysfunction.

-Taking other medicines

You should tell your doctor if you are taking the following medicine as it may interact with your Axcel Serratiopeptidase-5mg Tablet:

- Anticoagulant (for blood thinning).

It may still be all right for you to take Axcel Serratiopeptidase-5mg Tablet and your doctor will be able to decide what is suitable for you.

How to use Axcel Serratiopeptidase-5mg Tablet

-How much to use

For oral administration only.

To be taken 1-2 tablets. Dosage should be adjusted according to the individual age and condition.

-When to use it

Take Axcel Serratiopeptidase-5mg Tablet three times daily after meal or as directed by physician. Your doctor will decide when it is appropriate for you to use Axcel Serratiopeptidase-5mg Tablet according to the type of condition you have.

-How long to use it

Continue taking Axcel Serratiopeptidase-5mg Tablet for as long as doctor recommends.

-If you forget to use it

Consult your doctor or pharmacist on what you should do if you forget to use it.

Take the missed dose as soon as you remember. If it is almost time for your next dose, wait until then take the medicine and skip the missed dose. Do not take a double dose to make up for the missed dose.

-If you use too much (overdose)

Contact your doctor immediately or go to the Emergency Department of your nearest hospital, if you think you or anyone else may have taken too much of this medicine. Do this even if there are no signs of discomfort or poisoning. You may need urgent medical attention.

While you are using it

-Things you must do

- Take your medicine exactly as your doctor has told you.
- Tell all the doctors, dentists and pharmacists treating you that you are taking Axcel Serratiopeptidase-5mg Tablet.
- Tell your doctor immediately if you become pregnant while taking this medication.

-Things you must not do

- Do not stop taking the medicine unless advised by your doctor.
- Do not take any new medicines without consulting your doctor.
- Do not give Axcel Serratiopeptidase-5mg Tablet to anyone else, even if they have the same symptoms or condition as you.

-Things to be careful of

- No interference with the capability of driving or operating heavy machinery.

Side effects

Like all medicines, Axcel Serratiopeptidase-5mg Tablet can cause side effects although not everybody gets them. You are advised to consult doctor/ pharmacist for further information. The following

AXCEL SERRATIOPEPTIDASE-5MG TABLET

Serratiopeptidase (5mg)

occasional side effects have been reported:

- Rash or redness.
- Diarrhea.
- Loss of appetite.
- Gastric discomfort.
- Nausea or vomiting.
- Bleeding from the nose or bloody mucus.
- Shock (life-threatening condition).
- Allergic reaction symptoms.
- Inflammation of the liver.
- Jaundice (yellow discoloration of skin and eyes).

You may report any side effects or adverse drug reactions directly to the National Centre for Adverse Drug Reaction Monitoring by calling Tel: 03-78835550, or visiting the website npra.moh.gov.my (Public Reporting Medicinal Problems / Side Effects / AEFI / Vaccine Safety).

Storage and disposal of Axcel Serratiopeptidase-5mg Tablet

-Storage

Keep Axcel Serratiopeptidase-5mg Tablet out of reach of children. Store below 25°C. Keep container well closed. Protect from humidity and light.

-Disposal

Do not use Axcel Serratiopeptidase-5mg Tablet after the expiry date which is stated on the carton.

Medicines should not be disposed of via wastewater or household waste. Ask your pharmacist how to dispose of medicines no longer required. These measures will help to protect the environment.

Product description

-What it looks like

Axcel Serratiopeptidase-5mg Tablet is light pink, 8mm round, biconvex and enteric-coated tablets.

-Ingredients

- Active Ingredient:
Serratiopeptidase

- Inactive Ingredients:

Microcrystalline Cellulose (Ph 102), Tablelose 80 (Lactose for DC), Sodium Starch Glycolate, Magnesium Stearate, Aerosil R972V, Opadry II (85) White, Acryl-EZE pink and Purified Water.

-MAL Number

MAL10033139X

Manufacturer and product registration holder

Kotra Pharma (M) Sdn Bhd
No. 1, 2 & 3, Jalan TTC 12,
Cheng Industrial Estate,
75250 Melaka, Malaysia.

Date of revision

25/01/2018

Serial number

NPRA(R4/1)170118/00014