## TERNOLOL FILM COATED TABLET

Atenolol (50 mg, 100 mg)

#### What is in this leaflet?

- 1. What Ternolol is used for
- 2. How Ternolol works
- 3. Before you use Ternolol
- 4. How to use Ternolol
- 5. While you are using Ternolol
- 6. Side effects
- 7. Storage and Disposal of Ternolol
- 8. Product description
- 9. Manufacturer and Product Registration Holder
- 10. Date of Revision

#### What Ternolol is used for

This medicine contains the active ingredient Atenolol.

#### Used to treat:

- hypertension (high blood pressure)
- angina pectoris (chest pain or discomfort)
- acute myocardial infarction (heart attack)
- cardiac arrhythmias (irregular heart rhythm).

## **How Ternolol works**

Atenolol belongs to a family of medicines called beta blockers.

It works by slowing down or decreasing the heart's need for blood and oxygen. This results in reduced workload for the heart.

#### Before you use Ternolol

## - When you must not use it

Do not take this medicine if:

- you are allergic to atendlol or any of the excipients
- you have cardiogenic shock( failure of pump action of the heart).
- you have heartblock
- you have overt cardiac failure (heart failure)

• you have sinus bradycardia (a regular but usually slow heartbeat) or reversible obstructive airways disease.

#### - Before you start to use it

#### Tell your doctor if:

- you have congestive heart failure
- you have poor cardiac reserve (the ability of your heart to work in strenuous conditions)
- you have impaired kidney/liver function.

#### - Taking other medicines

Talk to your doctor or pharmacist if you are taking any other medicines in any form including any that you buy or obtained without a prescription.

Some medicines interfere with Ternolol. Tell your doctor if you are taking:

- Clonidine (used to treat ADHD (Attention Deficit Hyperactivity Disorder)
- Reserpine (used for the control of high blood pressure)
- Phenylephrine (used as decongestant)

Other interactions not stated may occur. Talk to your doctor or pharmacist if you have any other concerns.

#### How to use Ternolol

## - How much to use

#### Adults

## Treatment of agina

One 50 mg tablet once a day or increased gradually to one 100 mg tablet daily.

#### Treatment of high blood pressure

25 mg to 50 mg tablet once a day or increased gradually to 50 mg to 100 mg tablet daily.

# Acute myocardial infarction (Heart attack) management

One 50 mg tablet 10 minutes after last intravenous does, followed by another 50 mg 12 hour later.

You may be then given a dose of 100 mg once a day or 50 mg twice a day for 6 to 9 days or until discharged from the hospital.

Treatment of patients with kidney problems

• your dose may be reduced

Swallow the tablet with a full glass of water.

#### - When to use it

Take your medicine at about the same time each day will have the best effect.

#### - How long to use it

You are advised to take it as long as it is given to you by your doctor.

## - If you forget to use it

Do not take a double dose to make up for the forgotten dose.

If you forget to take your dose, skip the missed dose. Take the next dose as normal.

#### - If you use too much (overdose)

Contact your doctor immediately or go to the Emergency Department of your nearest hospital, if you think you or anyone else may have taken too much of this medicine. Do this even if there are no signs of discomfort or poisoning. You may need urgent medical attention.

If you take too much Ternolol, you may feel drowsiness, coma, hallucinations and have a fast or irregular heart beat.

#### While you are using Ternolol

#### - Things you must do

Tell your doctor as soon as possible if:

- you become pregnant.
- you are going to start taking any new medicines

#### - Things you must not do

Do not give this medicine to anyone else, even if their symptoms seem similar to yours.

Do not take your medicine to treat any other complaints unless you are told by your doctor or pharmacist.

Do not stop taking the medicine or change the dosage on your own before asking your doctor.

- Things to be careful of
- Driving and using machines

#### Side effects

All medicines can cause unwanted side effects. These effects can be serious but most of the time, they are mild and temporary.

Tell your doctor immediately if you think you are experiencing any of the following possible unwanted side effects while taking the medicine.

## Common side effects

- Headache
- Fatigue and tiredness
- Dizziness
- Nausea
- Dry mouth
- Allergic reactions
- Diarrhoea
- Edema

Other unwanted side effects not listed may occur. Tell your doctor immediately.

You may report any side effects or adverse drug reactions directly to the National Centre for Adverse Drug Reaction Monitoring by calling Tel: 03-78835550, or visiting the website portal.bpfk.gov.my (Consumers→Reporting)

#### Storage and Disposal of Ternolol

#### - Storage

Keep out of reach of children. Store below 25°C. Protect from light and moisture.

#### - Disposal

No specific instructions. Please ask your pharmacist how to dispose unused medicines properly.

#### **Product description**

#### - What it looks like?

Round, orange film-coated tablet with shallow convex faces bevel-edged and with a heart-shaped protrusion.

### - Ingredients

- Active ingredient Ternolol 50 film coated tablet contains Atenolol 50 mg per tablet

*Ternolol 100 film coated tablet* contains Atenolol 100 mg per tablet

#### - Inactive ingredients

Hydroxypropyl methylcellulose Stearic acid Sodium lauryl sulphate Sunset yellow lake Titanium dioxide Microcrystalline cellulose Pregelatinised starch Propylene glycol Isopropyl alcohol Magnesium stearate

Talc Crospovidone

Lactose

- MAL number (s): Ternolol 50 film coated tablet MAL19940057A

Ternolol 100 film coated tablet MAL19940058A

#### Manufacturer

HOVID Berhad Lot 56442, 7<sup>1</sup>/2 Miles, Jalan Ipoh / Chemor, 31200 Chemor, Malaysia.

#### **Product Registration Holder**

HOVID Bhd. 121, Jalan Tunku Abdul Rahman, 30010 Ipoh, Malaysia.

#### Date of revision

25/03/2014