

# TRIVEDON MR<sup>®</sup>

Trimetazidine Dihydrochloride (35 mg)

## What is in this leaflet

- 1 What TRIVEDON MR is used for
- 2 How TRIVEDON MR works
- 3 Before you use TRIVEDON MR
- 4 How to use TRIVEDON MR
- 5 While you are using it
- 6 Side effects
- 7 Storage and Disposal of TRIVEDON MR
- 8 Product Description
- 9 Manufacturer and Product Registration Holder
10. Date of Revision

## What TRIVEDON MR is used for

This medicine is intended for use in adult patient, to treat angina pectoris (chest pain caused by heart disease).

It is used in combination with other medicines to treat vertigo (dizziness), tinnitus (ringing in the ears) and impaired vision (decline of visual acuity and visual field disturbances).

## How TRIVEDON MR works

It protects the heart cells from the effects of a reduced oxygen supply during an episode of angina.

## Before you use TRIVEDON MR

### When you must not use it

#### **Do not take Trivedon MR**

if you are allergic to trimetazidine or any of the other ingredients of this medicine.

### Before you start to use it

Talk to your doctor or pharmacist before taking Trivedon MR. This medicine is not a cure for chest pain and should not be used as an initial treatment for unstable angina or heart attack.

### Taking other medicines

Please tell your doctor or pharmacist if you are taking or have recently taken any other medicines, including medicines obtained without a prescription and herbal medicines.

## How to use TRIVEDON MR

### How much to use

Always take Trivedon MR exactly as your doctor has told you. You should check with your doctor or pharmacist if you are not sure. The recommended dose of Trivedon MR is one tablet to be taken in the morning and evening during meals.

### When to use it

Take Trivedon MR during mealtimes

### How long to use it

You should discuss with your doctor before you stop taking your tablets. If you have any further questions on the use of this medicine, ask your doctor or pharmacist.

### If you forget to use it

If you forget to take one or more doses, do not take a double dose to make up for a forgotten dose.

### If you use too much (overdose)

If you take too many tablets, please contact your doctor or the nearest hospital Emergency Department immediately.

## While you are using TRIVEDON MR

### Things you must do

In the event of an angina attack, tell your doctor. Your treatment should be re-evaluated.

### Things you must not do

Trivedon MR is not recommended in children aged below 18 years.

## Pregnancy and lactation

As a precautionary measure, it is preferable not to take Trivedon MR during pregnancy or breast feeding.

If you are pregnant or breast-feeding, think you may be pregnant or are planning to have a baby, ask your doctor or pharmacist for advice before taking this medicine.

## Things to be careful of

### **Driving and using machines**

This medicine may make you feel dizzy and drowsy that may affect your ability to drive or use machinery.

### **Side effects**

Like all medicines, Trivedon MR can cause side effects, although not everybody gets them.

The most commonly encountered side-effects are: discomfort in the stomach, nausea, vomiting, headache, dizziness. Inform your doctor if you encounter any side effects listed above or not as your doctor will decide whether you should continue taking this medication

**You may report any side effects or adverse drug reactions directly to the National Centre for adverse Drug Reaction monitoring by calling Tel: 03-78835550 or visiting the web site: [portal.bpfk.gov.my](http://portal.bpfk.gov.my) (Consumers → Reporting).**

## Storage and Disposal of TRIVEDON MR

### **Storage**

Keep out of the reach and sight of children.

Store below 25°C.

If your doctor decides to stop your treatment, return any tablets left over to the pharmacist. Only keep them if your doctor tells you to.

You should not take your medicine after the expiry date (MM YYYY), which is clearly marked on the blister and on the outside of the box. The expiry date refers to the last day of that month.

### **Disposal**

Medicines should not be disposed of via wastewater or household waste.

Ask your pharmacist how to dispose of medicines no longer required. These measures will help to protect the environment

# TRIVEDON MR<sup>®</sup>

Trimetazidine Dihydrochloride (35 mg)

---

## Product description

### - What it looks like

White to off white circular biconvex, film coated tablets plain on both sides with aperture on one face

### - Ingredients

The active ingredient is Trimetazidine Dihydrochloride

### -Inactive ingredients:

The other ingredients are Sodium Chloride, Povidone, Magnesium Stearate, Cellulose acetate (CA-398-10), Hypromellose (HPMC 6 cps), Ethanol, Methylene Chloride

**MAL NO:** MAL12020015A

## Manufacturer

Cipla Ltd.  
L-139 to L-146  
Verna industrial Estate,  
Verna Salcette, Goa India

## Product Registration Holder

CIPLA MALAYSIA SDN BHD  
Suite 1101, Amcorp Tower,  
Amcorp Trade Centre,  
18 Persiaran Barat,  
46 050 Petaling Jaya,  
Selangor, Malaysia.

## Date of revision

23/07/2015

## Serial Number

BPFK(R4/1)070515/00095