

# WINGESIC 500 TABLET

Paracetamol (500mg)

## What is in the leaflet

1. What Wingsic 500 Tablet is used for
2. How Wingsic 500 Tablet works
3. Before you use Wingsic 500 Tablet
4. How to take Wingsic 500 Tablet
5. While you are using it
6. Side Effects
7. Storage and Disposal of Wingsic 500 Tablet
8. Product Description
9. Manufacturer and Product Registration Holder
10. Date of revision

## What Wingsic 500 Tablet is used for

Wingsic 500 Tablet is used for the relief of fever and mild to moderate pain including headache, sinus pain, toothache and body aches associated with cold and flu.

## How Wingsic 500 Tablet works

Wingsic 500 Tablet contains Paracetamol which belongs to a group of medicines called analgesic and antipyretic that helps to ease pain and reduce fever. It is rapidly and completely absorbed following oral administration and will exert its effect between 30 minutes to 2 hours later. This effect could last for 3 to 4 hours.

## Before you use Wingsic 500 Tablet

### - When you must not take it

Do not take this product if you are allergic (hypersensitivity) to Paracetamol or any other ingredients in this product.

### - Before you start to use it

This medicine should be given with care to patients with impaired liver or kidney function and to patients taking other drugs that affect the liver.

### - Taking other medicines

You should always check with your doctor if your child is taking any of these medicines:

- Anticoagulant drugs (Warfarin)
- Metoclopramide (medication for stomach problem)
- Propantheline (medication to treat peptic ulcers)
- Antidepressants with anticholinergic properties and narcotic analgesics
- Chloramphenicol (antibiotic)
- Enzymes such as alcohol and anticonvulsant agents
- Probenecid (medication to treat gout)
- Cholestyramine (medication to reduce cholesterol)

## How to take Wingsic 500 Tablet

### - How much to take

Adults and children above 12 years: Take 1 to 2 tablets every 4 to 6 hours. (Do not exceed 8 tablets per day)

### - When to take it

May be taken with or without food.

### - How long to take it

Take as directed by your doctor or pharmacist.

### - If you forget to take it

If you forget to give Wingsic 500 Tablet to your child, give it as soon as you remember. Never double up on the next dose to make up for the one missed.

### - If you take too much (overdose)

Symptoms include vomiting, abdominal pain, hypotension and sweating. In case of overdose, prompt specific treatment is essential.

Call National Poison Centre (1800 888 099) or your doctor or go to the accident and emergency centre at your nearest hospital immediately.

## While you are using Wingsic 500 Tablet

### - Things you must do

Do not exceed the recommended dosage and continual use should not exceed 5 days at a time (10 days for adult), without physician guidance.

If symptoms persist, please consult a doctor.

### - Things you must not do

Do not take this product to treat any other conditions unless it has been prescribed by your doctor.

### - Things to be careful of

This preparation contains Paracetamol. Do not take any other Paracetamol containing medicines at the same time.

## Side Effects

Allergy alert: Paracetamol may cause severe skin reactions. Symptoms may include skin reddening, blisters or rash. These could be signs of a serious condition. If these reactions occur, stop use and seek medical assistance right away.

You may report any side effects or adverse drug reactions directly to the National Centre for Adverse Drug Reaction Monitoring by calling Tel: 03-7883 5549, or visiting the website [npra.moh.gov.my](http://npra.moh.gov.my) (Public → Reporting Side Effects to Medicines (ConSERF or Vaccines (AEFI))).

## Storage and Disposal of Wingsic 500 Tablet

### - Storage

- Store below 30°C.
- Protect from light.
- Keep container tightly closed.
- Keep out of reach of children.

### - Disposal

Do not take this after the expiry date. An expiry date is printed on the label of the medicine.

## Product Description

### - How it looks like

Wingsic 500 Tablet is a pink coloured, biconvex and round shaped uncoated tablet.

### Ingredients

Active Ingredients

Each tablet contains Paracetamol 500mg

Inactive Ingredients

Gelatin, Magnesium Stearate, Corn Starch, Carmoisine lake, Sodium Starch Glycolate, Vinyl Pyrrolidone (Plasdone S630), Aerosil 200

### - MAL Number:

MAL08051671XZ

## Manufacturer and Product Registration Holder

WINWA MEDICAL SDN. BHD.  
1952, Lorong I.K.S. Bukit Minyak 3,  
Taman I.K.S. Bukit Minyak,  
14000 Bukit Mertajam,  
Pulau Pinang, Malaysia.

## Date of revision

14/5/2018

Serial Number:

NPRA(R4/1)140518/00145

X.LWG-PIL.01