

# REDAMOL 120 SYRUP

Paracetamol (120mg/5ml)

## What is in this leaflet

1. What Redamol 120 Syrup is used for
2. How Redamol 120 Syrup works
3. Before you use Redamol 120 Syrup
4. How to take Redamol 120 Syrup
5. While you are using Redamol 120 Syrup
6. Side effects
7. Storage and disposal of Redamol 120 Syrup
8. Product description
9. Manufacturer and Product registration holder
10. Date of revision

## What Redamol 120 Syrup is used for

Redamol 120 Syrup is used for the relief of fever, headache and symptoms of cold and flu, toothache, discomfort of teething and fever after vaccination.

## How Redamol 120 Syrup works

It helps to relieve pain by blocking the pain messengers called prostaglandins in the central nervous system.

## Before you use Redamol 120 Syrup

### -When you must not use it:

- If you are allergic to paracetamol or aspirin or any of the ingredients in this medicine. Allergy alert: Paracetamol may cause severe skin reactions. Symptoms may include skin reddening, blisters or rash. These could be signs of a serious condition. If these reactions occur, stop use and seek medical assistance right away.
- If you have alcohol addiction.
- If you have liver disease.
- If you have viral infection.
- If you have kidney function impairment.
- If you have heart and blood vessels disease.

### -Before you start to use it:

Please tell your doctor if you are suffering or have ever suffered from any of the following conditions or illnesses:

- Pregnancy or breastfeeding.

### -Taking other medicines:

Please tell your doctor or pharmacist if you are taking or have recently taken any other medicines, including medicines obtained without a prescription. Your doctor may need to change the dose of the other medications or take other precautions. In some cases you may have

to stop taking one of the medicines. This applies especially to the medicines listed below taken at the same time with Redamol 120 Syrup:

- Oral anticoagulant agents (used to prevent blood clots).
- Paracetamol (Do not take other paracetamol-containing medicine at the same time).

## How to take Redamol 120 Syrup

### -How much to use:

Children: 1 to 6 years-

Age	Dose
1 – 3 years	6ml – 10ml
4 – 6 years	10ml – 15ml

The above table is based on the average weight-age grid of children. If the body weight of the child known, it's always more accurate to use the weight to calculate the correct dose. Please refer below.

Using the body weight of the child, calculate the dose of Redamol 120 Syrup the child needs. Calculation is based on recommended dose of 15mg per body weight in kg (BW).

Table below provides the dose of Redamol 120 Syrup to be given, based on body weight:

Child's weight		Redamol 120 Syrup
In kg	In lb	
10	22	6 ml
15	33	9 ml
20	44	13 ml
30	66	19 ml

Note: 1kg = 2.2lb

### When to use it:

Give in 6 hours intervals, not more than 4 times a day. Take the syrup exactly as your doctor has told you to. The dose will be on the pharmacist's label. Check the label carefully. It is stated how many ml you need to take and how often. If you are not sure, ask your doctor or pharmacist.

### -How long to use it:

Keep taking the syrup for as long as you have been told unless you have any problems. In that case, check with your doctor.

### -If you forget to use it:

If you forget to take a dose, do not worry. Take it as soon as you remember then carry on as before. If you do not take your dose on one day, take your normal dose on the next day. Do not take a double dose to make up for forgotten individual doses.

### -If you use too much (overdose):

If you accidentally take too many dose, contact your doctor, pharmacist, or your nearest hospital emergency department immediately.

Symptoms of Redamol 120 Syrup overdose include diarrhoea, vomiting or nausea, loss of appetite, stomach cramps or pain and unusual increase sweating.

## While you are using Redamol 120 Syrup

### -Things you must do

- Take your medicine as advised from your doctor.
- Tell all the doctors, dentists and pharmacists treating you that you are taking Redamol 120 Syrup.
- Tell your doctor immediately if you are pregnant while taking this medication.

### -Things you must not do

- This medicine has been prescribed for you. Never give it to someone else. It may not be the right medicine for them even if their symptoms seem to be the same as yours.

### -Things to be careful of

- Special care is recommended during pregnancy. Take the medicine during pregnancy if the benefits against possible effects on the child had been assessed by the doctor.

## Side effects

Like all medicines, Redamol 120 Syrup can cause side effects, although not everybody gets them. Visit your doctor or pharmacist immediately if you experience any side effects after taking this medicine:

- Bloody or cloudy urine.
- Difficult or painful urination.
- Frequent urge to urinate.
- Sudden decrease in amount of urine.

# REDAMOL 120 SYRUP

Paracetamol (120mg/5ml)

---

- Sterile pyuria (presence of white cells in urine).
- Skin rash, hives or itching.
- Unexplained sore throat and fever.
- Unusual tiredness or weakness.
- Yellowing of eyes or skin.
- Cutaneous hypersensitivity reactions (including skin rashes, swelling under the skin, Steven Johnson Syndrome/Toxic Epidermal Necrolysis).

If you notice undesirable effects which have not been mentioned in this leaflet, please consult your doctor/pharmacist for further information.

You may report any side effects or adverse drug reactions directly to the National Centre for Adverse Drug Reaction Monitoring by calling Tel:03-78835550 or visiting the website [npra.moh.gov.my](http://npra.moh.gov.my) (*Public → Reporting Side effects to Medicines (ConSERF) or Vaccines (AEFI)*).

## Storage and disposal of Redamol 120 Syrup

### -Storage:

Store in a dry place below 30°C. Protect from light.

### -Disposal:

Medicines should not be disposed of via wastewater or household waste. Ask your pharmacist how to dispose the medicines you no longer required. These measures will help to protect the environment.

## Product description

### -What it looks like:

Red colour, fruity flavored syrup.

### -Ingredients:

-Active Ingredient  
Paracetamol.

### -Inactive Ingredient

Colour ponceau, strawberry flavour, raspberry flavour, propyl paraben, sodium saccharin, glycerine, refined sugar, citric acid anhydrous, sodium benzoate, polyethylene glycol, methyl paraben and purified water.

### -MAL number:

MAL19910340XZ

## Manufacturer & Product Registration Holder

Royce Pharma Manufacturing Sdn. Bhd.  
PT 1663, Nilai Industrial Estate,  
71800 Nilai, Negeri Sembilan,  
Malaysia.

## Date of revision

03/05/2018

## Serial Number

NPRA(R4/1)100418/00103