

TARIVID OTIC SOLUTION

Ofloxacin (3mg/mL)

What is in this leaflet

1. What Tarivid Otic Solution is used for
2. How Tarivid Otic Solution works
3. Before you use Tarivid Otic Solution
4. How to use Tarivid Otic Solution
5. While you are using it
6. Side effects
7. Storage and Disposal of Tarivid Otic Solution
8. Product Description
9. Manufacturer and Product Registration Holder
10. Date of revision

What Tarivid Otic Solution is used for
Tarivid Otic Solution is a topical antibacterial agent used to treat external and middle ear infections.

How Tarivid Otic Solution works

Tarivid Otic Solution interferes with bacterial enzymes to prevent bacterial growth, thereby killing the susceptible bacteria.

Before you use Tarivid Otic Solution

- When you must not use it

You should not use Tarivid Otic Solution if you have a history of hypersensitivity to ofloxacin.

Pregnancy and lactation

Ask your doctor or pharmacist for advice before taking any medicine.

- Before you start to use it

- Use Tarivid Otic Solution at a temperature as similar to that of the body temperature as possible.

- Taking other medicines

Tell your doctor if you are taking any other medicines, including any that you buy from a pharmacy, supermarket or health food shop without a prescription.

How to use Tarivid Otic Solution

- How much to use

For adults, 6-10 drops of Tarivid Otic Solution should be applied into the infected ear 2 times a day.

For children, dosage may be decreased.

- When to use it

Use as directed by your doctor or pharmacist.

- How long to use it

Generally, duration of treatment with this medicine should be 4 weeks.

- If you forget to use it

Take the missed dose as soon as you remember.

Do not double the dose

- If you use too much

Contact your doctor immediately.

While you are using Tarivid Otic Solution

- Things you must do

The eardrop should be allowed to maintain in the ear for 10 minutes.

You should complete the full course of treatment.

- Things you must not do

Do not stop using the medicine unless advised by your doctor.

- Things to be careful of

Route of administration: Use only for instillation.

If the solution is cold, dizziness may occur.

Thus, use the medicine at a temperature as similar to that of the body temperature as possible.

Side Effects

Worsening of muscles weakness

Tarivid Otic Solution is generally well tolerated. Earache may occur infrequently.

You may report any side effects or adverse drug reactions directly to the National Centre for Adverse Drug Reaction Monitoring by calling Tel: 03-78835550, or visiting the website npra.moh.gov.my (Public → Reporting Medicinal Problems / Side Effects / AEFI / Vaccine Safety).

Storage and Disposal of Tarivid Otic Solution

- Storage

Keep out of the reach and sight of children.

Store below 30°C.

Stability after opening:

This product exhibits good stability for 4 months at room temperature (25°C) when it is stored in an attached light protective vinyl plastic bag after opening.

- Disposal

Medicines should not be disposed of via wastewater or household waste. Ask your pharmacist how to dispose of medicines no longer required. These measures will help to protect the environment.

Product description

- What it looks like

Pale yellow to light yellow and clear solution.

- Ingredients

- Active ingredient(s)

Ofloxacin

- Inactive ingredients

Benzalkonium chloride,

sodium chloride

- MAL number:

MAL20122266ACZ

Manufacturer

Santen Pharmaceutical Co. Ltd. Noto plant, 2-14, Shikinami, Hodatsushimizu-cho, Hakui-gun, Ishikawa, Japan.

Product Registration Holder

First Pharmaceutical Sdn Bhd
No. 20, Jalan SS 19/5
47500 Subang Jaya
Selangor, Malaysia

Date of Revision

22/2/2018

Serial Number

NPRA(R4/1)141217/00358