LEZPAIN FILM COATED TABLET

Mebeverine Hydrochloride (135mg)

What is in this leaflet

- 1. What LEZPAIN FILM COATED TABLET is used for
- 2. How LEZPAIN FILM COATED TABLET works
- 3. Before you use LEZPAIN FILM COATED TABLET
- 4. How to use LEZPAIN FILM COATED TABLET
- 5. While you are using it
- Side effects
- 7. Storage and disposal LEZPAIN FILM COATED TABLET
- 8. Product description
- 9. Manufacturer and Product Registration Holder
- 10. Date of revision

What LEZPAIN FILM COATED TABLET is used for

LEZPAIN FILM COATED TABLET is used for the symptomatic treatment of abdominal pain and cramps, bowel disturbances and intestinal discomfort related to irritable bowel syndrome.

How LEZPAIN FILM COATED TABLET works

Mebeverine is an antispasmodic, with a direct action on the smooth muscle of the gastrointestinal tract, relieving spasm without affecting normal gut movement.

Before you use LEZPAIN FILM COATED TABLET

-When you must not use it

Do not take LEZPAIN FILM COATED TABLET if you have known hypersensitivity to mebeverine hydrochloride or have shown hypersensitivity to any component of this product.

-Before you start to use it

Tell your doctor or pharmacist if you are allergic to any medicines or if you have any other medical conditions.

Pregnancy and lactation

Please consult your doctor before using LEZPAIN FILM COATED TABLET if you are pregnant, trying to get pregnant or think you may be pregnant and if you are breast-feeding. LEZPAIN FILM COATED TABLET can be used during breast-feeding as mebeverine is not found in breastmilk. Ask your doctor or pharmacist for advice before taking any medicine.

-Taking other medicines

Tell your doctor or pharmacist if you are taking any other medications, including those without a prescription.

How to take LEZPAIN FILM COATED TABLET

-How much to use

Adult and children over 10 years: 135mg 3 times daily

-When to use it

To be taken 20 minutes before meals

-How long to use it

Consult your doctor on the duration of consumption.

-If you forget to use it

If you miss a dose of LEZPAIN FILM COATED TABLET and you are taking it regularly, take it as soon as possible. If several hours have passed or if it is approaching time for the next dose, do not double the dose to make up for the forgotten dose, unless advised by your health care provider. Do not take 2 doses at once.

-If you use too much (overdose)

Seek for medical attention from your doctor immediately. Theoretically, it may be predicted that central nervous system (CNS) excitability will occur in cases of overdosage.

While you are using it

-Things you must do

Follow your doctor's instruction when you are taking LEZPAIN FILM COATED TABLET.

-Things you must not do

Do not start taking this medicine or change the dosage without first checking with your doctor. Do not give this medicine to anyone else, even if their symptoms seem similar to yours.

-Things to be careful of

Driving and using machines

When taken as prescribed, mebeverine has no effect on psychomotor skills associated with driving and operating machinery.

Side effects

Like all medicines, LEZPAIN FILM COATED TABLET can cause side effects, although not everybody gets them.

Visit your doctor or pharmacist immediately if you experience any side effects after taking this medicine.

You may report any side effects or adverse drug reactions directly to the National Centre for Adverse Drug Reaction Monitoring by calling Tel: 03-78835550, or visiting the website portal.bpfk.gov.my

 $(Consumers \rightarrow Reporting)$

Storage and Disposal of LEZPAIN FILM COATED TABLET

-Storage

Store below 30°C, protect from light & moisture.

-Disposal

Dispose when you stop taking the tablets or the tablets have passed their expiry date. Ask your doctor or pharmacist what to do with any tablets that are left over.

Product description

-What it looks like

A white color film coated round tablet.

-Ingredients

-Active Ingredient-

Mebeverine Hydrochloride

-Inactive Ingredients-

Povidone

Silicon Dioxide

Magnesium Stearate

Lactose

Cellulose

Polyethylene Glycol

Dimethicone

Titanium Dioxide

Isopropyl Alcohol

<u>-MAL number</u> MAL06011697A

Manufacturer

Y.S.P Industries (M) Sdn. Bhd. Lot 3,5 & 7, Jalan P/7, Section 13, Kawasan Perindustrian Bandar Baru Bangi, 43000 Kajang, Selangor Darul Ehsan, Malaysia.

Product Registration Holder

Y.S.P Industries (M) Sdn. Bhd. Lot 3,5 & 7, Jalan P/7, Section 13, Kawasan Perindustrian Bandar Baru Bangi, 43000 Kajang, Selangor Darul Ehsan, Malaysia

Date of revision

04/02/2016

Serial Number

BPFK(R4/1)030216/00026