

LASTUSS COUGH SYRUP

Detromethorphan Hydrobromide (30mg/5ml)

What is in this leaflet

1. What LASTUSS is used for
2. How LASTUSS works
3. Before you use LASTUSS
4. How to use LASTUSS
5. While you are using it
6. Side effects
7. Storage and Disposal of cough suspension
8. Product Description
9. Manufacturer and Product Registration Holder
10. Date of revision

What is used for LASTUSS

For dry and unproductive cough

How LASTUSS works

Dextromethorphan is an antitussive to help stop persistent coughing

Before you use LASTUSS

- When you must not use it

- Hypersensitivity to Dextromethorphan
- Persistent or long term cough accompanied by excessive secretions
- For those receiving monoamine-oxidase inhibitors
- Liver function impairment

- Before you start use it

Talk to your doctor or pharmacist.

- If you are pregnant or breast-feeding
- If you are taking alcohol.
- If you have an intolerance to some sugars.
- If you are taking any other medicines.
- Dextromethorphan is not suitable to be used in children below 2 years old.
- Dextromethorphan should be used cautiously and to follow the advice of physician or pharmacist for children between the ages of 2 to 6 years

- Taking other medicines

Alcohol and medicines causing central nervous system depression should be avoided when taking dextromethorphan

How to use LASTUSS

- How much to use

Adult: 2 teaspoonfuls
 Children (6-12 years): 1 teaspoonful
 Children (2-6 years): half teaspoonful
 or as directed by the physician

- When to use it

To be taken twice a day or use as directed by your doctor or pharmacist.

- How long to use it

Continue taking LASTUSS for as long as your doctor recommends.

- If you forget to use it

Take the missed dose as soon as you remember. If it is almost time for the next dose, ignore the missed dose and wait to take the next dose at regular intervals. Do not take double doses to replace the missed doses.

- If you use too much (overdose)

Contact your doctor immediately or go to the Emergency Department of your nearest hospital.
 Manifestations following over- dosage of dextromethorphan have included nausea, vomiting, drowsiness, dizziness, blurred vision, and involuntary eye movement, lack of muscle coordination, short breaths, unable to urinate, insensibility, substance- induced psychosis and coma.

While you are using it

- Things you must do

Take your medicine exactly as your doctor has told you.

Tell all the doctors, dentists and pharmacists treating you that you are taking LASTUSS.

Tell your doctor immediately if you become pregnant while taking this medication.

- Things you must not do

Do not stop taking the medicine unless advised by your doctor.

Do not take any new medicines without consulting your doctor or pharmacist.

Do not give LASTUSS to anyone else, even if they have the same symptoms or condition as you.

- Things to be careful of

Shake thoroughly before use

Side effects

Occasional nausea, drowsiness and dizziness

You may report any side effects or adverse drug reactions directly to the National Centre for Adverse Drug Reaction Monitoring by calling Tel: 03-78835549, or visiting the website npra.moh.gov.my (Public Reporting Side effects to Medicines (ConSERF) or Vaccines (AEFI)).

Storage and Disposal of LASTUSS

- Storage

Keep in a cool, dark place below 30 °C.

- Disposal

Do not use your medicine after the date shown as an expiry date on the packaging.

Medicines should not be disposed of via wastewater or household waste. Ask your pharmacist how to dispose of medicines no longer required. These measures will help to protect the environment.

LASTUSS COUGH SYRUP

Detromethorphan Hydrobromide (30mg/5ml)

Product Description

- What it looks like

Orange colored suspension, pleasantly flavored, having sweet taste

- Ingredients

- Active ingredient

Dextromethorphan

- Inactive ingredients

Colloidal Silicon Dioxide, Butylated Hydroxy Toluene , Butylated Hydroxyanisole, Anhydrous Citric Acid, Propylene Glycol , Orange Sweet, Purified Water , Sucrose , Sorbitol Solution, Methyl Paraben, Propyl Paraben, Colour Sunset Yellow FCF, Polysorbate 80 , Xanthan Gum.

- MAL number:

MAL 19972991AZ

Manufacturer

FDC Limited
B-8, MIDC Industrial area,
Waluj, Aurangabad – 431 136,
Maharashtra

Product Registration Holder

Unimed Sdn Bhd
53, Jalan Tembaga SD 5/2B,
Bandar Sri Damansara,
52200, Kuala Lumpur, Malaysia.

Date of revision

30/5/2018

Serial number:

NPRA(R4/1)150518/00147