EUSOL I AND II

Chlorinated lime (2.5% w/v)

What is in this leaflet

- 1. What Eusol is used for
- 2. How Eusol works
- 3. Before you use Eusol
- 4. How to use Eusol
- 5. While you are using Eusol
- 6. Side effects
- 7. Storage and Disposal of Eusol
- 8. Product description
- Manufacturer and Product Registration holder

10.Date of revision

What Eusol is used for

It is used for use as an antiseptic wound dressing and cleansing agent.

How Eusol works

Chlorinated lime on exposure to air becomes moist and gradually decomposes, carbon dioxide being absorbed and chlorine evolved. It is decomposed by hydrochloric acid with evolution of available chlorine. It is not used internally.

Before you use Eusol

- When you must not use it
- allergic to the active and non-active ingredients of the product.
- antimicrobial activity of chlorine is reduced by the presence of organic material.

- Before you start to use it

It is locally applied, chlorinated lime may dissolve blood clots and cause bleeding. For external use only. It has a bleaching action.

- Taking other medicines

The medicine is for topical application and will usually have minimal effect on other medication. However, please inform the doctor or pharmacist if taking other medicines.

How to use Eusol

- How much to use

Used externally. Add EUSOL I into equal amount of EUSOL II before using as an antiseptic wound dressing. The mixture should be used within 2 weeks of its preparation.

- When to use it

Please follow the advice of the doctor or pharmacist.

- How long to use it

You should check with your doctor or pharmacist.

- If you forget to use it

If you forget to use this medicine, use it as soon as you remember. Then continue using it at your usual time.

- If you use too much (overdose)

If you have used too much Eusol, contact your doctor, an emergency department or a poison center immediately.

While you are using Eusol

- Things you must do

Inform your doctor of all the medications you are taking.

- Things you must not do

Do not stop using medication although you feel better after a few days, until the prescribed course is finished.

- Things to be careful of

If you are pregnant, become pregnant, or breastfeeding, please inform the doctor or pharmacist.

Side effects

Chlorinated lime may dissolve blood clots and cause bleeding.

You may report any side effects or adverse drug reactions directly to the National Centre for Adverse Drug Reaction Monitoring by calling Tel: 03-78835550, or visiting the website npra.moh.gov.my (Public → Reporting Medicinal Problems / Side Effects / AEFI / Vaccine Safety).

Storage and Disposal of Eusol

- Storage

Store in a dry place below 30°C. Protect from light. Keep out of reach of children.

- Disposal

Ask your pharmacist how to dispose of medicines no longer required.

Product description

- What it looks like

This product is packed in 2 separate bottles:

Eusol I: chlorinated lime

Eusol II: excipients of sodium acetate and acetic acid glacial.

- Ingredients:
- Active ingredients: chlorinated lime
- Inactive ingredient: sodium acetate, acetic acid glacial and deionised water
- <u>MAL number</u>: MAL07101360X

Manufacturer and Product Registration Holder

Prime Pharmaceutical Sdn. Bhd. 1505, Lorong Perusahaan Utama 1 Taman Perindustrian Bukit Tengah 14000 Bukit Mertajam, Penang, Malaysia.

Date of revision

04/12/2017

Serial number

NPRA(R4/1)301117/00332