

# **BRONPHYLLIN SYRUP**

Each 5ml contains Theophylline Anhydrous  
25mg

## **What is in this leaflet:**

1. What Bronphyllin Syrup is used for
2. How Bronphyllin Syrup works
3. Before you use Bronphyllin Syrup
4. How to take Bronphyllin Syrup
5. While you are using Bronphyllin Syrup
6. Side effects
7. After using Bronphyllin Syrup
8. Product Description
9. Manufacturer

## **What Bronphyllin Syrup is used for**

It is used for the relief of reversible bronchospasm which may occur in association with asthma, bronchitis or emphysema.

## **How Bronphyllin Syrup Works**

The active ingredient in the medicine is 'Theophylline Anhydrous'. Theophylline Anhydrous acts mainly on tissue linings in the lung. The medication relaxes the lung smooth muscles thereby improving the ease to breathe in patients with reversible obstructive airway disease. It also suppresses the airway from being oversensitive to allergens.

## **Before you use Bronphyllin Syrup**

### **When you must not take it**

- If you are allergic to Theophylline Anhydrous or any other ingredients in the product

### **Before you start to take it**

You should check with your doctor:-

- If you are or think you may be pregnant
- If you are planning to become pregnant
- If you are breastfeeding
- If you have irregular heart rhythm
- If you have active peptic ulcer disease
- If you have history of seizure disorder
- If you have congestive heart failure
- If you have kidney disease
- If you have liver disease
- If you have thyroid disorder
- If you are a smoker

## **Taking Other Medicines**

There are a few medicines which may not mix with Bronphyllin Syrup. It is important to tell your doctor or pharmacist about all the medicines that you are taking, including those obtained without doctor's prescription. You should talk to your doctor or pharmacist if you are on the following:-

- Antibiotics
- Ulcer healing medications
- Beta blockers
- Antihypertensives
- Oral contraceptives
- Antiepileptic drugs
- Lithium and other tranquilizers

## **How to take Bronphyllin Syrup**

### **How much to Take**

You should take Bronphyllin Syrup exactly as advised by your doctor or pharmacist. The amount you take each day will depend on your condition. Do not take more or less than what your doctor prescribes.

The usual dosage for adults is 25mg every 6 hours.

For children over 2 years, the recommended dosage is 1ml/kg body weight every 6 hours, to maximum of 25ml per dose.

It is not recommended to be used for children under 2 years of age except on the advice of a physician.

### **When to take**

Bronphyllin Syrup is to be taken after meal.

### **How long to take it**

It is important to take Bronphyllin Syrup as long as your doctor prescribes.

### **If you forget to take it**

Do not take extra dose to make up for a missed dose. Just take your next dose at usual the time.

## **If you take too much (overdose)**

If you take too much of the medication by mistake, contact your doctor immediately or seek medical attention.

## **While you are using Bronphyllin Syrup**

### **Things you must do**

Take Bronphyllin Syrup exactly the way as prescribed.

### **Things you must not do**

Do not share your medicines with others even if they have the same diagnosis as you

### **Things to be careful of**

If you experience any side effects, consult your doctor or pharmacist for advice

## **Side Effects**

The common side effects are nausea, vomiting, tremor, increased heart rate, palpitations, headache, insomnia. These symptoms are normally short-lived.

Careful monitoring by physician is required to avoid serious side effects.

Please refer to your doctor or pharmacist if you experience any other symptoms while taking this medication.

## **After using Bronphyllin Syrup**

### **Storage**

Store at temperature 25°C. Protect from light.

### **Disposal**

If you have any unwanted medicines, do not put them in waste water or household rubbish. Ask your pharmacist how to dispose of the medicine.

## **Product Description**

### **What it looks like**

A clear, orange coloured syrup with orange flavor

## **Date of Revision**

4 September 2011

## **Market Authorization Holder**

Sunward Pharmaceutical Sdn. Bhd.

***Ingredients***

Each 5ml contains Theophylline Anhydrous  
25mg as active ingredient.

Other non-active ingredients are:

Sunset yellow, sweet orange flavor, liquid sorbitol 70%, Syrup Simplex, citric acid, propylene glycol, sodium benzoate, sodium saccharin

***Product Registration Number***

MAL20020814A

---

**Manufacturer**

Sunward Pharmaceutical Sdn. Bhd.  
9&11, Jalan Kempas 4,  
Taman Perindustrian Tampoi Indah,  
81200 Johor Bahru, Johor, Malaysia.

**Date of Revision**  
4 September 2011

**Market Authorization Holder**  
Sunward Pharmaceutical Sdn. Bhd.

# **BRONPHYLLIN SYRUP**

Setiap 5ml mengandungi Theophylline anhydrous 25mg

## **Kandungan**

1. Kegunaan Bronphyllin Syrup.
2. Cara Bronphyllin Syrup berfungsi.
3. Perhatian sebelum mengambil Bronphyllin Syrup.
4. Cara pengambilan Bronphyllin Syrup.
5. Semasa pengambilan Bronphyllin Syrup.
6. Kesan sampingan.
7. Selepas mengambil Bronphyllin Syrup.
8. Gambaran produk.
9. Pengilang.

## **Kegunaan Bronphyllin Syrup**

Rawatan spasma bronkus berbalik yang berlaku akibat daripada asma, keradangan bronkus atau emfisema.

## **Cara Bronphyllin Syrup Berfungsi**

Theophylline anhydrous sebagai bahan aktif bertindak pada unjuran tisu di dalam peparu yang akan merehatkan otot-otot licin peparu. Tindakan ini memudahkan pernafasan bagi pesakit yang menghidapi penyakit salur pernafasan obstruktif berbalik. Mengurangkan kesensitifan salur pernafasan terhadap allergen.

## **Perhatian Sebelum Mengambil Bronphyllin Syrup**

### **Jangan ambil jika**

- Anda alergi kepada Theophylline anhydrous atau bahan-bahan lain yang terkandung di dalam produk ini.

### **Sebelum mengambil**

Dapatkan nasihat doktor jika anda:-

- Mengandung/ Disyaki mengandung.
- Merancang untuk mengandung.
- Menyusukan bayi.
- Mengalami degupan jantung tak sekata.
- Menghidapi penyakit ulser peptic.
- Pernah mengalami sawan.
- Menghidapi kegagalan jantung kongestif.
- Menghidapi penyakit buah pinggang.
- Menghidapi masalah hati.
- Menghidapi penyakit tiroid.
- Merokok.

### **Date of Revision**

4 September 2011

### **Mengambil ubat lain**

Terdapat beberapa jenis ubatan yang tidak digalakkkan diambil bersama dengan Bronphyllin Syrup. Anda harus memaklumkan doktor dan ahli farmasi anda mengenai ubatan yang sedang diambil termasuk ubatan yang dibeli di farmasi tanpa preskripsi doktor. Maklumkan doktor dan ahli farmasi jika anda sedang mengambil ubatan berikut:-

- Antibiotik.
- Ubat rawatan ulser.
- Penyekat beta.
- Antihipertensi
- Pil perancang
- Ubat antiepileptik
- Lithium dan ubat-ubat penenang yang lain.

## **Cara Mengambil Bronphyllin Syrup**

### **Arahan pengambilan**

Ikut arahan doktor atau ahli farmasi anda. Jangan ambil lebih atau kurang daripada aman yang disyorkan.

Dos biasa untuk dewasa adalah 25mg setiap 6 jam.

Kanak-kanak melebihi 2 tahun, dos saranan adalah 1ml/kg berat badan setiap 6 jam. Maksima adalah 25ml per dos.

Ubat ini tidak disarankan bagi kegunaan untuk kanak-kanak dibawah umur 2 tahun kecuali dengan nasihat doktor.

### **Waktu pengambilan**

Bronphyllin Syrup perlu diambil selepas makan.

### **Jangka masa pengambilan**

Ikut arahan yang telah ditetapkan oleh doktor atau ahli farmasi anda.

### **Jika terlupa mengambil**

Jangan gandakan dos anda. Ambil dos seperti biasa bila tiba waktu untuk dos seterusnya.

### **Kandungan**

### **Jika terlebih dos**

Hubungi doktor anda atau dapatkan bantuan kecemasan

## **Semasa Pengambilan Bronphyllin Syrup**

### **Perkara yang perlu dilakukan**

Ambil Bronphyllin Syrup mengikut arahan doktor.

### **Larangan**

Jangan kongsi ubatan anda dengan orang lain walaupun mereka mengalami masalah yang sama.

### **Perhatian**

Rujuk doktor atau ahli farmasi anda jika anda mengalami mana-mana kesan sampingan yang dinyatakan.

## **Kesan Sampingan**

Kesan sampingan biasa adalah loya, muntah, mengigil, peningkatan kadar jantung, jantung berdebar, sakit kepala dan sukar tidur. Kesan-kesan ini hanyalah untuk sementara waktu sahaja.

Rujuk doktor atau ahli farmasi anda jika anda mengalami kesan-kesan lain semasa pengambilan ubat ini.

## **Selepas Mengambil Bronphyllin Syrup**

### **Penyimpanan**

Simpan pada suhu 25°C. Lindungi daripada cahaya.

### **Pelupusan**

Jangan buang ubatan yang berlebihan secara sebarang. Dapatkan nasihat ahli farmasi anda mengenai cara-cara pelupusan ubatan.

## **Gambaran Produk**

### **Rupa produk**

Sirap berwarna dan berperasa oren.

### **Kandungan**

Setiap 5ml mengandungi Theophylline anhydrous 25mg sebagai bahan aktif.

## **Market Authorization Holder**

Sunward Pharmaceutical Sdn. Bhd.

Setiap 5ml mengandungi Theophylline anhydrous 25mg sebagai bahan aktif.

Lain-lain kandungan bahan tidak aktif:

Sunset yellow, perasa oren manis, cecair sorbitol 70%, Sirap Simplex, Asid sitrik, propylene glycol, sodium benzoate, sodium saccharin.

**Nombor Pendaftaran Produk**

MAL20020814A

---

**Pengilang**

Sunward Pharmaceutical Sdn. Bhd.

9&11, Jalan Kempas 4,  
Taman Perindustrian Tampoi Indah,  
81200 Johor Bahru, Johor, Malaysia.

**Date of Revision**  
4 September 2011

**Market Authorization Holder**  
Sunward Pharmaceutical Sdn. Bhd.