

COVERSYL PLUS 4 mg/1.25 mg, tablet

Perindopril tert-butylamine 4mg/Indapamide 1.25mg

What is in this leaflet

1. What COVERSYL PLUS 4mg/1.25mg is used for
2. How COVERSYL PLUS 4mg/1.25mg works
3. Before you use COVERSYL PLUS 4mg/1.25mg
4. How to take COVERSYL PLUS 4mg/1.25mg
5. While you are using COVERSYL PLUS 4mg/1.25mg
6. Possible side effects
7. After using COVERSYL PLUS 4mg/1.25mg
8. Product description

What is COVERSYL PLUS 4mg/1.25mg is and what it is used for

COVERSYL PLUS 4mg/1.25mg is a combination of two active ingredients, perindopril and indapamide. It is an anti-hypertensive and is used in the treatment of high blood pressure (hypertension).

How COVERSYL PLUS 4mg/1.25mg works

Perindopril belongs to a class of medicines called ACE inhibitors. These work by widening the blood vessels, which makes it easier for your heart to pump blood through them. Indapamide is a diuretic. Diuretics increase the amount of urine produced by the kidneys. However, indapamide is different from other diuretics, as it only causes a slight increase in the amount of urine produced. Each of the active ingredients reduces blood pressure and they work together to control your blood pressure.

Before you use COVERSYL PLUS 4 mg/1.25 mg, tablet

Do not take COVERSYL PLUS 4 mg/1.25 mg, tablet:

- if you are allergic (hypersensitive) to perindopril or to any other angiotensin-converting enzyme inhibitor, to indapamide or to any other sulphonamide or to any of the other ingredients of COVERSYL PLUS 4mg/1.25mg,
- if you have already experienced symptoms such as wheezing, swelling of the face or tongue, intense itching or cutaneous eruptions during previous treatment with an ACE inhibitor or if a member of your family has already experienced these symptoms, whatever the circumstances (condition called angioedema or Quincke's oedema),
- if you have a serious liver disease or you suffer from a condition called hepatic encephalopathy (severe neurological disorders encountered in severe liver failure),
- if you have a severe kidney disease or if you undergo dialysis,
- if you have abnormally high or low levels of blood potassium,
- in the event of suspicion of non treated cardiac decompensation (considerable water retention, difficulty in breathing),
- if you are over 3 months pregnant (It is also preferable to avoid taking COVERSYL PLUS 4 mg/1.25 mg, tablet in the beginning of the pregnancy, see Pregnancy section)
- if you are breastfeeding.

Take special care with COVERSYL PLUS 4 mg/1.25 mg, tablet:

Inform your doctor before taking COVERSYL PLUS:

- if you have aortic stenosis (narrowing of the main vessel

supplying the heart) or hypertrophic cardiomyopathy (heart muscle disease) or renal artery stenosis (narrowing of the artery bringing blood to the kidney),

- if you have any other heart or kidney problem,
- if you have liver problems,
- if you suffer from a collagen disease (skin disease) such as systemic lupus erythematosus or scleroderma,
- if you suffer from atherosclerosis (hardening of the arteries),
- if you suffer from hyperparathyroidism (dysfunction of the parathyroid gland),
- if you suffer from gout,
- if you are diabetic,
- if you follow a low-salt diet or if you are using a salt substitute containing potassium,
- if you are taking lithium or potassium-sparing diuretics (spironolactone, triamterene) as their use at the same time as COVERSYL PLUS 4 mg/1.25 mg, tablet should be avoided (see "Taking other medicines").

You must inform your doctor if you believe you are (or likely to become) pregnant.

COVERSYL PLUS 4 mg/1.25 mg, tablet is not recommended in the beginning of pregnancy, and must not be taken if you are over 3 months pregnant, as it may seriously harm your child if used from that stage of pregnancy (see section Pregnancy).

Also inform your doctor that you are taking COVERSYL PLUS 4 mg/1.25 mg, tablet:

- if you are to undergo an anaesthetic and/or a surgical procedure,
- if you have recently suffered from diarrhoea and vomiting, or if you are dehydrated,

- if you are to undergo dialysis or apheresis of low-density lipoproteins (LDL) (which consists of removing cholesterol from your blood using a machine),
- if you are to receive desensitisation treatment to reduce the effects of an allergy to bee or wasp stings,
- if you are to undergo a medical examination requiring an injection of an iodinated contrast medium (a substance that renders organs such as the kidney or the stomach visible on X-rays).

The attention of athletes is drawn to the fact that COVERSYL PLUS 4 mg/1.25 mg, tablet contains an active ingredient (indapamide) that may induce a positive reaction during anti-doping control tests.

COVERSYL PLUS 4 mg/1.25 mg, tablet must not be used in children.

Taking other medicines

Tell your doctor or pharmacist if you are taking or have recently taken another medicine, including a medicine obtained without prescription.

You must avoid taking COVERSYL PLUS 4 mg/1.25 mg, tablet with:

- lithium (used to treat depression),
- potassium-sparing diuretics (spironolactone, triamterene), potassium salts,

Coversyl PLUS 4 mg/1.25 mg, tablet treatment can be affected by other medicines. If you are taking one of the following medicines, talk to your doctor, as special precautions may be necessary:

- other medicines to treat hypertension,
- procainamide (for the treatment of irregular heartbeats),
- allopurinol (for the treatment of gout),

- terfenadine or astemizole (antihistamines for hay fever or allergies),
- corticosteroids used to treat various disorders including severe asthma and rheumatoid polyarthritis,
- immunosuppressant agents used for the treatment of autoimmune diseases or following transplant surgery to prevent rejection (e.g. ciclosporine),
- drugs for the treatment of cancer,
- erythromycin by injection (an antibiotic),
- halofantrine (used to treat certain types of malaria),
- pentamidine (used to treat pneumonia),
- gold injection (used to treat rheumatoid polyarthritis),
- vincamine (used to treat symptomatic cognitive disorders in the elderly, in particular memory disorders),
- bepridil (used to treat angina pectoris),
- medicines used for cardiac rhythm problems (for ex.: quinidine, hydroquinidine, disopyramide, amiodarone, sotalol),
- digoxin or other digitalis glycosides (for the treatment of cardiac problems),
- baclofen (to treat muscular stiffness occurring notably in multiple sclerosis),
- medicines to treat diabetes, such as insulin or metformin,
- calcium, including calcium supplements,
- stimulant laxatives (e.g. senna),
- non-steroidal anti-inflammatories (e.g. ibuprofen) or high doses of salicylates (e.g. aspirin),
- amphotericin B by injection (to treat (severe) fungal infection (due to microscopic fungi),
- medicines to treat mental disorders such as depression, anxiety, schizophrenia... (e.g. tricyclic antidepressants, neuroleptics, eg: amisulpride, sulpiride, sultopride, tiapride, haloperidol, droperidol),
- tetracosactide (to treat Crohn's disease).

How to take COVERSYL PLUS 4 mg/1.25 mg, tablet

Always comply with your doctor's prescription. If you are not sure consult your doctor or pharmacist. The usual dose is one tablet once daily. Your doctor may decide to increase the dose to 2 tablets a day or to modify the dose if you suffer from renal insufficiency. The tablet must be swallowed with a glass of water in one daily administration, preferably in the morning and before a meal.

If you have taken more COVERSYL PLUS 4 mg/1.25 mg, tablet than you should have

If you have taken too many tablets, contact your doctor or the nearest hospital immediately.

The most likely event in case of overdose is hypotension. If marked hypotension occurs (associated with dizziness or vertigo), it can be treated by lying the patient down with his/her legs raised.

If you forgot to take COVERSYL PLUS 4 mg/1.25 mg, tablet

It is important to take your medicine every day as directed as a regular treatment is more effective.

However if you forgot to take COVERSYL PLUS 4 mg/1.25 mg, tablet, simply take your treatment the next day as usual.

Do not take a double dose to make up for a forgotten dose.

If you stop taking COVERSYL PLUS 4 mg/1.25 mg, tablet :

Since the treatment of hypertension is generally a long-term treatment, you must ask your doctor for advice before stopping it.

If you have other questions on the use of this medicine, ask your

doctor or pharmacist for more information.

While you are using COVERSYL PLUS 4mg/1.25mg

Food and drinks

It is recommended to take COVERSYL PLUS 4 mg/1.25 mg, tablet before meals.

Pregnancy

You must inform your doctor if you think that you are (or might become) pregnant. Your doctor will normally recommend that you stop taking COVERSYL PLUS 4 mg/1.25 mg, tablet before becoming pregnant or as soon as you find out you are pregnant. He/she will recommend that you take another medicine instead of COVERSYL PLUS 4 mg/1.25 mg, tablet. COVERSYL PLUS 4 mg/1.25 mg, tablet is not recommended in the beginning of pregnancy and it must not be taken if you are over 3 months pregnant, as it may seriously harm your child.

Breastfeeding

You must not take COVERSYL PLUS 4 mg/1.25 mg, tablet if you are breastfeeding.

Inform your doctor if you are breastfeeding or about to start breastfeeding. Consult your doctor immediately.

Ask your doctor or pharmacist for advice before taking any medicine.

Driving vehicles and using machines

Coversyl PLUS 4 mg/1.25 mg, tablet does not usually affect alertness but individual reactions such as dizziness or fatigue related to reduced blood pressure can occur in some patients. As a consequence, your ability to drive vehicles or use machines may be diminished.

Important information concerning certain ingredients

of COVERSYL PLUS 4 mg/1.25 mg, tablet

Coversyl PLUS 4 mg/1.25 mg, tablet contains lactose. If your doctor has informed you that you are intolerant to certain sugars, contact him/her before taking this medicine.

Possible side effect

Like all medicines, COVERSYL PLUS 4 mg/1.25 mg, tablet can cause side effects, though not everybody gets them.

If you experience one of the following side effects, stop taking COVERSYL PLUS 4 mg/1.25 mg, tablet at once and contact your doctor immediately:

- swelling of the face, lips, mouth, tongue or throat, difficulty in breathing,
- severe dizziness or fainting,
- unusually fast or irregular heart beats.

In decreasing order of frequency, side effects may include:

- Common (<1/10, ≥1/100,): headaches, dizziness, vertigo, pins and needles, visual disturbances, tinnitus (sensations of noises in the ears), dizziness due to lowered blood pressure, cough, breathlessness, gastrointestinal disorders (nausea, epigastric pain, anorexia, vomiting, abdominal pain, taste disorders, dry mouth, dyspepsia or difficult digestion, diarrhoea, constipation), allergic reactions (such as skin eruptions, itching), muscle cramps, feeling of tiredness,
- Uncommon (<1/100, ≥1/1000): mood swings, sleep disorders, bronchospasm (feeling of tightness in the chest, short and noisy breath), angioedema (symptoms such as wheezing, swelling of the face and tongue), urticaria, purpura (red dots on the skin) kidney disorders, impotence, sweating,

- Very rare (<1/10 000): confusion, cardiovascular disorders (irregular heart beats, angina pectoris, heart attack), eosinophilic pneumonia (a rare form of pneumonia), rhinitis (blocked or runny nose), severe skin manifestations such as erythema multiforme. If you suffer from systemic lupus erythematosus (a form of collagen disease), this might get worse. Cases of photosensitivity reactions (change in skin appearance) after exposure to the sun or to UVA rays have also been reported. Kidney, liver or pancreas disorders and modifications in laboratory parameters (blood tests) can occur. Your doctor may have to prescribe blood tests to monitor your condition. In the event of liver failure (liver disorders), hepatic encephalopathy may occur (serious neurologic disorders encountered in severe liver failure).

If you notice any side effects not listed in this leaflet, or if some side effects get serious, please inform your doctor or pharmacist.

After using COVERSYL PLUS 4 mg/1.25 mg

Storage

Keep out of the reach and sight of children.

Do not use COVERSYL PLUS 4 mg/1.25 mg, tablet after the expiry date indicated on the box. The expiry date refers to the last day of the month.

Store at a temperature below 30°C.

Disposal

Medicines should not be disposed of via wastewater or household waste. Ask your pharmacist how to dispose of medicines no longer required. These measures will help to protect the environment.

Product description

**What does COVERSYL PLUS
4 mg/1.25 mg, tablet contain**

The active substances are:

Perindopril tert-butylamine and indapamide. Each tablet contains 4 mg of perindopril tert-butylamine (i.e. 3.338 mg of perindopril) and 1.25 mg of indapamide.

The other ingredients are:

Lactose monohydrate, magnesium stearate (E470B), colloidal hydrophobic silica and microcrystalline cellulose.

No 2 Jalan SS 21/37,
Damansara Uptown
47400 Petaling Jaya
Selangor Darul Ehsan

Date of revision: 09.2009

**What does COVERSYL PLUS
4 mg/1.25 mg, tablet looks like
and contents of the outer
packaging**

CoverSYL PLUS 4 mg/1.25 mg, tablet tablets are white and rod-shaped. One tablet contains 4 mg of perindopril tert-butylamine and 1.25 mg of indapamide.

The tablets are available in heat-sealed blisters, overwrapped in a sachet, of 30 tablets.

Registration number

MAL20013354A
(Les Laboratoires Servier Industrie)
MAL09011458ACS
(Kotra Malaysia Sdn Bhd)

Manufacturer

Les Laboratoires Servier Industrie
905 route de Saran
45520 Gidy
France

And

Kotra Malaysia Sdn Bhd
No. 1 Jalan TTC 12
Cheng Industrial Estate
75250 Melaka
Malaysia

Product Licence Holder

Servier Malaysia Sdn Bhd
1301 Level 13, Uptown 2

COVERSYL PLUS 4mg/1.25mg tablet

Perindopril tert-butylamine 4mg/indapamide 1.25mg

Dalam sisip bungkusan ini:

1. Apa itu COVERSYL PLUS 4mg/1.25mg dan kegunaannya
2. Bagaimana COVERSYL PLUS 4mg/1.25mg bertindak
3. Sebelum mengambil COVERSYL PLUS 4mg/1.25mg
4. Cara pengambilan COVERSYL PLUS 4mg/1.25mg
5. Semasa mengambil COVERSYL PLUS 4mg/1.25mg
6. Kesan sampingan yang mungkin berlaku
7. Selepas menggunakan COVERSYL PLUS 4mg/1.25mg
8. Maklumat lanjut

Apa itu COVERSYL PLUS 4mg/1.25mg dan kegunaannya

COVERSYL PLUS 4mg/1.25mg adalah kombinasi dua bahan aktif, perindopril dan indapamide. Ia adalah ubat yang digunakan untuk merawat tekanan darah tinggi (hipertensi).

Bagaimana COVERSYL PLUS 4mg/1.25mg bertindak

Perindopril merupakan ubat didalam kumpulan yang dipanggil (ACE) inhibitor. Ia bertindak dengan membesarakan saluran darah, seterusnya memudahkan jantung anda mengepam darah didalamnya. Indapamide adalah sejenis ubat diuretik. Diuretik meningkatkan kandungan air kencing yang dikeluarkan oleh ginjal. Walau bagaimanapun, indapamide adalah berbeza daripada diuretik lain kerana ia hanya menyebabkan peningkatan kandungan air kencing yang sedikit sahaja. Setiap bahan aktif yang tersebut diatas mengurangkan tekanan darah dan

ianya bertindak bersama-sama untuk mengawal tekanan darah anda.

Sebelum mengambil COVERSYL PLUS 4mg/1.25mg

Jangan mengambil COVERSYL PLUS 4mg/1.25mg

- jika anda alah terhadap perindopril atau ACE inhibitor yang lain atau indapamide atau sulfonamide lain atau sebarang kandungan di dalam COVERSYL PLUS 4mg/1.25mg
- jika anda mengalami simptom seperti kesukaran bernafas, bengkak muka atau lidah, gatal yang teruk atau alahan kulit yang teruk dengan rawatan ACE inhibitor yang sebelumnya atau ahli keluarga anda mempunyai simptom tersebut (keadaan yang dipanggil 'angioedema' atau Quincke's oedema),
- jika anda mempunyai masalah hati atau mempunyai keadaan yang dipanggil 'hepatic encephalopathy' (gangguan neurological yang teruk di dalam kegagalan hati)
- jika anda mempunyai masalah ginjal yang teruk atau memerlukan dialisis,
- jika anda mempunyai kandungan potassium yang terlalu tinggi atau rendah dalam darah,
- jika anda disyaki mempunyai kegagalan jantung yang tidak dirawat (bengkak air yang teruk, kesukaran bernafas),
- jika anda mengandung lebih daripada 3 bulan (elak daripada mengambil COVERSYL PLUS 4mg/1.25mg pada awal kehamilan, lihat Seksyen Mengandung)
- jika anda menyusukan anak.

Ambil perhatian yang khusus dengan COVERSYL PLUS 4mg/1.25mg

Sila berbincang dengan doktor anda jika perkara yang berikut berkaitan dengan anda :

- jika anda mempunyai stenosis aorta (pengucupan salur darah utama yang keluar dari jantung) atau hypertrophic cardiomyopathy (penyakit otot jantung) atau stenosis arteri ginjal (pengucupan salur darah yang menyalurkan darah ke ginjal),
- jika anda ada masalah jantung lain,
- jika anda ada masalah hati,
- jika anda mengalami penyakit kolagen vaskular (penyakit tisu penghubung) seperti systemic lupus erythematosus atau scleroderma,
- jika anda mempunyai arteriosklerosis (pengerasan tisu arteri jantung)
- jika anda mempunyai hyperparathyroidism (kelenjar paratiroid yang tidak berfungsi dengan betul)
- jika anda mempunyai masalah gout,
- jika anda mempunyai penyakit kencing manis,
- jika anda sedang dalam diet garam terkawal atau menggunakan pengganti garam yang mengandungi potassium,
- jika anda mengambil lithium atau diuretik jenis 'potassium-sparing' (spironolactone, triamterene) dan penggunaannya bersama COVERSYL PLUS 4mg/1.25mg harus dielakkan.

Anda mesti memberitahu doktor anda jika anda rasa (atau mungkin) mengandung.

COVERSYL PLUS 4mg/1.25mg tidak disarankan semasa mengandung trimester pertama dan ia boleh menyebabkan kemudaratan yang serius selepas

3 bulan mengandung (lihat Seksyen Mengandung).

Anda patut memberitahu doktor atau pegawai perubatan jika anda mengambil COVERSYL PLUS 4mg/1.25mg:

- jika anda akan melalui proses anaesthesia atau pembedahan,
- jika anda baru mengalami cirit birit atau muntah-muntah atau kehilangan air badan yang berlebihan,
- jika anda melalui proses dialysis atau ‘LDL apheresis’ (pembuangan kolesterol dari saluran darah menggunakan mesin)
- jika anda akan mendapat rawatan untuk menjadikan anda kurang sensitif terhadap kesan alahan lebah atau sengatan tebuian,
- jika anda melalui ujian perubatan yang memerlukan suntikan ‘iodinated contrast agent’ (bahan yang membuatkan organ dalaman seperti ginjal atau perut dapat dilihat melalui X-ray).

Ahli sukan perlu mengetahui bahawa COVERSYL PLUS 4mg/1.25mg mengandungi bahan aktif indapamide yang mungkin memberi kesan positif terhadap ujian darah.

COVERSYL PLUS 4mg/1.25mg tidak disarankan untuk kegunaan kanak-kanak.

Pengambilan ubat lain

Sila beritahu doktor atau ahli farmasi anda jika anda akan atau telah mengambil ubat-ubatan lain termasuk ubat yang tidak memerlukan preskripsi doktor.

Anda patut elak COVERSYL PLUS 4mg/1.25mg dengan:

- litium untuk rawatan depresi,
- diuretik jenis ‘potassium sparing’ (spironolactone, triamterene), garam potassium.

Rawatan dengan COVERSYL PLUS 4mg/1.25mg boleh diganggu oleh ubat-ubatan lain. Sila beritahu doktor jika anda akan atau telah mengambil ubat-ubatan lain yang kerana perhatian yang khusus mungkin diperlukan:

- ubat lain untuk tekanan darah tinggi,
- procainamide (untuk rawatan degupan jantung tidak normal),
- allopurinol (untuk rawatan gout),
- terfenadine atau astemizole (antihistamin untuk alergi atau ‘hay fever’),
- kortikosteroid yang digunakan untuk merawat asma yang teruk dan arthritis reumatoид
- ubat immunosupresant (untuk mengurangkan mekanisma penghalang badan) yang digunakan untuk rawatan penyakit auto imun atau selepas pembedahan transplant (eg ciclosporin),
- ubat untuk rawatan kanser,
- suntikan erythromycin (antibiotik),
- halofantrine (untuk rawatan malaria),
- pentamidine (untuk rawatan paru-paru berair),
- suntikan ‘gold’ (untuk rawatan rheumatoid polyarthritis),
- vincamine (untuk rawatan simptom penyakit kognitif orange dewasa),
- bepridil (untuk rawatan angina pektoris),
- sultopride (untuk rawatan psychoses)
- ubat untuk merawat masalah degupan jantung (eg. quinidine, hydroquinidine, isopyramide, amiodarone, otolol),
- digoxin (untuk rawatan masalah jantung),
- baclofen (untuk rawatan kekejangan otot yang terjadi pada penyakit ‘multiple sclerosis’),
- ubat untuk rawatan kencing manis seperti insulin atau metformin,
- kalsium termasuk suplemen kalsium,

- laksatif perangsang (seperti senna),
- ubat NSAID (seperti ibuprofen) atau dos tinggi salicylates (seperti aspirin),
- amphotericin B melalui suntikan (untuk rawatan penyakit fungus),
- ubat untuk rawatan penyakit mental seperti kemurungan, keresahan, schizophrenia (seperti tricyclic antidepressant, neuroleptics),
- tetracosactide (untuk rawatan Crohn).

Cara pengambilan COVERSYL PLUS 4mg/1.25mg

Selalu mematuhi arahan doktor. Anda perlu hubungi mereka jika anda tidak pasti. Dos biasa adalah 1 tablet sehari. Doktor anda mungkin akan meningkatkan dos ubat kepada 2 tablet sehari atau mengubah pengambilan dos anda jika anda mempunyai kerosakan ginjal. Telan 1 tablet sehari dengan segelas air disebelah pagi dan sebelum makan.

Jika anda mengambil terlebih dos COVERSYL PLUS 4mg/1.25mg daripada yang perlu

Jika anda ambil terlalu banyak tablet, hubungi bahagian kecemasan atau beritahu doktor anda secepat mungkin. Kesan terlebih dos biasanya yang akan berlaku adalah hipotensi (yang dikaitkan rasa pening atau vertigo). Jika ini berlaku, baring dengan menaikkan kaki anda ke atas.

Jika anda terlupa mengambil COVERSYL PLUS 4mg/1.25mg

Adalah penting anda mengambil ubat ini setiap hari seperti yang diarahkan untuk rawatan yang berkesan. Walau bagaimanapun, jika anda terlupa, ambil dos yang berikutnya pada masa yang sama. Jangan menggandakan dos untuk mengganti dos yang terlupa itu.

Jika anda berhenti mengambil COVERSYL PLUS 4mg/1.25mg

Oleh kerana rawatan dengan COVERSYL PLUS 4mg/1.25mg adalah sepanjang hayat, anda perlu berbincang dengan doktor sebelum berhenti mengambil ubat ini.

Jika anda ada sebarang pertanyaan berkaitan penggunaan ubat ini, sila bertanya pada doktor atau ahli farmasi anda.

Semasa mengambil COVERSYL PLUS 4mg/1.25mg

Pengambilan COVERSYL PLUS 4mg/1.25mg dengan makanan dan minuman

Adalah lebih baik COVERSYL PLUS 4mg/1.25mg diambil sebelum makan.

Mengandung

Anda mesti beritahu doktor anda jika anda rasa anda mengandung (atau akan mengandung). Doktor anda akan menasihatkan anda supaya berhenti mengambil COVERSYL PLUS 4mg/1.25mg sebaik saja mengetahui yang anda hamil. Ubat hipertensi yang sesuai mesti digantikan sebelum anda mula hamil. COVERSYL PLUS 4mg/1.25mg tidak disarankan sewaktu awal kehamilan dan boleh menyebabkan kesan mudarat yang serius kepada kandungan jika diambil selepas 3 bulan kehamilan.

Menyusukan anak

Jangan ambil COVERSYL PLUS 4mg/1.25mg jika anda menyusukan bayi.

Sila beritahu doktor anda jika anda menyusukan bayi atau ingin mula menyusukan bayi. Berbincang dengan doktor anda dengan segera.

Sila dapatkan nasihat dari doktor atau ahli farmasi sebelum mengambil sebarang ubat.

Pemanduan dan penggunaan mesin

COVERSYL PLUS 4mg/1.25mg tidak mempunyai kesan ke atas keterjagaan tetapi rasa pening dan lemah yang disebabkan tekanan darah rendah boleh terjadi ke atas sesetengah pesakit. Jika anda mendapat kesan ini, kebolehan anda untuk memandu dan mengendalikan mesin mungkin terjejas.

Informasi penting tentang kandungan COVERSYL PLUS 4mg/1.25mg

Coversyl PLUS 4mg/1.25mg mengandungi laktosa. Jika anda telah diberitahu oleh doktor yang anda mempunyai alahan keatas sesetengah gula, hubungi doktor anda sebelum mengambil ubat ini.

Kesan sampingan yang mungkin berlaku

Seperti ubat-ubatan lain, COVERSYL PLUS 4mg/1.25mg boleh menyebabkan kesan sampingan, walaupun tidak semua pesakit akan merasainya.

Jika anda mengalami sebarang kesan dibawah ini, sila berhenti mengambil ubat dan maklumkan kepada doktor anda secepat mungkin:

- bengkak di bahagian muka, bibir, mulut, lidah atau tekak, kesukaran bernafas
- pening yang teruk atau pengsan
- degupan jantung yang cepat luar biasa atau tidak tetap

Diantara kesan sampingan yang mungkin terjadi (urutan menurun), kesan yang :

- Selalu berlaku (kurang dari 1 dalam setiap 10 pesakit tetapi lebih dari 1 dalam 100 pesakit): sakit kepala, pening, vertigo, pin & needles, gangguan penglihatan, tinnitus (kedengaran bunyi di dalam telinga), kepala rasa pening disebabkan tekanan darah yang rendah, batuk, sesak nafas, gangguan perut-usus (loya, sakit bahagian

perut, pengurangan selera makan, muntah, sakit abdomen, gangguan rasa, mulut kering, dyspepsia atau kesukaran penghadaman, cirit-birit, sembelit), reaksi alergi (seperti masalah kulit, gatal), kejang otot, rasa penat.

- Jarang berlaku (terjadi pada kurang dari 1 dalam 100 pesakit tetapi lebih dari 1 di dalam 1000 pesakit): perubahan emosi, gangguan tidur, bronkospasm (dada rasa ketat, nafas yang pendek dan berbunyi), angioedema (simptom seperti wheezing, bengkak, muka atau lidah), urticaria, purpura (bintik merah pada kulit), masalah ginjal, impoten, berpeluh.
- Sangat jarang berlaku (terjadi pada kurang dari 1 dalam 10,000 pesakit): kecelaruan minda, gangguan kardiovaskular (degupan jantung tidak tetap, angina pectoris, serangan jantung), eosinophilic pneumonia (pneumonia yang sangat jarang berlaku), rhinitis (hidung tersumbat atau selsema), manifestasi kulit yang teruk seperti erythema multiforme. Jika anda mengalami systemic lupus erythematosus (sejenis penyakit kolagen), ia mungkin akan menjadi lebih teruk. Kejadian reaksi photosensitivity (perubahan pada permukaan kulit) selepas terdedah kepada cahaya matahari atau UVA. Gangguan darah, ginjal, hepar atau pankreas dan perubahan parameter makmal (ujian darah) boleh berlaku. Doktor anda mungkin akan melakukan ujian darah untuk memerhatikan keadaan anda. Dalam kes kegagalan hepar (masalah hepar), hepatic encephalopathy mungkin berlaku (gangguan neurologic terjadi dalam kes kegagalan hepar yang teruk).

Jika sebarang kesan sampingan menjadi serius atau anda

mengalami kesan sampingan yang tidak disenaraikan seperti yang diatas, sila beritahu doktor atau ahli farmasi anda.

Selepas menggunakan COVERSYL PLUS 4mg/1.25mg

Bagaimana menyimpan COVERSYL PLUS 4mg/1.25mg

Jauhi dari dicapai atau dilihat oleh kanak kanak.

Jangan mengambil COVERSYL PLUS 4mg/1.25mg selepas tarikh luput yang tercetak pada kotak. Tarikh luput adalah merujuk hari terakhir untuk bulan tersebut.

Simpan di bawah suhu 30°C.

Cara pembuangan

Ubat ini tidak sepatutnya dibuang melalui sistem pembuangan air rumah atau bersama barang buangan lain. Sila bertanya ahli farmasi anda cara-cara membuang ubat yang tidak diperlukan. Langkah ini boleh membantu menjaga persekitaran kita.

Maklumat lanjut

Apa yang terkandung di dalam COVERSYL PLUS 4mg/1.25mg

Bahan aktif:

Perindopril tert-butylamine dan indapamide. Setiap tablet bersalut ini mengandungi 4 mg perindopril tert-butylamine (i.e. 3.338 mg of perindopril) dan 1.25 mg of indapamide.

Bahan lain di dalam tablet:

Lactose monohydrate, magnesium stearate (E470B), colloidal hydrophobic silica, microcrystalline cellulose.

Bagaimana rupa bentuk dan pembungkusan COVERSYL PLUS 4mg/1.25mg

COVERSYL PLUS 4mg/1.25mg adalah tablet yang berwarna putih, tablet bersalut berbentuk rod. Satu tablet bersalut

mengandungi 4mg perindopril tert-butylamine dan 1.25mg indapamide.

Mengandungi 30 tablet yang dipek dalam bentuk blister dan dibungkus di dalam sachet.

No. pendaftaran PBKD:

MAL20013354A
(Les Laboratoires Servier Industrie)
MAL09011458ACS
(Kotra Malaysia Sdn Bhd)

Pengilang:

Les Laboratoires Servier Industrie
905 route de Saran
45520 Gidy - France

Dan

Kotra Malaysia Sdn Bhd
No. 1 Jalan TTC 12
Cheng Industrial Estate
75250 Melaka
Malaysia

Pemegang Pendaftaran:

Servier Malaysia Sdn Bhd
1301 Level 13, Uptown 2
No 2 Jalan SS 21/37,
Damansara Uptown
46400 Petaling Jaya
Selangor

Tarikh revisi: 09.2009