

CONTROLOC 20MG TABLETS

Pantoprazole sodium (20mg)

What is in this leaflet

1. What Controloc is used for
2. How Controloc works
3. Before you use Controloc
4. How to use Controloc
5. While you are using it
6. Side Effects
7. Storage and Disposal of Controloc
8. Product Description
9. Manufacturer and Product Registration Holder
10. Date of Revision

What Controloc is used for

Controloc is used in adults and children 12 years and older:

- Mild reflux disease and associated symptoms (e.g. heartburn, acid regurgitation, pain on swallowing)
- Long-term management and prevention of relapse reflux esophagitis (inflammation or irritation of esophagus caused by a return flow of fluid from stomach)

In adults:

- Prevention of stomach and intestinal ulcer induced by Non-Steroidal Anti-Inflammatory Drugs (NSAIDs)

How Controloc works

Controloc belongs to a medicine group known as proton pump inhibitors (PPIs). Controloc reduces the amount of acid in your stomach. Controloc may help your acid-related symptoms, but you could still have serious stomach problems. Talk with your doctor.

Before you use Controloc

- When you must not use it
You should not take Controloc if you are allergic to pantoprazole or any of the other ingredients of this medicine.

- Before you start to use it

Before you take Controloc, tells your doctor if you:

- have liver problems
- Are pregnant or plan to become pregnant. It is not known if Controloc can harm your unborn baby.
- are breast-feeding or planning to breast-feed. Your doctor should decide if you will take Controloc or breast-feed. You should not do both without first talking with your doctor.
- Have reduced body stores or risk factors for reduced vitamin B12 absorption
- Doctor will plan a few tests if you have any alarm symptom (e.g. significant unintentional weight loss, recurrent vomiting, swallowing difficulty, vomiting blood, low red blood cells or black, tarry stools) and when gastric ulcer is suspected or present, cancer need to be diagnosed.

- Taking other medicines

Tell your doctor about all the medicines you take including prescription and non-prescription medicines, vitamins, and herbal supplements. Controloc may affect how other medicines work, and other medicines may affect how Controloc works. Especially tell your doctor if you take:

- Atazanavir or nelfinavir (medicines for Human Immunodeficiency Virus (HIV))
- Methotrexate (chemotherapy medication used to treat cancer)
- Tacrolimus (prevent transplant rejection),
- Fluvoxamine (for

depression or anxiety disorder)

- Warfarin, phenprocoumon (prevent blood clotting)

How to use Controloc

- How much to use

- Take 1 Controloc 20mg each day for symptom relieve
- Take 1 Controloc 20mg for long-term management and 2 Controloc 20mg if relapse occur.
- One Controloc 20mg for prevention of stomach and intestinal ulcer induced by NSAIDs.

- When to use it

Controloc is taken as advised by your doctor.

- How long to use it

- 2-4-weeks for symptom relief and 4 week treatment for healing of associated esophagitis (inflammation of esophagus). If this is not sufficient, healing will usually be achieved within a further 4 weeks.

- If you forget to use it

If you miss a dose, take it as soon as you remember. If you do not remember until it is time for your next dose, skip the missed dose, and take the next dose at your regular time. Do not take 2 doses of Controloc at the same time.

- If you use too much (overdose)

If you take too much Controloc, call your doctor or go to the nearest hospital emergency room right away.

While you are using it

- Things you must do

Controloc should be swallowed

CONTROLOC 20MG TABLETS

Pantoprazole sodium (20mg)

whole with water.

- Things you must not do
Controloc should not be chewed or crushed.

- Things to be careful of
- There is increased risk for bone fracture for people who received high-doses, or used more than a year or longer.
 - There is an increased risk of bacteria infection.
 - Liver enzymes should be monitored for long-term use.
 - Low magnesium level (hypomagnesemia) has been rarely reported in individuals treated with PPIs for at least three months (in most cases after a year of therapy). Serious consequences of hypomagnesemia include muscle spasm, irregular heart beat, and fits. For those expected to be on prolonged treatment or who take PPIs with medications such as digoxin or medicines that may cause hypomagnesemia (e.g. diuretics), doctor may consider measuring magnesium levels before starting PPI treatment and periodically during treatment.

Side Effects

The uncommon side effects of Controloc include:

- Sleep disorder
- Headache
- Dizziness
- Diarrhoea

- Nausea
- Vomiting
- Constipation
- Dry mouth, abdominal pain and discomfort
- Liver enzymes increased
- Rash
- fatigue

You may report any side effects or adverse drug reactions directly to the National Centre for Adverse Drug Reaction Monitoring by calling Tel: 03-78835550, or visiting the website npra.moh.gov.my (Public → Reporting Medicinal Problems / Side Effects / AEFI / Vaccine Safety).

Storage and Disposal of Controloc

- Storage
Store Controloc below 30°C. Keep Controloc and all medicines out of the reach of children.

- Disposal
Do not throw away any medicines via wastewater or household waste. Ask your pharmacist how to throw away medicines you no longer use. These measures will help protect the environment.

Product Description

- What it looks like
Controloc is supplied as film-coated tablet that are yellow, biconvex and printed with P20 in brown ink on one side.

- Ingredients

- Active ingredient
- Pantoprazole Sodium
- Inactive ingredients
- Sodium carbonate; Mannitol; Crospovidone; Povidone K90; Calcium stearate; Hypromellose 2910; Povidone K25; Titanium dioxide E171;

Yellow ferric oxide E172; Propylene glycol; Triethyl citrate; Methacrylic acid – ethyl acrylate Copolymer (1:1); Polysorbate 80; Sodium Laurisulfate; Printing ink.

- MAL Number
MAL20032151AZ

Manufacturer

Takeda GmbH,
Production Site Oranienburg
Lehnitzstrasse 70-98
D-16515 Oranienburg
Germany

Product Registration Holder

Takeda Malaysia Sdn Bhd
Unit TB-L13-1, Level 13,
Tower B, Plaza 33,
No. 1, Jalan Kemajuan,
Seksyen 13,
46200 Petaling Jaya, Selangor,
Malaysia.

Date of Revision

29/08/2017 (CCDS v3.1)

Serial Number

NPRA(R4/1)290817/00235