

ATEBLOC TABLET 50MG

ATEBLOC TABLET 100MG

Atenolol 50mg / Atenolol 100mg

What is in this leaflet

- What *Atebloc* is used for
- How *Atebloc* work
- Before you use *Atebloc*
- How to take *Atebloc*
- While you are using *Atebloc*
- Side effects
- After using *Atebloc*
- Product description
- Manufacturer
- Marketing authorization holder
- Date of revision

What *Atebloc* is used for

For treatment of high blood pressure, chest pain, irregular heartbeat and preventive treatment of heart attack.

How *Atebloc* work

Atenolol belongs to a group of medicines called beta blockers. It has properties to reduce high blood pressure, relieve chest pains and control irregular heartbeats. Its relief in chest pains is probably due to reduced oxygen demand by heart muscles, its control effect on irregular heartbeats due to blocking stimulation of pacemaker of the heart, and reduction in high blood pressure effect may include reduced output of the heart, decreased sympathetic effect to the peripheral blood vessels and inhibition of enzyme renin release by the kidneys.

Before you use *Atebloc*

When you must not take it

Avoid if there is a heart failure, shock caused by heart disorders, second and third degree heart block, or when heart rate is less than 45 beats/min).

Before you start to take it

Caution if there is history of allergy, mental depression, asthma, heart failure, low blood sugar or renal function impairment.

Effects on ability to drive and use Machines

Atenolol is unlikely to impair a patient's ability to drive or use machinery.

Taking other medications

Concurrent use with general anaesthetics may increase the risk of depression of the heart and low blood pressure. Co-administration of calcium channel blocking agents or clonidine may enhance reduction of blood pressure effects. Concomitant therapy with sympathomimetic amines may result in mutual inhibition of the therapeutic effects.

How to take *Atebloc*

How much to take

Adults: Antianginal: oral, initially, 50mg once daily, the dosage being increased gradually to 100mg after one week if necessary and tolerated. Some patients may require up to 200mg per day. Antihypertensive: Oral, initially 25 to 50mg per day, the dosage being increased to 50 – 100mg a day after two weeks if necessary and tolerated.

Children: Dosage has not been established.

When to take it

Take atenolol exactly as your doctor has told you.

How long to take it

You should check with your doctor or pharmacist.

If you forget to take it

Do not take a double dose to make up for a forgotten tablet.

If you take too much (overdose)

Signs of overdose include slow heartbeat, severe dizziness, fainting, difficulty in breathing, bluish-coloured fingernails or palms of hands, or seizures. For treatment of overdose, consult doctor immediately if symptoms exist.

While you are using *Atebloc*

Things you must do

Inform doctor of medications you are taking. Check with doctor before

discontinuing medication, gradual dosage reduction may be necessary in the treatment of coronary artery disease.

Things you must not do

You must take your medication as directed.

Things to be careful of

If you are pregnant or become pregnant or breast-feeding during treatment, you should not take atenolol unless your doctor tells you to. This is because the risk to your unborn baby is not yet known.

Side effects

Fatigue, depression, cold extremities, skin rash (allergic reactions), diarrhoea, dizziness, headache, nausea or vomiting and weakness may occur occasionally.

After using *Atebloc*

Storage

Store in cool, dry place below 30°C. Protect from light. Keep out of reach of children.

Disposal

Do not dispose via wastewater or household waste. Ask your pharmacist how to dispose of medicines no longer required.

Product description

What it looks like

Atebloc 50: Yellow, round, biconvex, top-scored.

Atebloc 100: Orange, round, flat, top-scored

Ingredients

Atebloc 50: lactose monohydrate, cornstarch, magnesium stearate, dioctyl sodium sulfosuccinate and quinoline yellow lake.

Atebloc 100: lactose monohydrate, cornstarch, magnesium stearate, dioctyl sodium sulfosuccinate and sunset yellow lake.

MAL NO: Atebloc 50: MAL19930047A

Atebloc 100: MAL19930046A

Manufacturer

Prime Pharmaceutical Sdn. Bhd.

1505, Lorong Perusahaan Utama 1

Taman Perindustrian Bukit Tengah

14000 Bukit Mertajam, Penang, Malaysia.

Date of revision

27-10-2011

Marketing Authorization Holder

Prime Pharmaceutical Sdn. Bhd.

Penang, Malaysia.

ATEBLOC TABLET 50MG

ATEBLOC TABLET 100MG

Atenolol 50mg / Atenolol 100mg

Apakah yang ada pada risalah ini

- Apakah kegunaan *Atebloc*
- Bagaimana *Atebloc* berfungsi
- Sebelum mengambil *Atebloc*
- Cara menggunakan *Atebloc*
- Semasa menggunakan *Atebloc*
- Kesan-kesan sampingan
- Cara penyimpanan dan pelupusan *Atebloc*
- Maklumat lanjut
- Pengilang
- Tarikh kemaskini RiMUP

Apakah kegunaan *Atebloc*

Untuk rawatan penyakit darah tinggi, sakit dada, denyutan jantung tak teratur dan rawatan pencegahan serangan jantung.

Bagaimana *Atebloc* berfungsi

Atenolol tergolong dalam kumpulan ubat penghalang beta. Ia mempunyai sifat penurunan tekanan darah, melegakan sakit dada dan mengawal denyutan jantung tak teratur. Kesan melegakan sakit dadanya adalah akibat pengurangan keperluan oksigen otot jantung, kesan mengawal denyutan jantung adalah akibat penghalangan rangsangan ke atas perentak jantung, dan kesan penurunan tekanan darah adalah akibat pengurangan keluaran jantung, pengurangan aliran keluar kesan simpatetik ke saluran darah pinggiran dan rencatan pengeluaran renin oleh ginjal.

Sebelum mengambil *Atebloc*

Bila tidak boleh mengambil
Elakkan pengambilan jika jelas terdapat kegagalan jantung, kejutan jantung tahap 2 atau 3, sekatan jantung atau kadar denyutan jantung kurang daripada 45 denyutan/min.

Sebelum mula mengambil
Berhati-hati jika terdapat kesan alergi, depresi mental, asma, kegagalan jantung, kepekatan gula darah rendah atau ketidaksempurnaan fungsi ginjal. Atenolol sepatutnya tidak mengganggu keupayaan memandu dan mengendalikan mesin.

Tarikh kemaskini RiMUP

27-10-2011

Jika mengambil ubat-ubat lain

Penggunaan serentak dengan anestetik am mungkin meningkatkan risiko depresi otot jantung dan tekanan darah rendah. Penggunaan bersama agen penyekat terusan kalsium atau klonidin mungkin meningkatkan kesan penurunan tekanan darah. Terapi bersama dengan ubat amina simpatomimetik mungkin mengakibatkan rencatan salingan kesan terapeutik.

Cara menggunakan *Atebloc*

Berapa banyak harus diambil
Dewasa: Sakit dada: oral, mulanya 50mg setiap hari dengan peningkatan ansuran kepada 100mg selepas seminggu jika perlu. Ada pesakit yang memerlukan dos sehingga 200mg setiap hari. Tekanan darah tinggi: Oral, mulanya 25 - 50mg setiap hari, dos ditingkatkan kepada 50 – 100mg setiap hari selepas dua minggu, jika perlu. Kanak-kanak: Dos tidak ditetapkan.

Bila perlu diambil

Ambil atenolol mengikut arahan oleh doktor.

Berapa lama perlu diambil

Perlu mengikut nasihat doktor atau ahli farmasi.

Jika terlupa mengambil

Jangan mengambil dos berganda sebagai gantian dos yang terlepas.

Jika mengambil berlebihan(terlebih dos)

Tanda-tanda lebihan dos ialah denyutan jantung perlahan, pening, pengsan, kerumitan bernafas, warna kebiruan kuku atau tapak tangan, atau kekejangan. Perlu dapatkan rawatan segera doktor jika terdapat tandanya terlebih dos.

Semasa menggunakan *Atebloc*

Perkara yang perlu dilakukan

Beritahu doktor ubat yang sedang diambil. Dapatkan nasihat doktor sebelum hentikan pengambilan ubat, pengurangan dos perlu dilakukan secara beransuran dalam rawatan penyakit arteri jantung.

Perkara yang tidak boleh dilakukan

Pengambilan ubat mestilah seperti yang diarahkan.

Perkara yang perlu diberi perhatian

Beritahu doktor jika hamil atau penyusuan semasa rawatan, atenolol tidak boleh diambil tanpa arahan doktor. Ini adalah kerana tiada maklumat mengenai risiko kepada bayi dalam kandungan.

Kesan-kesan sampingan

Letih, depresi, hujung kaki dan tangan yang sejuk, ruam kulit (reaksi alergik), cirit-birit, pening, sakit kepala, loya atau muntah dan kelemahan mungkin berlaku.

Cara penyimpanan dan pelupusan

Atebloc

Penyimpanan

Simpan di tempat yang dingin dan kering, bawah 30°C. Jauhkan dari cahaya. Jauhkan dari kanak-kanak.

Pelupusan

Jangan dibuang dalam air sisa atau sisa buangan. Dapatkan nasihat ahli farmasi cara pelupusan ubat yang ingin dibuang.

Maklumat lanjut

Rupa dan warna produk

Atebloc 50: Tablet kuning, bulat, biconveks dengan permukaan atas skor.

Atebloc 100: Tablet oren, bulat, rata dengan permukaan atas skor.

Bahan-bahan kandungan

Atebloc 50: laktos monohidrat, tepung jagung, magnesium stearat, dioktil sodium sulfosuksinat dan quinolin kuning lake.

Atebloc 100: laktos monohidrat, tepung jagung, magnesium stearat, dioktil sodium sulfosuksinat dan sunset kuning lake.

MAL NO: *Atebloc50: MAL19930047A*
Atebloc100: MAL19930046A

Pengilang

Prime Pharmaceutical Sdn. Bhd.
1505, Lorong Perusahaan Utama 1
Taman Perindustrian Bukit Tengah
14000 Bukit Mertajam, Pulau Pinang,
Malaysia.

Pemegang pendaftaran keluaran

Prime Pharmaceutical Sdn. Bhd.
Pulau Pinang, Malaysia.