

# ROYCE ENEMA

Glycerin and Sodium Chloride (25% w/v / 15% w/v)

## What is in this leaflet

1. What Royce Enema is used for
2. How Royce Enema works
3. Before you use Royce Enema
4. How to take Royce Enema
5. While you are using Royce Enema
6. Side effects
7. Storage and disposal of Royce Enema
8. Product description
9. Manufacturer and Product registration holder
10. Date of revision

## What Royce Enema is used for

Royce Enema is used for the treatment of constipation.

## How Royce Enema works

Glycerin promotes movement of the intestines and evacuation of the bowel by its irritant action.

Sodium chloride is the bile salt involved in maintaining the osmotic pressure of the blood and fluid changes.

## Before you use Royce Enema

### - When you must not use it:

- If you are allergic to glycerin or any of the ingredients in this medicine.
- If you are a children with less than 6 years of age.

Ask your doctor or pharmacist for advice before taking any medicine.

### - Before you start to use it:

Please tell your doctor if you are suffering or have ever suffered from any of the following conditions or illnesses:

- Impairment of renal (kidney) function.
- Calcium channel blockers (prevent calcium into smooth muscle cells), diuretics, or other medicine which may affect electrolyte levels

### - Taking other medicines:

The medicine is for rectal use only. Please tell your doctor or pharmacist if you are taking or have recently taken any other medicines, including medicines obtained without a prescription. Your doctor may need to change the dose of the other medications or take other precautions.

## How to take Royce Enema

### - How much to use:

Your doctor or pharmacist can advise you how many doses are needed each

day and for how long you may need to take it. This depends on your condition and whether or not you are taking any other medicines.

The recommended dosages are as follows:

For rectal use only.

One dose (an ampoule) to be administered each time. If bowel movement does not commence after half an hour, another dose to be administered. Remove the stopper, insert the tip into the rectum and press out the content. Discontinue use if resistance is encountered. Forcing the enema can result in injury.

### - When to use it:

Your doctor will decide when it is appropriate for you to use Royce Enema according to the type of condition you have.

### - How long to use it:

Continue taking Royce Enema for as long as your doctor recommends.

### - If you forget to use it:

Consult your doctor or pharmacist on what you should do if you forget to use it. Take the missed dose as soon as you remember. If it is almost time for your next dose, wait until then to take the medicine and skip the missed dose. Do not take a double dose to make up for the missed dose.

### - If you use too much (overdose):

Contact your doctor immediately or go to the Emergency Department of your nearest hospital, if you think you or anyone else may have taken too much of this medicine. Do this even if there are no signs of discomfort or poisoning. You may need urgent medical attention.

## While you are using Royce Enema

### - Things you must do

- Take your medicine exactly as your doctor has told you.
- Tell all the doctors, dentists and pharmacists treating you that you are taking Royce Enema.
- Tell your doctor immediately if you become pregnant while taking this medication.

### - Things you must not do

- Do not stop taking the medicine unless advised by your doctor.
- Do not take any new medicines without consulting your doctor or pharmacist.
- Do not give Royce Enema to anyone else, even if they have the same symptoms or condition as you.

### - Things to be careful of

- Treatment should be discontinued if severe irritations exist.
- Prolonged use of laxative may result in dependence.
- For use in adults and children above 6 years with intestinal parasitic disease (condition in which a parasite infects the gastro-intestinal tract).

## Side effects

Like all medicines, Royce Enema can cause side effects, although not everybody gets them. Visit your doctor or pharmacist immediately if you experience any of the side effects after taking this medicine:

- Rare gastro (stomach) intestinal irritation.
- Irritation near the rectum (opening for the bowel).

You may report any side effects or adverse drug reactions directly to the National Centre for Adverse Drug Reaction Monitoring by calling Tel:03-78835549 or visiting the website [npra.moh.gov.my](http://npra.moh.gov.my) (Consumer → Reporting Side Effects to Medicines (ConSERF) or Vaccines (AEFI).

## Storage and disposal of Royce Enema

### - Storage:

Store in a dry place below 30°C. Keep out of reach of children.

### - Disposal:

Medicines should not be disposed of via wastewater or household waste. Ask your pharmacist how to dispose the medicines you no longer required. These measures will help to protect the environment.

## Product description

### - What it looks like:

A clear, colourless liquid.

### - Ingredients:

*Active Ingredient:*

Glycerin and sodium chloride.

*Inactive Ingredients:*  
Purified water.

-MAL No.:  
MAL19910319X

**Manufacturer and Product  
Registration Holder**

Royce Pharma Manufacturing Sdn.Bhd.  
PT 1663, Nilai Industrial Estate,  
71800 Nilai, Negeri Sembilan, Malaysia.

**Date of revision**  
18/05/2018

**Serial Number**  
NPRA(R4/1)180518/00161